

THE POSET OF ALL LOGICS II: LEIBNIZ CLASSES AND HIERARCHY

R. JANSANA AND T. MORASCHINI

ABSTRACT. A Leibniz class is a class of logics closed under the formation of term-equivalent logics, compatible expansions, and non-indexed products of sets of logics. We study the complete lattice of all Leibniz classes, called the Leibniz hierarchy. In particular, it is proved that the classes of truth-equational and assertional logics are meet-prime in the Leibniz hierarchy, while the classes of protoalgebraic and equivalential logics are meet-reducible. However, the last two classes are shown to be determined by Leibniz conditions consisting of meet-prime logics only.

1. INTRODUCTION

When ordered under *intepretability* [23], the class of (propositional) logics forms a preorder. Its associated partial order Log , called the *poset of all logics*, consists of equivalence classes of equi-interpretable logics. Building on this formalism, in this paper we introduce and study the notion of a *Leibniz class* of logics.

From an order-theoretic point of view, Leibniz classes are classes of logics that can be faithfully identified with the upsets of Log that are closed under infima of arbitrarily large sets. Equivalently, they can be characterized in terms of closure properties as the classes of logics closed under the formation of term-equivalent logics, compatible expansions, and non-indexed products of sets of logics (see [23] for the relevant definitions).

Part of the interest of Leibniz classes lies in the fact that they allow to clarify the yet informal concept of the *Leibniz hierarchy*, i.e., a taxonomy in which logics are classified in terms of syntactic principles (up to now recognized on empirical grounds) corresponding to the behaviour of the so-called Leibniz operator [12, 18, 19, 20]. More precisely, the road we take is to identify the Leibniz hierarchy with the complete lattice of Leibniz classes ordered under inclusion. Remarkably, this abstraction preserves the fact that Leibniz classes are collections of logics globally satisfying some syntactic principles, here called *Leibniz conditions*, consisting of special sequences of logics indexed by all ordinals.

One of the main advantages of this point of view is that it allowed to unify in [24] the theory of the Leibniz hierarchy with that of the *Maltsev hierarchy* of universal algebra, i.e., a taxonomy in which varieties are classified by means of syntactic principles related to the structure of congruence lattices [22, 25, 29, 31, 36, 37, 39].

In this paper we restrict our attention to the study of Leibniz classes. First we show that the majority of classes of logics traditionally associated with the Leibniz hierarchy happen to be Leibniz classes. Among them we count the classes of protoalgebraic, equivalential, and assertional logics, whose respective Leibniz conditions are also found (Theorems 3.4,

2010 *Mathematics Subject Classification.* 03G27,03C05,08B05.

Key words and phrases. Abstract algebraic logic, Leibniz hierarchy, Leibniz condition, Maltsev condition, interpretability, poset of all logics.

3.7, and 3.11). Then we begin an order-theoretic investigation of the Leibniz hierarchy, understood as the complete lattice of all Leibniz classes. More precisely, we focus on the problem of determining whether the most prominent Leibniz classes are meet-prime or meet-irreducible elements of the Leibniz hierarchy. Affirmative answers to these questions can then be interpreted as stating that the Leibniz classes under consideration capture primitive or fundamental concepts. Similar problems were studied in the setting of the Maltsev hierarchy for instance in [21] (see also [2, 30, 38]).

Some of our results can be summarized as follows. On the one hand the Leibniz classes of truth-equational and assertional logics are shown to be meet-prime (Theorems 5.10 and 5.11). On the other hand, it is proved that the classes of protoalgebraic and equivalential logics are meet-reducible (Theorems 6.7 and 6.7). This negative result is compensated by the observation that the Leibniz condition determining the class of protoalgebraic (resp. equivalential) logics consists of logics whose equivalence classes are meet-prime in the poset Log (Theorems 7.2 and 7.5).

2. LEIBNIZ CONDITIONS

We use the same notation as in [23]. Recall that OR is the class of ordinals.

Definition 2.1. A *strong Leibniz condition* Φ is simply a logic \vdash_Φ . A logic \vdash is said to *satisfy* Φ if $\vdash_\Phi \leq \vdash$, and the class of logics satisfying Φ is denoted by $\text{Log}(\Phi)$. Similarly, a *Leibniz condition* Φ is a class $\{\Phi_\alpha : \alpha \in \text{OR}\}$ of strong Leibniz conditions such that if $\alpha \leq \beta$, then $\vdash_{\Phi_\beta} \leq \vdash_{\Phi_\alpha}$. A logic \vdash is said to *satisfy* Φ if $\vdash_{\Phi_\alpha} \leq \vdash$ for some $\alpha \in \text{OR}$, and the class of logics satisfying Φ is denoted by $\text{Log}(\Phi)$.

Accordingly, a class \mathbb{K} of logics is a (resp. *strong*) *Leibniz class* if it is of the form $\text{Log}(\Phi)$ for some (resp. strong) Leibniz condition Φ . \square

Recall that Log is a set-complete meet-semilattice [23, Thm. 4.6]. A subcollection $F \subseteq \text{Log}$ is a *set-complete filter* if it is a non-empty upset closed under infima of sets. Similarly, $F \subseteq \text{Log}$ is a *principal filter* if it is a principal upset of Log . Finally, given a class \mathbb{K} of logics, we set

$$\mathbb{K}^\dagger := \{\llbracket \vdash \rrbracket : \vdash \in \mathbb{K}\}.$$

The following result is instrumental to construct concrete Leibniz classes.

Theorem 2.2. *Let \mathbb{K} be a class of logics. The following conditions are equivalent:*

- (i) \mathbb{K} is a Leibniz class.
- (ii) \mathbb{K} is closed under term-equivalence, compatible expansions, and non-indexed products of sets of logics.
- (iii) The collection \mathbb{K}^\dagger is a set-complete filter of Log , and $\mathbb{K} = \{\vdash : \llbracket \vdash \rrbracket \in \mathbb{K}^\dagger\}$.

Proof. We rely on the observation that, given a family $\{\vdash_i : i \in I\}$ of logics, the infimum of $\{\llbracket \vdash_i \rrbracket : i \in I\}$ in Log is $\llbracket \otimes_{i \in I} \vdash_i \rrbracket$ [23, Thm. 4.6].

(i) \Rightarrow (iii): Suppose that \mathbb{K} is a Leibniz class, i.e., that there is a Leibniz condition $\Phi = \{\Phi_\alpha : \alpha \in \text{OR}\}$ such that $\mathbb{K} = \text{Log}(\Phi)$. We begin by proving that the collection \mathbb{K}^\dagger is a set-complete filter of Log .

It is clear that \mathbb{K}^\dagger is a non-empty upset of Log . Then consider a set of logics $\{\vdash_i : i \in I\} \subseteq \mathbb{K}$. Since $\mathbb{K} = \text{Log}(\Phi)$, for every $i \in I$ there exists an ordinal α_i such that $\vdash_{\Phi_{\alpha_i}} \leq \vdash_i$. Let β be the supremum of $\{\alpha_i : i \in I\}$. Since Φ is a Leibniz condition, we have $\vdash_{\Phi_\beta} \leq \vdash_{\Phi_{\alpha_i}} \leq \vdash_i$ for every $i \in I$. Hence we obtain $\vdash_{\Phi_\beta} \leq \otimes_{i \in I} \vdash_i$ which, together with $\mathbb{K} = \text{Log}(\Phi)$, implies

that $\bigotimes_{i \in I} \vdash_i \in \mathbb{K}$. Since $\llbracket \bigotimes_{i \in I} \vdash_i \rrbracket$ is the infimum of $\{\llbracket \vdash_i \rrbracket : i \in I\}$ in Log , we conclude that \mathbb{K}^\dagger is closed under infima of sets. This establishes that \mathbb{K}^\dagger is a set-complete filter of Log .

Now, from the definition of \mathbb{K}^\dagger it follows that $\mathbb{K} \subseteq \{\vdash : \llbracket \vdash \rrbracket \in \mathbb{K}^\dagger\}$. To prove the other inclusion, consider a logic \vdash such that $\llbracket \vdash \rrbracket \in \mathbb{K}^\dagger$. By the definition of \mathbb{K}^\dagger , there is a logic $\vdash' \in \mathbb{K}$ such that $\vdash' \leq \vdash$. Since $\vdash' \in \mathbb{K}$, there is an ordinal α such that $\vdash_{\Phi_\alpha} \leq \vdash' \leq \vdash$. Since $\mathbb{K} = \text{Log}(\Phi)$, this implies that $\vdash \in \mathbb{K}$ as desired.

(iii) \Rightarrow (ii): Suppose that $\vdash \in \mathbb{K}$, and consider a logic \vdash' that is either term-equivalent to \vdash or a compatible expansion of \vdash . By [23, Prop. 3.8] we have that $\vdash \leq \vdash'$ and, therefore, $\llbracket \vdash \rrbracket \leq \llbracket \vdash' \rrbracket$. Since \mathbb{K}^\dagger is an upset of Log and $\llbracket \vdash \rrbracket \in \mathbb{K}^\dagger$, we obtain that $\llbracket \vdash' \rrbracket \in \mathbb{K}^\dagger$. Together with the fact that $\mathbb{K} = \{\vdash'' : \llbracket \vdash'' \rrbracket \in \mathbb{K}^\dagger\}$, this yields $\vdash' \in \mathbb{K}$. Hence we conclude that \mathbb{K} is closed under term-equivalence and compatible expansions.

Then consider a family $\{\vdash_i : i \in I\} \subseteq \mathbb{K}$. By assumption the infimum of $\{\llbracket \vdash_i \rrbracket : i \in I\}$ in Log belongs to \mathbb{K}^\dagger . This amounts to the fact that $\llbracket \bigotimes_{i \in I} \vdash_i \rrbracket \in \mathbb{K}^\dagger$. As $\mathbb{K} = \{\vdash : \llbracket \vdash \rrbracket \in \mathbb{K}^\dagger\}$, we conclude that $\bigotimes_{i \in I} \vdash_i \in \mathbb{K}$.

(ii) \Rightarrow (i): Consider the cumulative hierarchy $\{V_\alpha : \alpha \in \text{OR}\}$ of set theory. For every ordinal α we set

$$\mathbb{K}_\alpha := \mathbb{K} \cap V_\alpha \text{ and } \vdash_\alpha := \bigotimes \mathbb{K}_\alpha.$$

Note that if $\mathbb{K}_\alpha = \emptyset$, then \vdash_α is the inconsistent logic in the empty language over ω variables. Also note that if $\alpha \leq \beta$, then $\mathbb{K}_\alpha \subseteq \mathbb{K}_\beta$ and, therefore, $\vdash_\beta \leq \vdash_\alpha$. In particular, this implies that the following is a Leibniz condition:

$$\Phi := \{\vdash_\alpha : \alpha \in \text{OR}\}.$$

To conclude the proof, it suffices to show that $\text{Log}(\Phi) = \mathbb{K}$. To prove the right-to-left inclusion, consider a logic $\vdash \in \mathbb{K}$. Since \vdash is a set, there is an ordinal α such that $\vdash \in V_\alpha$. This implies that $\vdash \in \mathbb{K}_\alpha$ and, therefore, that $\vdash_\alpha \leq \vdash$. Hence we conclude that $\vdash \in \text{Log}(\Phi)$.

To prove the other inclusion, consider $\vdash \in \text{Log}(\Phi)$. There exists an ordinal α such that $\vdash_\alpha \leq \vdash$. By [23, Prop. 3.8] this implies that \vdash is term-equivalent to a compatible expansion of $\bigotimes \mathbb{K}_\alpha$. As $\mathbb{K}_\alpha \subseteq \mathbb{K}$ and \mathbb{K} is closed under non-indexed products of sets of logics, compatible expansions, and term-equivalence, this implies $\vdash \in \mathbb{K}$. \square

Corollary 2.3. *A class \mathbb{K} of logics is a strong Leibniz class if and only if \mathbb{K}^\dagger is a principal filter of Log , and $\mathbb{K} = \{\vdash : \llbracket \vdash \rrbracket \in \mathbb{K}^\dagger\}$.*

Proof sketch. The proof of the “only if” part is an easier variant of the one of part (i) \Rightarrow (iii) of Theorem 2.2. Then we sketch the “if” part only. By the assumption we know that \mathbb{K}^\dagger is the upset generated by $\llbracket \vdash \rrbracket$, for some logic $\vdash \in \mathbb{K}$. Let Φ be the strong Leibniz condition determined by \vdash . It is not hard to see that $\text{Log}(\Phi) = \mathbb{K}$ and, therefore, that \mathbb{K} is a strong Leibniz class. \square

Remark 2.4. Typical applications of Theorem 2.2 comprise proofs that certain collections of logics are Leibniz classes. For instance, in [23, Props. 3.8 and 6.1] it was shown that the class of equivalential logics is closed under term-equivalence, compatible expansions, and non-indexed products of sets of logics. By Theorem 2.2(ii) we conclude that it is a Leibniz class. \square

Remark 2.5. From Theorem 2.2(ii) it follows that Leibniz classes are closed under the formation of extensions of logics, as these are special cases of compatible expansions. \square

In this paper we identify the intuitive concept of the *Leibniz hierarchy* with the poset of all Leibniz classes ordered under the inclusion relation.

Proposition 2.6. *The Leibniz hierarchy is a complete lattice in which infima are intersections and suprema are obtained as follows for every collection $\{\mathbb{K}_i : i \in I\}$ of Leibniz classes, where I can be a proper class:*

$$\bigvee_{i \in I} \mathbb{K}_i = \{\vdash : \vdash \text{ is a logic and } \bigotimes_{j \in J} \vdash_j \leq \vdash \text{ for some subset } \{\vdash_j : j \in J\} \subseteq \bigcup_{i \in I} \mathbb{K}_i\}.$$

Proof. Immediate from Theorem 2.2. □

Remark 2.7. The statement of Proposition 2.6 presupposes that we can meaningfully speak of very large intersections and unions of classes (of logics), as the collection $\{\mathbb{K}_i : i \in I\}$ is in general a collection of classes. However, for our purposes this problem is immaterial as we will only work with finite joins and meets of Leibniz classes. □

3. EXAMPLES OF LEIBNIZ CLASSES

In this section we show that a range of classes of logics, traditionally associated with the yet informal concept of the Leibniz hierarchy in abstract algebraic logic, are indeed Leibniz classes.

A logic \vdash is said to be *protoalgebraic* [7, 12] if there is a non-empty set¹ $\Delta(x, y, \vec{z})$ of formulas such that for every $\langle A, F \rangle \in \text{Mod}(\vdash)$ and $a, b \in A$,

$$\langle a, b \rangle \in \Omega^A F \iff \Delta^A(a, b, \vec{c}) \subseteq F, \text{ for every } \vec{c} \in A.$$

In this case, we say that $\Delta(x, y, \vec{z})$ is a set of *congruence formulas with parameters* for \vdash .

Theorem 3.1. *Let \vdash be a logic. The following conditions are equivalent:*

- (i) \vdash is protoalgebraic.
- (ii) \vdash has theorems and $\text{Mod}^\equiv(\vdash) = \mathbb{R}(\text{Mod}(\vdash))$.
- (iii) There is a non-empty set of formulas $\nabla(x, y)$ such that

$$\emptyset \vdash \nabla(x, x) \quad x, \nabla(x, y) \vdash y.$$

In this case, the following is a set congruence formulas with parameters for \vdash :

$$\hat{\nabla}(x, y, \vec{z}) := \{\varphi(\psi(x, \vec{z}), \psi(y, \vec{z})) : \varphi(x, y) \in \nabla \text{ and } \psi(x, \vec{z}) \in \text{Fm}_{\mathcal{L}_\vdash}(\omega)\}.$$

Proof. For the equivalence between (i), (ii), and (iii), see [18, Thms. 6.7, 6.17, and 6.57]. The fact that $\hat{\nabla}$ is a set of equivalence formulas with parameters for \vdash follows from [9, Thm. 13.5] (see also [16, Prop. 3.2]). □

Our aim is to prove that protoalgebraic logics form a Leibniz class. To this end, it is convenient to introduce the following concept:

Definition 3.2. Given an infinite cardinal κ , let $\mathcal{L}_\vdash^\kappa$ be the language consisting of the binary symbols $\{\neg_\alpha : \alpha < \kappa\}$ and the n -ary symbols $\{\ast_{n\alpha} : \alpha < \kappa\}$ for $0 < n \in \omega$. We set

$$\nabla_\kappa(x, y) := \{x \neg_\alpha y : \alpha < \kappa\}.$$

¹In the literature the set Δ is not required to be non-empty. However, this restriction is almost immaterial as, in a fixed language, there is a unique protoalgebraic logic with an empty Δ , namely the almost inconsistent logic [18, Prop. 6.11.5].

The *basic protoalgebraic logic of rank κ* is the logic $\vdash_{\mathfrak{P}}^{\kappa}$ formulated on $Fm_{\mathcal{L}_+^{\kappa}}(\omega)$ determined by the rules

$$\emptyset \triangleright \nabla_{\kappa}(x, x) \quad x, \nabla_{\kappa}(x, y) \triangleright y. \quad \boxtimes$$

The following result explains the role of $\vdash_{\mathfrak{P}}^{\kappa}$.

Proposition 3.3. *Let \vdash be a logic.*

- (i) *\vdash is protoalgebraic if and only if $\vdash_{\mathfrak{P}}^{\kappa} \leq \vdash$ for every (equiv. some) infinite cardinal $\kappa \geq |\mathcal{L}_+|$.*
- (ii) *If κ is an infinite cardinal and $\vdash_{\mathfrak{P}}^{\kappa} \leq \vdash$, then \vdash is protoalgebraic.*

Proof. (ii): Let τ be an interpretation of $\vdash_{\mathfrak{P}}^{\kappa}$ into \vdash . From [23, Prop. 3.3] we obtain $\emptyset \vdash \tau[\nabla_{\kappa}(x, x)]$ and $x, \tau[\nabla_{\kappa}(x, y)] \vdash y$. Then the set $\tau[\nabla_{\kappa}(x, y)]$ witnesses the validity of condition (iii) of Theorem 3.1, whence \vdash is protoalgebraic.

(i): As a special instance of (ii) we obtain that if $\vdash_{\mathfrak{P}}^{\kappa} \leq \vdash$ for some infinite cardinal $\kappa \geq |\mathcal{L}_+|$, then \vdash is protoalgebraic. Then suppose that \vdash is protoalgebraic, and consider any infinite cardinal $\kappa \geq |\mathcal{L}_+|$. By Theorem 3.1 there is a set of formulas $\nabla(x, y) \subseteq Fm(\vdash)$ such that

$$\emptyset \vdash \nabla(x, x) \text{ and } x, \nabla(x, y) \vdash y. \quad (1)$$

Now, observe that $|\nabla| \leq \max\{\omega, |\mathcal{L}_+|\} \leq \kappa$. Since $\nabla \neq \emptyset$, there is a surjective map

$$f: \{-\circ_{\alpha}: \alpha < \kappa\} \rightarrow \nabla(x_1, x_2).$$

Similarly, since $|Fm_{\mathcal{L}_+}(\omega)| \leq \max\{\omega, |\mathcal{L}_+|\} \leq \kappa$, for every $0 < n \in \omega$ there is a surjective map

$$g_n: \{*_n\alpha: \alpha < \kappa\} \rightarrow \{\varphi \in Fm_{\mathcal{L}_+}(\omega): \varphi = \varphi(x_1, \dots, x_n)\}.$$

Observe that the maps f and $\{g_n: 0 < n \in \omega\}$ can be turned in the natural way into a single translation τ of $\mathcal{L}_{\mathfrak{P}}^{\kappa}$ into \mathcal{L}_+ .

Recall by Theorem 3.1 that $\hat{\nabla}_{\kappa}$ and $\hat{\nabla}$ are sets of congruence formulas with parameters for $\vdash_{\mathfrak{P}}^{\kappa}$ and \vdash respectively. We claim that

$$\hat{\nabla}(x, y, \vec{z}) = \tau[\hat{\nabla}_{\kappa}(x, y, \vec{z})]. \quad (2)$$

We begin by proving the inclusion from left to right. Consider $\varphi \in \hat{\nabla}$. There are $\psi(x, y) \in \nabla(x, y)$ and $\gamma(x, z_1, \dots, z_{n-1}) \in Fm_{\mathcal{L}_+}(\omega)$ such that

$$\varphi = \psi(\gamma(x, z_1, \dots, z_{n-1}), \gamma(y, z_1, \dots, z_{n-1})).$$

Since f and g_n are surjective, there are $\alpha, \beta < \kappa$ such that $\tau(-\circ_{\alpha}) = \psi(x_1, x_2)$ and $\tau(*_n\beta) = \gamma(x_1, \dots, x_n)$. Clearly,

$$\varphi = \tau(*_n\beta(x, z_1, \dots, z_{n-1}) -\circ_{\alpha} *_n\beta(y, z_1, \dots, z_{n-1})).$$

Moreover, the definition of $\hat{\nabla}_{\kappa}$ guarantees

$$*_n\beta(x, z_1, \dots, z_{n-1}) -\circ_{\alpha} *_n\beta(y, z_1, \dots, z_{n-1}) \in \hat{\nabla}_{\kappa}.$$

Hence we conclude that $\varphi \in \tau[\hat{\nabla}_{\kappa}]$. This establishes the inclusion from left to right in (2).

To prove the other inclusion, consider $\varphi \in \tau[\hat{\nabla}_{\kappa}]$. Then there are $\alpha < \kappa$ and $\gamma(x, \vec{z}) \in Fm(\vdash_{\mathfrak{P}}^{\kappa})$ such that $\varphi = \tau(\gamma(x, \vec{z}) -\circ_{\alpha} \gamma(y, \vec{z}))$. Set

$$\gamma'(x, \vec{z}) := \tau(\gamma(x, \vec{z})) \text{ and } \psi(x, y) := \tau(x -\circ_{\alpha} y).$$

From the definition of $\hat{\nabla}$ and the fact that $f(-\circ_{\alpha}) \in \nabla(x_1, x_2)$, it follows

$$\varphi = \tau(\gamma(x, \vec{z}) -\circ_{\alpha} \gamma(y, \vec{z})) = \psi(\gamma'(x, \vec{z}), \gamma'(y, \vec{z})) \in \hat{\nabla}.$$

This establishes the equality in (2).

To conclude the proof, it suffices to show that τ is an interpretation of $\vdash_{\mathbb{P}}^{\kappa}$ into \vdash . To this end, consider a matrix $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$. We begin by showing that $\langle A^{\tau}, F \rangle$ is a model of $\vdash_{\mathbb{P}}^{\kappa}$. Observe that $\nabla(x, y) = \tau[\nabla_{\kappa}(x, y)]$. Together with (1) and $\langle A, F \rangle \in \text{Mod}(\vdash)$, this yields that the matrix $\langle A, F \rangle$ is a model of the rules $\emptyset \triangleright \tau[\nabla_{\kappa}(x, x)]$ and $x, \tau[\nabla_{\kappa}(x, y)] \triangleright y$. As a consequence, $\langle A^{\tau}, F \rangle$ is a model of the rules $\emptyset \triangleright \nabla_{\kappa}(x, x)$ and $x, \nabla_{\kappa}(x, y) \triangleright y$, whence it is a model of $\vdash_{\mathbb{P}}^{\kappa}$.

Now, for every $a, b \in A$,

$$\begin{aligned} a = b &\iff \langle a, b \rangle \in \Omega^A F \\ &\iff \hat{\nabla}^A(a, b, \vec{c}) \subseteq F \text{ for every } \vec{c} \in A \\ &\iff \tau[\hat{\nabla}_{\kappa}]^A(a, b, \vec{c}) \subseteq F \text{ for every } \vec{c} \in A \\ &\iff \hat{\nabla}_{\kappa}^{A^{\tau}}(a, b, \vec{c}) \subseteq F \text{ for every } \vec{c} \in A \\ &\iff \langle a, b \rangle \in \Omega^{A^{\tau}} F. \end{aligned}$$

The above equivalences are justified as follows: the first is a consequence of the fact that $\langle A, F \rangle$ is reduced by Theorem 3.1(ii), the second follows from the fact that $\langle A, F \rangle \in \text{Mod}(\vdash)$ and that $\hat{\nabla}$ is a set of congruence formulas with parameters for \vdash , the third from (2), the fourth is straightforward, and the fifth from the fact that $\langle A^{\tau}, F \rangle \in \text{Mod}(\vdash_{\mathbb{P}}^{\kappa})$ and that $\hat{\nabla}_{\kappa}$ is a set of congruence formulas with parameters for $\vdash_{\mathbb{P}}^{\kappa}$. The above display implies that the congruence $\Omega^{A^{\tau}} F$ is the identity relation. As a consequence, we obtain $\langle A, F \rangle \in \mathbb{R}(\text{Mod}(\vdash_{\mathbb{P}}^{\kappa})) = \text{Mod}^{\equiv}(\vdash_{\mathbb{P}}^{\kappa})$. This establishes that τ is an interpretation of $\vdash_{\mathbb{P}}^{\kappa}$ into \vdash , as desired. \square

For every ordinal α , let $\vdash_{\mathbb{P}}^{\alpha}$ be the logic $\vdash_{\mathbb{P}}^{\omega+|\alpha|}$.

Theorem 3.4. *The sequence $\Phi = \{\vdash_{\mathbb{P}}^{\alpha} : \alpha \in \text{OR}\}$ is a Leibniz condition and $\text{Log}(\Phi)$ coincides with the class of protoalgebraic logics. In particular, protoalgebraic logics form a Leibniz class.*

Proof. To prove that Φ is a Leibniz condition, consider $\alpha, \beta \in \text{OR}$ such that $\alpha \leq \beta$. The logic $\vdash_{\mathbb{P}}^{\alpha}$ is protoalgebraic by Proposition 3.6(ii). This fact and

$$|\mathcal{L}_{\alpha}| = \omega + |\alpha| \leq \omega + |\beta|$$

allow us to apply Proposition 3.6(i), obtaining $\vdash_{\mathbb{P}}^{\beta} \leq \vdash_{\mathbb{P}}^{\alpha}$. Hence we conclude that Φ is a Leibniz condition. Finally, the fact that $\text{Log}(\Phi)$ is the class of protoalgebraic logics is a direct consequence of Proposition 3.6. \square

By Remark 2.4 we know that the collection Equiv of equivalential logics is a Leibniz class. It is therefore sensible to wonder whether we can find an intelligible Leibniz condition Φ such that $\text{Equiv} = \text{Log}(\Phi)$. This can be done with a simple adaptation of the method employed in the case of protoalgebraic logics.

Definition 3.5. Given an infinite cardinal κ , let $\mathcal{L}_{\mathbb{E}}^{\kappa}$ be the language consisting of the binary symbols $\{\neg_{\alpha} : \alpha < \kappa\}$. We set

$$\Delta_{\kappa}(x, y) := \{x \neg_{\alpha} y : \alpha < \kappa\}.$$

The *basic equivalential logic of rank κ* is the logic $\vdash_{\mathbb{E}}^{\kappa}$ formulated on $\text{Fm}_{\mathcal{L}_{\mathbb{E}}^{\kappa}}(\omega)$ determined by the following rules, stipulated for every $\alpha < \kappa$,

$$\emptyset \triangleright \Delta_{\kappa}(x, x) \quad x, \Delta_{\kappa}(x, y) \triangleright y$$

$$\Delta_\kappa(x_1, y_1) \cup \Delta_\kappa(x_2, y_2) \triangleright \Delta_\kappa(x_1 \multimap_\alpha x_2, y_1 \multimap_\alpha y_2). \quad \boxtimes$$

The importance of the logic \vdash_E^κ is justified as follows:

Proposition 3.6. *Let \vdash be a logic.*

- (i) \vdash is equivalential if and only if $\vdash_E^\kappa \leq \vdash$ for every (equiv. some) infinite cardinal $\kappa \geq |\mathcal{L}_\vdash|$.
- (ii) If κ is an infinite cardinal and $\vdash_E^\kappa \leq \vdash$, then \vdash is equivalential.

Proof. Condition (ii) follows from the fact that equivalential logics form a Leibniz class by Remark 2.4, while the proof of (i) is analogous to that of Proposition 3.6(i). \boxtimes

For every ordinal α , let \vdash_E^α be the logic $\vdash_E^{\omega+|\alpha|}$.

Theorem 3.7. *The sequence $\Phi = \{\vdash_E^\alpha : \alpha \in \text{OR}\}$ is a Leibniz condition and $\text{Log}(\Phi)$ coincides with the class of equivalential logics. In particular, equivalential logics form a Leibniz class.*

Proof. Analogous to the one of Theorem 3.4. \boxtimes

A logic \vdash is said to be *assertional* [1, 32] if F is a singleton for every $\langle A, F \rangle \in \text{Mod}^\equiv(\vdash)$.

Proposition 3.8. *Let \vdash be a logic.*

- (i) If $x, y, \varphi(x, \vec{z}) \vdash \varphi(y, \vec{z})$ for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash)$, then $|F| \leq 1$ for every $\langle A, F \rangle \in \text{Mod}^\equiv(\vdash)$.
- (ii) \vdash is assertional if and only if it has theorems and $x, y, \varphi(x, \vec{z}) \vdash \varphi(y, \vec{z})$ for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash)$.

Proof. See [1, Thm. 10]. The result is attributed to Suszko in [11], see also [34]. \boxtimes

To prove that assertional logics form a *strong* Leibniz class, we need few more concepts.

Definition 3.9. The *basic assertional logic* is the logic \vdash_A formulated in countably many variables and in the language comprising just a unary connective $\top(x)$, axiomatized by the rule $\emptyset \triangleright \top(x)$.

A *pointed set* is an algebra $A = \langle A; \top^A \rangle$ where \top^A is a unary constant map on A . In this case, we denote by \top_*^A the element of A defined by the map $\top^A: A \rightarrow A$.

Proposition 3.10. *The logic \vdash_A is assertional and*

$$\text{Mod}^\equiv(\vdash_A) = \{\langle A, \{\top_*^A\} \rangle : A \text{ is a pointed set}\}.$$

Proof. Due to the poor signature of \vdash_A and the fact that $\emptyset \vdash_A \top(x)$, it is easy to see that $\emptyset \vdash_A \varphi$ for every formula $\varphi \in \text{Fm}(\vdash_A)$ that is not a variable. As a consequence, we obtain $x, y, \varphi(x, \vec{z}) \vdash \varphi(y, \vec{z})$ for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash_A)$. Moreover, the logic \vdash_A has theorems, e.g., $\top(x)$. Hence, we can apply Proposition 3.8(ii) obtaining that \vdash_A is assertional.

Now we turn to prove the equality in the statement. First consider a matrix $\langle A, F \rangle \in \text{Mod}^\equiv(\vdash_A)$. Since \vdash_A is assertional, $F = \{a\}$ for some $a \in A$. Together with the fact that $\top(x)$ is a theorem, this yields $\top^A(c) = a$ for all $c \in A$. Hence \top^A is a constant map on A and $a = \top_*^A$. This establishes the inclusion from left to right.

To prove the other inclusion, let A be a pointed set. It is clear that the matrix $\langle A, \{\top_*^A\} \rangle$ is a model of the rule $\emptyset \triangleright \top(x)$ and, therefore, of \vdash_A . It only remains to prove that the congruence $\tilde{\Omega}_{\vdash_A}^A \{\top_*^A\}$ is the identity relation. To this end, consider two distinct elements $a, c \in A$. We can assume without loss of generality that $c \neq \top_*^A$. Then, by the compatibility

of the Leibniz congruence, we obtain $\langle a, c \rangle \notin \Omega^A\{\top_*^A, a\}$. Moreover, observe that the matrix $\langle A, \{\top_*^A, a\} \rangle$ is a model of the rule $\emptyset \triangleright \top(x)$ and, therefore, of the logic \vdash_A , whence $\tilde{\Omega}_{\vdash_A}^A\{\top_*^A\} \subseteq \Omega^A\{\top_*^A, a\}$. Hence we conclude that $\langle a, c \rangle \notin \tilde{\Omega}_{\vdash_A}^A\{\top_*^A\}$. This shows that $\tilde{\Omega}_{\vdash_A}^A\{\top_*^A\}$ is the identity relation and, therefore, $\langle A, \{\top_*^A\} \rangle \in \text{Mod}^{\equiv}(\vdash_A)$. \square

As a consequence, we obtain the following:

Theorem 3.11. *Assertional logics form the strong Leibniz class $\text{Log}(\vdash_A)$.*

Proof. It suffices to show that a logic \vdash is assertional if and only if $\vdash_A \leq \vdash$. To prove the “if” part, suppose that $\vdash_A \leq \vdash$ and let τ be an interpretation of \vdash_A into \vdash . Then consider $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$. We have that $\langle A^\tau, F \rangle \in \text{Mod}^{\equiv}(\vdash_A)$. Together with Proposition 3.10, this yields that F is a singleton. Hence we conclude that \vdash is assertional.

To prove the “only if” part, suppose that \vdash is assertional. From Proposition 3.8(ii) we know that \vdash has a theorem $\varphi(x)$. Then let τ be the translation of \mathcal{L}_{\vdash_A} into \mathcal{L}_{\vdash} that sends $\top(x)$ to $\varphi(x)$. We shall see that τ is an interpretation of \vdash_A into \vdash . To this end, consider a matrix $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$. Since \vdash is assertional, there is $a \in A$ such that $F = \{a\}$. Together with the fact that $\varphi(x)$ is a theorem of \vdash , this implies that φ^A is the constant map with value a . Hence $A^\tau = \langle A; \varphi^A \rangle$ is essentially a pointed set B , and $\langle A^\tau, F \rangle = \langle B, \{\top_*^B\} \rangle$. By Proposition 3.10 this guarantees that $\langle A^\tau, F \rangle \in \text{Mod}^{\equiv}(\vdash_A)$. Hence we conclude that τ is an interpretation of \vdash_A into \vdash . \square

A logic \vdash is *truth-equational* [32] if there is a set $E(x)$ of equations such that for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$, and $a \in A$,

$$a \in F \iff A \vDash E(a).$$

Similarly, a logic \vdash is said to be *parametrically truth-equational* [26] if there is a set $E(x, \vec{y})$ of equations such that for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ with $F \neq \emptyset$, and $a \in A$,

$$a \in F \iff A \vDash E(a, \vec{c}) \text{ for every } \vec{c} \in A.$$

Theorem 3.12. *Parametrically truth-equational logics, and truth-equational logics form Leibniz classes.*

Proof. We detail only the proof of the fact that parametrically truth-equational logics form a Leibniz class. By Theorem 2.2 it will be enough to show that parametrically truth-equational logics are closed under term-equivalence, compatible expansions, and non-indexed products of sets. The fact that they are closed under term-equivalence, and compatible expansions is clear. Then consider a family $\{\vdash_i : i \in I\}$ of parametrically truth-equational logics. For every $j \in I$, let $E_j(x, \vec{y})$ be the set of equations witnessing the fact that \vdash_j is parametrically truth-equational. For every formula $\varphi(x, \vec{y})$ of \vdash_j , we denote by $\hat{\varphi}$ the sequence $\langle \varphi_i : i \in I \rangle$ in which $\varphi_i = x$ for every $i \in I \setminus \{j\}$, and $\varphi_j = \varphi$. Observe that $\hat{\varphi}(x, \vec{y})$ is a basic operation of $\otimes_{i \in I} \vdash_i$. Bearing this in mind, we define

$$E(x, \vec{y}) := \{ \hat{\varphi} \approx \hat{\psi} : \varphi \approx \psi \in \bigcup_{i \in I} E_i(x, \vec{y}) \}.$$

Then consider a matrix $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\otimes_{i \in I} \vdash_i)$ such that $F \neq \emptyset$. By [23, Prop. 4.5] we have $\langle \mathbf{A}, F \rangle \leq_{sd} \otimes_{i \in I} \langle \mathbf{A}_i, F_i \rangle$ for some $\langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ with $F_i \neq \emptyset$. For every $a \in A$,

$$\begin{aligned} a \in F &\iff a(i) \in F_i \text{ for every } i \in I \\ &\iff \mathbf{A}_i \vDash E_i(a(i), \vec{c}) \text{ for every } \vec{c} \in A_i \text{ and } i \in I \\ &\iff \mathbf{A} \vDash E(a, \vec{c}) \text{ for every } \vec{c} \in A. \end{aligned}$$

The above equivalences are justified as follows: the first is straightforward, the second follows from the fact the set $E_i(x, \vec{y})$ witnesses that \vdash_i is parametrically truth-equational, and the third is a consequence of the fact that the projection $\pi_i: A \rightarrow A_i$ is surjective for all $i \in I$. From the display above we obtain that $\otimes_{i \in I} \vdash_i$ is parametrically truth-equational, as desired. \square

Problem 1. Is it possible to find a transparent Leibniz condition Φ such that $\text{Log}(\Phi)$ is the class of (parametrically) truth-equational logics?

A logic \vdash is said to be *order algebraizable* [33] if there is a set $\Delta(x, y)$ of formulas and a set $E(x)$ of inequalities such that for every $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ the relation \preceq_F^A on A defined as follows is a partial order: for every $a, c \in A$,

$$a \preceq_F^A c \iff \Delta^A(a, c) \subseteq F,$$

and for every $a \in A$,

$$a \in F \iff \langle \mathbf{A}, \preceq_F^A \rangle \vDash E(e).$$

Theorem 3.13. *Order algebraizable logics, and logics with theorems form Leibniz classes.*

Proof. The result can be established by checking condition (ii) of Theorem 2.2 with ideas similar to the ones in the proof of Theorem 3.12. \square

We expect that a Leibniz condition defining the class of order algebraizable logics could be extracted from [33, Thm. 7.1(ii)].

Remark 3.14. Recall from Proposition 2.6 that every class of logics that can be written as the intersection of two Leibniz classes is still a Leibniz class. This observation can be exploited to show that some well-known collections of logics are Leibniz classes. For instance, a logic \vdash is said to be *algebraizable* [8] (resp. *weakly algebraizable* [13]) if it is equivalential (resp. protoalgebraic) and truth-equational. From Theorems 3.4, 3.12, and 3.7 it follows that (weakly) algebraizable logics form a Leibniz class. \square

We conclude this section by providing some examples of collections of logics that are not Leibniz classes. By Remark 2.5 we know that every class of logics that is not closed under the formation of extensions will serve this purpose. Among these we count, for instance, the class of logics with the Craig deductive interpolation property [14], and the class of logics with the infinite (resp. finite) Beth definability property [4, 5, 28].

To describe another interesting class of logics that is not a Leibniz class, recall that a logic \vdash has an *algebraic semantics* [8, 10] if there are a set of equations $E(x)$ and a class of algebras \mathbf{K} such that for every $\Gamma \cup \{\varphi\} \subseteq \text{Fm}(\vdash)$,

$$\Gamma \vdash \varphi \iff \bigcup \{E(\gamma) : \gamma \in \Gamma\} \vDash_{\mathbf{K}} E(\varphi)$$

where $\vDash_{\mathbf{K}}$ is the equational consequence relative to \mathbf{K} . Surprisingly enough, every logic is term-equivalent to one with an algebraic semantics, as shown essentially in [10, Thm.

3.1] (see [27, Cor. 6.3] for a detailed proof). Together with the fact that some logics lack an algebraic semantics [6], this yields that the class of logics with an algebraic semantics is not closed under term-equivalence, whence it is not a Leibniz class by Theorem 2.2(ii).

The reader familiar with abstract algebraic logic may be interested to know that also (fully) selfextensional, and (fully) Fregean logics [19] do not form Leibniz classes, since these collections are not closed under compatible expansions.

Problem 2. The majority of well-known Leibniz classes can be characterized in terms of the behaviour of the so-called *Leibniz operator*, i.e., the map $\Omega^A : \mathcal{P}(A) \rightarrow \text{Con}A$, defined on every algebra A , that associates a subset $F \subseteq A$ with the Leibniz congruence $\Omega^A F$. Is it possible to establish a precise relation between Leibniz classes and the behaviour of the Leibniz operator?

Later on we make use of the following well-known observation [18].

Proposition 3.15. *The classes of protoalgebraic, equivalential, assertional, order algebraizable, and truth-equational logics comprise only logics with theorems.*

4. IRREDUCIBILITY, PRIMALITY AND THEIR BOUNDARIES

Part of the speculative power of the identification of the Leibniz hierarchy with the lattice of all Leibniz classes comes from the fact that it allows to apply order-theoretic methods and intuitions to the study of the first. To explain how, recall that an element a of a lattice $\langle A; \wedge, \vee \rangle$ is said to be *meet-irreducible* if for every pair $b, c \in A$,

$$\text{if } a = b \wedge c, \text{ then either } a = b \text{ or } a = c.$$

Similarly, a is said to be *meet-prime* if for every pair $b, c \in A$,

$$\text{if } b \wedge c \leq a, \text{ then either } b \leq a \text{ or } c \leq a.$$

Accordingly, a is said to be *meet-reducible* when it is not meet-irreducible. It is clear that every meet-prime element of $\langle A; \wedge, \vee \rangle$ is meet-irreducible, while the converse is not true in general.

Since the Leibniz hierarchy is a lattice (Proposition 2.6), it makes sense to ask whether a Leibniz class is meet-irreducible or meet-prime. An affirmative answer to this question can then be regarded as a certificate that the Leibniz class under consideration captures a primitive or fundamental concept.

Remark 4.1. Since infima in the Leibniz hierarchy are intersections, a Leibniz class \mathbb{K} is meet-prime if there is no pair of logics $\vdash_1, \vdash_2 \notin \mathbb{K}$ such that $\vdash \in \mathbb{K}$, for every logic \vdash such that $\vdash_1, \vdash_2 \leq \vdash$. \square

Example 4.2. The Leibniz class of algebraizable (resp. weakly algebraizable) logics is meet-reducible, since it can be obtained as the intersection of the strictly larger Leibniz classes of equivalential (resp. protoalgebraic) logics and truth-equational logics (Remark 3.14). Moreover, there are meet-irreducible Leibniz classes that are not meet-prime, e.g., the class of logics \vdash for which there is no three-element algebra A and $a \in A$ such that $\langle A, \{a\} \rangle \in \text{Mod}^\equiv(\vdash)$. Even if we do not pursue the details here, the proof of this fact is an adaptation of an argument in [21, pag. 54]. \square

The next results put some boundaries to the expectation that well-known Leibniz classes should be meet-irreducible or meet-prime.

Proposition 4.3. *All Leibniz classes properly included into the class of logics with theorems are meet-reducible.*

Proof sketch. Given a logic \vdash with theorems, we denote by \vdash_{\emptyset} the logic on $Fm(\vdash)$ defined for every $\Gamma \cup \{\varphi\} \subseteq Fm(\vdash)$ as

$$\Gamma \vdash_{\emptyset} \varphi \iff \Gamma \neq \emptyset \text{ and } \Gamma \vdash \varphi.$$

It is clear that \vdash_{\emptyset} lacks theorems. It is not hard to see that

$$\text{Mod}^{\equiv}(\vdash_{\emptyset}) = \text{Mod}^{\equiv}(\vdash) \cup \{\langle A, \emptyset \rangle : \langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash) \text{ for some } F \subseteq A\}. \quad (3)$$

Then consider a Leibniz class \mathbb{K} properly included into the Leibniz class Thrms of logics with theorems. We set

$$\mathbb{K}_{\emptyset} := \mathbb{K} \cup \{\vdash_{\emptyset} : \vdash \in \mathbb{K}\}.$$

It is clear that $\mathbb{K} \subsetneq \mathbb{K}_{\emptyset}$, since \mathbb{K}_{\emptyset} contains logics without theorems. Moreover, $\mathbb{K} = \mathbb{K}_{\emptyset} \cap \text{Thrms}$. Therefore, to conclude that \mathbb{K} is meet-reducible, it will be enough to show that \mathbb{K}_{\emptyset} is a Leibniz class.

To prove this, consider a Leibniz condition $\Phi = \{\vdash^{\alpha} : \alpha \in \text{OR}\}$ such that $\mathbb{K} = \text{Log}(\Phi)$. We define $\Phi_{\emptyset} := \{\vdash_{\emptyset}^{\alpha} : \alpha \in \text{OR}\}$. From the fact that Φ is a Leibniz condition and (3) it follows that Φ_{\emptyset} is also Leibniz condition. We shall prove that $\mathbb{K}_{\emptyset} = \text{Log}(\Phi_{\emptyset})$. To this end, consider a logic $\vdash \in \mathbb{K}$. Then there is $\alpha \in \text{OR}$ such that $\vdash^{\alpha} \leq \vdash$. Together with (3), this implies that $\vdash_{\emptyset}^{\alpha} \leq \vdash, \vdash_{\emptyset}$. Hence we obtain $\mathbb{K}_{\emptyset} \subseteq \text{Log}(\Phi_{\emptyset})$. To prove the other inclusion, consider $\alpha \in \text{OR}$ and a logic \vdash with an interpretation τ of $\vdash_{\emptyset}^{\alpha}$ into \vdash . If \vdash has theorems, then we can use (3) to conclude that τ is also an interpretation of \vdash^{α} into \vdash , whence $\vdash \in \text{Log}(\Phi) = \mathbb{K} \subseteq \mathbb{K}_{\emptyset}$. Then we consider the case where \vdash lacks theorems. Since $\vdash^{\alpha} \in \mathbb{K}$, the logic \vdash^{α} has a theorem $\varphi(x)$. Let y be a variable different from x and observe that $y \vdash_{\emptyset}^{\alpha} \varphi(x)$. By [23, Prop. 3.3] this yields $y \vdash \tau(\varphi(x))$. Then consider the logic \vdash^{+} on $Fm(\vdash)$ induced by the class of matrices

$$\{\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash) : F \neq \emptyset\}. \quad (4)$$

Since $y \vdash \tau(\varphi(x))$, it is clear that $\emptyset \vdash^{+} \tau(\varphi)$. In particular, this guarantees that $\text{Mod}^{\equiv}(\vdash^{+})$ is the class of matrices in (4). This fact, together with (3), yields that τ is an interpretation of \vdash^{α} into \vdash^{+} . As a consequence, we obtain $\vdash^{+} \in \mathbb{K}$. Since \vdash coincides with \vdash_{\emptyset}^{+} , we conclude that $\vdash \in \mathbb{K}_{\emptyset}$. This establishes that $\mathbb{K}_{\emptyset} = \text{Log}(\Phi_{\emptyset})$ and, therefore, that \mathbb{K} is a Leibniz class. \square

Remark 4.4. Proposition 4.3 indicates that Leibniz classes comprising only logics with theorems (but not all of them) cannot be meet-irreducible in an absolute sense. For this reason, we say that a Leibniz class \mathbb{K} is meet-reducible *among logics with theorems* if for every pair of Leibniz classes \mathbb{K}_1 and \mathbb{K}_2 comprising logics with theorems only, if $\mathbb{K} = \mathbb{K}_1 \cap \mathbb{K}_2$, then either $\mathbb{K} = \mathbb{K}_1$ or $\mathbb{K} = \mathbb{K}_2$. A similar definition applies to the case of meet-prime Leibniz classes. \square

Proposition 4.5. *All Leibniz classes properly included into the class of assertional logics are not meet-prime among Leibniz classes with theorems.*

Proof. Let \mathbb{F} be the class of logics \vdash with theorems such that if $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$, then either the matrix $\langle A, F \rangle$ is trivial or $|F| \geq 2$. Using the characterization of Leibniz classes given in Theorem 2.2(ii) it is not hard to see that \mathbb{F} is indeed a Leibniz class.

Now, let Asrt be the Leibniz class of assertional logics, and consider an arbitrary Leibniz class \mathbb{K} properly included into Asrt . It is clear that both Asrt and \mathbb{F} are not included into \mathbb{K} . Bearing this in mind, it only remains to show that $\text{Asrt} \cap \mathbb{F} \subseteq \mathbb{K}$. To this end, consider a logic $\vdash \in \text{Asrt} \cap \mathbb{F}$. Observe that every matrix $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ is such that F is a singleton (as $\vdash \in \text{Asrt}$) and, therefore, is trivial (as $\vdash \in \mathbb{F}$). Thus $\text{Mod}^{\equiv}(\vdash)$ is the class of trivial matrices in the language of \mathcal{L}_{\vdash} . As a consequence, the logic \vdash is inconsistent [23, Lem. 7.1]. In particular, this guarantees that $\llbracket \vdash \rrbracket$ is the maximum of Log [23, Thm. 7.3]. By Theorem 2.2(iii) the collection \mathbb{K}^{\dagger} is a non-empty filter of Log which implies $\llbracket \vdash \rrbracket \in \mathbb{K}^{\dagger}$. Together with $\mathbb{K} = \{\vdash' : \llbracket \vdash' \rrbracket \in \mathbb{K}^{\dagger}\}$, this yields $\vdash \in \mathbb{K}$, as desired. \square

5. MEET-PRIME LEIBNIZ CLASSES

In this section we show that the Leibniz class of all logics with theorems, and the Leibniz class of truth-minimal logics that we introduce below are meet-prime in the absolute sense. On the other hand, it is proved that the Leibniz classes of assertional and truth-equational logics are meet-prime among the logics with theorems. As both assertional and truth-equational logics have theorems, Proposition 4.3 guarantees that the restriction to logics with theorems cannot be dropped here. For the present purpose, it is convenient to start the discussion from the new class of truth-minimal logics.

5.1. Truth-minimal logics.

Definition 5.1. A logic \vdash is *truth-minimal* if for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$,

if $\langle A, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ and $G \subseteq F$, then either $G = F$ or $G = \emptyset$.

The next result is instrumental to construct examples of truth-minimal logics.

Proposition 5.2. Let κ be an infinite cardinal and M a class of matrices such that $|F| \leq 1$ for every $\langle A, F \rangle \in M$. The logic \vdash induced by M formulated in κ variables is truth-minimal.

Proof. The assumption that $|F| \leq 1$ for every $\langle A, F \rangle \in M$ implies that $x, y, \varphi(x, \vec{z}) \vdash \varphi(y, \vec{z})$ for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash)$. By Proposition 3.8(i) we conclude that \vdash is truth-minimal. \square

Our first aim is to show that truth-minimal logics constitute a Leibniz class. To this end, we make use of the following observation:

Lemma 5.3. Let $\{\vdash_i : i \in I\}$ be a family of logics, and $A \leq_{\text{sd}} \bigotimes_{i \in I} A_i$ where A_i is an \mathcal{L}_{\vdash_i} -algebra for every $i \in I$. If $G \neq \emptyset$ is a deductive filter of $\bigotimes_{i \in I} \vdash_i$ on A , then for every $j \in I$ there is a deductive filter G_j of \vdash_j on A_j such that $G = A \cap \prod_{i \in I} G_i$.

Proof. For every $j \in I$, we denote the natural projection map by $\pi_j : A \rightarrow A_j$, and set $G_j := \pi_j[G]$.

Then we turn to prove that $G = A \cap \prod_{i \in I} G_i$. The inclusion from left to right is clear. To prove the other one, consider an element $a \in A \cap \prod_{i \in I} G_i$. For every $j \in I$ there is $c_j \in G$ such that $c_j(j) = a(j)$. Then consider the basic operation $x \dashv\!\!\!\dashv_j y$ of $\bigotimes_{i \in I} \vdash_i$, whose j -th component is the projection on the first coordinate, and whose i -th component is the projection on the second coordinate for every $i \in I \setminus \{j\}$. Since $c_j(j) = a(j)$,

$$a \dashv\!\!\!\dashv_j^A c_j = c_j \in G. \quad (5)$$

Now, consider some distinct variables $\{y_j: j \in I\} \cup \{x\} \subseteq Fm(\otimes_{i \in I} \vdash_i)$. Bearing in mind that $\otimes_{i \in I} \vdash_i$ is the logic induced by the class $\otimes_{i \in I} \text{Mod}^{\equiv}(\vdash_i)$, it is easy to see that the following rule is valid in $\otimes_{i \in I} \vdash_i$:

$$\{x \multimap_j y_j: j \in I\} \triangleright x.$$

Together with (5) and the fact that G is a deductive filter of $\otimes_{i \in I} \vdash_i$ on A , this implies $a \in G$. Hence we conclude

$$G = A \cap \prod_{i \in I} G_i. \quad (6)$$

Consider an index $j \in I$. To conclude the proof, it suffices to show that G_j is a filter of \vdash_j on A_j . To prove this, suppose that $\Gamma \vdash_j \varphi$ and consider a homomorphism $h: Fm(\vdash_j) \rightarrow A_j$ such that $h[\Gamma] \subseteq G_j$. For every n -ary term $\psi(z_1, \dots, z_n) \in Fm(\vdash_j)$, let $\hat{\psi}(z_1, \dots, z_n)$ be an arbitrary basic n -ary operation of $\otimes_{i \in I} \vdash_i$ whose j -th component is ψ . Moreover, consider a variable $y \in Fm(\otimes_{i \in I} \vdash_i)$ not occurring in Γ .² Again bearing in mind that $\otimes_{i \in I} \vdash_i$ is the logic induced by the class $\otimes_{i \in I} \text{Mod}^{\equiv}(\vdash_i)$, it is clear that the following rule is valid in $\otimes_{i \in I} \vdash_i$:

$$\{\hat{\gamma} \multimap_j y: \gamma \in \Gamma\} \triangleright \hat{\varphi} \multimap_j y. \quad (7)$$

Now, choose an element $a \in G$, and let $f: Fm(\otimes_{i \in I} \vdash_i) \rightarrow A$ be a homomorphism such that $f(y) = a$ and $f(x)(j) = h(x)$ for every variable $x \in Fm(\vdash_j)$. Observe that there exists such an f , since the projection $\pi_j: A \rightarrow A_j$ is surjective. For every $i \in I \setminus \{j\}$ and $\gamma \in \Gamma$, we have

$$\begin{aligned} f(\hat{\gamma} \multimap_j y)(i) &= a(i) \in \pi_i[G] = G_i \\ f(\hat{\gamma} \multimap_j y)(j) &= h(\gamma) \in G_j. \end{aligned}$$

Together with (6), this implies

$$f(\hat{\gamma} \multimap_j y) \in A \cap \prod_{i \in I} G_i = G, \text{ for every } \gamma \in \Gamma.$$

The above display, together with (7) and the fact that G is a deductive filter of $\otimes_{i \in I} \vdash_i$ on A , yields that $f(\hat{\varphi} \multimap_j y) \in G$. As a consequence, we obtain

$$h(\varphi) = f(\hat{\varphi} \multimap_j y)(j) \in \pi_j[G] = G_j.$$

Hence we conclude that G_j is a deductive filter of \vdash_j on A_j . \(\square\)

As a consequence, we obtain the following:

Theorem 5.4. *The class of truth-minimal logics is a Leibniz class.*

Proof. It is straightforward that the class of truth-minimal logics is closed under term-equivalence and compatible expansions. In the light of Theorem 2.2, it only remains to prove that this class is closed under the formation of non-indexed products of sets.

To prove this, consider a family $\{\vdash_i: i \in I\}$ of truth-minimal logics. Moreover, consider $\langle A, F \rangle, \langle A, G \rangle \in \text{Mod}^{\equiv}(\otimes_{i \in I} \vdash_i)$ with $\emptyset \neq G \subseteq F$. We need to show that $F = G$. By [23, Prop. 4.5] there is a family of matrices $\{\langle A_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i): i \in I\}$ such that $\langle A, F \rangle \leq_{\text{sd}}$

²In case there is no such a variable y , we replace in the proof $\Gamma \cup \{\varphi\}$ by $\sigma[\Gamma \cup \varphi]$, where σ is a substitution on $Fm(\vdash_j)$ that maps variables to variables, and that is injective but not surjective.

$\bigotimes_{i \in I} \langle A_i, F_i \rangle$. By Lemma 5.3, for every $j \in I$ there is a deductive filter G_j of \vdash_j on A_j such that

$$\emptyset \neq G_j \subseteq F_j \quad (8)$$

and $G = A \cap \prod_{i \in I} G_i$. Now, from the fact that G_j is a deductive filter of \vdash_j and $G_j \subseteq F_j$ it follows that $\tilde{\Omega}_{\vdash_j}^A G_j \subseteq \tilde{\Omega}_{\vdash_j}^A F_j$. Therefore, bearing in mind that $\tilde{\Omega}_{\vdash_j}^A F_j$ is the identity relation, we conclude that the same holds for $\tilde{\Omega}_{\vdash_j}^A G_j$. In particular, this yields

$$\langle A_j, G_j \rangle \in \text{Mod}^{\equiv}(\vdash_j). \quad (9)$$

Finally, for every $j \in J$ we can apply the fact that \vdash_j is truth-minimal to (8, 9), obtaining that $F_j = G_j$. As a consequence, we get

$$G = A \cap \prod_{i \in I} G_i = A \cap \prod_{i \in I} F_i = F.$$

Hence we conclude that $\bigotimes_{i \in I} \vdash_i$ is truth-minimal. \square

To prove that the Leibniz class of truth-minimal logics is meet-prime, we rely on the following technical observations:

Lemma 5.5. *Let \vdash be a logic. If $\langle A, A \rangle \in \text{Mod}^{\equiv}(\vdash)$, then A is the trivial algebra.*

Proof. From [23, Prop. 2.2(ii)] we obtain $\tilde{\Omega}_{\vdash}^A A = A \times A$. On the other hand, since $\langle A, A \rangle \in \text{Mod}^{\equiv}(\vdash)$, the congruence $\tilde{\Omega}_{\vdash}^A A$ is the identity relation. But the fact that $A \times A$ is the identity relation implies that A is a singleton, as desired. \square

Lemma 5.6. *Let \vdash be a logic, and $\langle A, F \rangle, \langle A, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \neq G \subsetneq F$. There are $\langle B, F^{\dagger} \rangle, \langle B, G^{\dagger} \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \neq G^{\dagger} \subsetneq F^{\dagger} \subsetneq B$ and*

$$\omega \leq |B| = |G^{\dagger}| = |F^{\dagger} \setminus G^{\dagger}| = |B \setminus F^{\dagger}|.$$

Proof. Choose an infinite cardinal $\kappa \geq |A|$ and define $B := A^{\kappa} \times A^{\kappa}$. Then set

$$F^{\dagger} := F^{\kappa} \times F^{\kappa} \text{ and } G^{\dagger} := G^{\kappa} \times G^{\kappa}.$$

Since $\text{Mod}^{\equiv}(\vdash)$ is closed under direct products [23, Lem. 2.3], we obtain

$$\langle B, F^{\dagger} \rangle, \langle B, G^{\dagger} \rangle \in \text{Mod}^{\equiv}(\vdash).$$

Moreover, from $\emptyset \neq G \subsetneq F$ it follows $\emptyset \neq G^{\dagger} \subsetneq F^{\dagger}$.

Observe that since $\emptyset \neq G \subsetneq F$, the set F contains at least two distinct elements. This fact and $\emptyset \neq G \subsetneq F$ guarantee that

$$2^{\kappa} \leq |F^{\kappa} \times G^{\kappa}| = |G^{\dagger}| \quad (10)$$

$$2^{\kappa} \leq |F^{\kappa}| \leq |F^{\kappa} \times (F^{\kappa} \setminus G^{\kappa})| = |F^{\dagger} \setminus G^{\dagger}| \quad (11)$$

Since F contains at least two elements and $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$, we can apply Lemma 5.5 obtaining $F \subsetneq A$. Bearing this in mind, we get $F^{\dagger} \subsetneq B$ and

$$2^{\kappa} \leq |A^{\kappa}| \leq |A^{\kappa} \times (A^{\kappa} \setminus F^{\kappa})| \leq |B \setminus F^{\dagger}|. \quad (12)$$

Finally, from $\kappa \geq \omega + |A|$ it follows $|B| = |A^{\kappa} \times A^{\kappa}| = 2^{\kappa}$. But, together with (10, 11, 12), this yields

$$\omega \leq |B| = |G^{\dagger}| = |F^{\dagger} \setminus G^{\dagger}| = |B \setminus F^{\dagger}|. \quad \square$$

Lemma 5.7. *Let \vdash_1 and \vdash_2 be logics, and for every $i = 1, 2$ let $\langle A_i, F_i \rangle, \langle A_i, G_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $\emptyset \neq G_i \subsetneq F_i$. There are sets $\emptyset \neq G \subsetneq F \subsetneq B$, an \mathcal{L}_{\vdash_1} -algebra \mathbf{B}_1 , and an \mathcal{L}_{\vdash_2} -algebra \mathbf{B}_2 such that $B_1 = B_2 = B$, and*

$$\begin{aligned} \langle \mathbf{B}_1, F \rangle, \langle \mathbf{B}_1, G \rangle &\in \text{Mod}^{\equiv}(\vdash_1) \\ \langle \mathbf{B}_2, F \rangle, \langle \mathbf{B}_2, G \rangle &\in \text{Mod}^{\equiv}(\vdash_2). \end{aligned}$$

Proof. By Lemma 5.6 for every $i = 1, 2$ there are $\langle C_i, F_i^+ \rangle, \langle C_i, G_i^+ \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $\emptyset \neq G_i^+ \subsetneq F_i^+ \subsetneq C_i$ and $\omega \leq |C_i| = |G_i^+| = |F_i^+ \setminus G_i^+| = |C_i \setminus F_i^+|$.

Let then $\kappa := \max\{|C_1|, |C_2|\}$. Consider also $i = 1, 2$. Since $\text{Mod}^{\equiv}(\vdash_i)$ is closed under direct powers [23, Lem. 2.3], we obtain

$$\langle C_i^\kappa, (F_i^+)^{\kappa} \rangle, \langle C_i^\kappa, (G_i^+)^{\kappa} \rangle \in \text{Mod}^{\equiv}(\vdash_i). \quad (13)$$

Moreover, since $\emptyset \neq G_i^+ \subsetneq F_i^+ \subsetneq C_i$, we have

$$\emptyset \neq G_i^{\kappa} \subsetneq F_i^{\kappa} \subsetneq C_i^\kappa.$$

Finally, since $\omega \leq |C_i| = |G_i^+| = |F_i^+ \setminus G_i^+| = |C_i \setminus F_i^+| \leq \kappa$, we have

$$\kappa = |C_i^\kappa| = |G_i^{\kappa}| = |F_i^{\kappa} \setminus G_i^{\kappa}| = |C_i^\kappa \setminus F_i^{\kappa}|. \quad (14)$$

Then consider some sets $\emptyset \neq G \subsetneq F \subsetneq B$ such that

$$\kappa = |B| = |G| = |F \setminus G| = |B \setminus F|.$$

From (13) and (14) it easily follows that for each $i = 1, 2$ there is an \mathcal{L}_{\vdash_i} -algebra \mathbf{B}_i with universe B such that $\langle \mathbf{B}_i, F \rangle \cong \langle C_i^\kappa, (F_i^+)^{\kappa} \rangle$, $\langle \mathbf{B}_i, G \rangle \cong \langle C_i^\kappa, (G_i^+)^{\kappa} \rangle$ and, therefore, $\langle \mathbf{B}_i, F \rangle, \langle \mathbf{B}_i, G \rangle \in \text{Mod}^{\equiv}(\vdash_i)$. \square

We are now ready to prove the main result of this part:

Theorem 5.8. *The Leibniz class of truth-minimal logics is meet-prime.*

Proof. Consider two logics \vdash_1 and \vdash_2 that are not truth-minimal. It will be enough to construct a logic \vdash that is not truth-minimal and in which \vdash_1 and \vdash_2 are interpretable.

To this end, observe that for every $i = 1, 2$ there are $\langle A_i, F_i \rangle, \langle A_i, G_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $\emptyset \neq G_i \subsetneq F_i$. From Lemma 5.7 we obtain

$$\langle \mathbf{B}_1, F \rangle, \langle \mathbf{B}_1, G \rangle \in \text{Mod}^{\equiv}(\vdash_1) \text{ and } \langle \mathbf{B}_2, F \rangle, \langle \mathbf{B}_2, G \rangle \in \text{Mod}^{\equiv}(\vdash_2)$$

for some sets $\emptyset \neq G \subsetneq F \subsetneq B$, an \mathcal{L}_{\vdash_1} -algebra \mathbf{B}_1 , and an \mathcal{L}_{\vdash_2} -algebra \mathbf{B}_2 such that $B_1 = B_2 = B$.

Let \mathbf{B} be the common expansion of \mathbf{B}_1 and \mathbf{B}_2 with all finitary operations on B . Moreover, let \vdash be the logic, formulated in $\kappa := \max\{|Fm(\vdash_1)|, |Fm(\vdash_2)|\}$ variables induced by the set of matrices

$$\mathbf{K} := \{ \langle \mathbf{B}, F \rangle, \langle \mathbf{B}, G \rangle \}.$$

We claim that \vdash is equivalential. To prove this, consider elements $0 \in B \setminus F$ and $1 \in G$ (this is possible, since $\emptyset \neq G$ and $F \subsetneq B$). Then let $x \multimap y$ be the binary basic operation of \mathbf{B} defined as follows for every $a, c \in B$,

$$a \multimap^{\mathbf{B}} c := \begin{cases} 1 & \text{if } a = c \\ 0 & \text{otherwise.} \end{cases}$$

Bearing in mind that $0 \notin F \cup G$ and $1 \in F \cap G$, it is not hard to see that the set $\Delta(x, y) := \{x \multimap y\}$ satisfies the conditions in the right hand side of [23, Thm. 2.7]. As a consequence we obtain that \vdash is equivalential, establishing the claim.

Since \mathbf{B} is endowed with all unary constant maps, we have $\mathbb{S}(\mathbf{K}) = \mathbf{K}$. Moreover, for each pair of different $a, c \in B$, we have

$$a \multimap^B c = 0 \notin G \cup F \text{ and } a \multimap^B a = 1 \in G \cap F.$$

By [23, Prop. 2.2(i)] this implies that the matrices in \mathbf{K} are reduced. This fact, together with the claim and $\mathbb{S}(\mathbf{K}) = \mathbf{K}$, allows us to apply [23, Prop. 3.9] obtaining that the identity maps are interpretations of \vdash_1 and \vdash_2 into \vdash . Moreover, \vdash is not truth-minimal, as witnessed by the fact that $\mathbf{K} \subseteq \text{Mod}^{\equiv}(\vdash)$ and $\emptyset \neq G \subsetneq F$. \square

The proof strategy described above can be adapted to the case of truth-equational and assertional logics, as we proceed to explain.

5.2. Truth-equational logics. To prove that the Leibniz class of truth-equational logics is meet-prime among logics with theorems, it is convenient to recall the following characterization result.

Theorem 5.9. *A logic \vdash is truth-equational if and only if for every algebra A ,*

$$\text{if } \langle A, F \rangle, \langle A, G \rangle \in \text{Mod}^{\equiv}(\vdash), \text{ then } F = G.$$

Proof. See [32, Prop. 17 and Thm. 2.8]. \square

As a consequence we obtain the desired result.

Theorem 5.10. *The Leibniz class of truth-equational logics is meet-prime among logics with theorems.*

Proof. Consider two logics \vdash_1 and \vdash_2 with theorems that are not truth-equational. It will be enough to construct a logic \vdash that is not truth-equational and in which \vdash_1 and \vdash_2 are interpretable.

First we claim that for every $i = 1, 2$ there are $\langle A_i, F_i \rangle, \langle A_i, G_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $\emptyset \neq G_i \subsetneq F_i$. To prove this, consider $i = 1, 2$. Since the logic \vdash_i is not truth-equational, we can apply Theorem 5.9 obtaining an algebra A_i and distinct $U_i, V_i \subseteq A_i$ such that $\langle A_i, U_i \rangle, \langle A_i, V_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$. We can assume without loss of generality that $V_i \setminus U_i \neq \emptyset$. Bearing this in mind, we define $F_i := V_i$ and $G_i := V_i \cap U_i$. It is clear that $\langle A_i, G_i \rangle \in \text{Mod}(\vdash_i)$ and $G_i \subsetneq F_i$. Since \vdash_i has theorems, the fact that $\langle A_i, G_i \rangle \in \text{Mod}(\vdash_i)$ implies that $G_i \neq \emptyset$. Moreover, since $G_i \subseteq F_i = V_i$, we have

$$\tilde{\Omega}_{\vdash_i}^{A_i} G_i \subseteq \tilde{\Omega}_{\vdash_i}^{A_i} V_i.$$

Recall that $\tilde{\Omega}_{\vdash_i}^{A_i} V_i$ is the identity relation, since $\langle A_i, V_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$. Hence $\tilde{\Omega}_{\vdash_i}^{A_i} G_i$ is also the identity relation and, therefore, $\langle A_i, G_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$. Finally, by the assumptions, $\langle A_i, F_i \rangle = \langle A_i, V_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$. This concludes the proof of the claim.

Together with Lemma 5.7, the claim implies that there are

$$\langle \mathbf{B}_1, F \rangle, \langle \mathbf{B}_1, G \rangle \in \text{Mod}^{\equiv}(\vdash_1) \text{ and } \langle \mathbf{B}_2, F \rangle, \langle \mathbf{B}_2, G \rangle \in \text{Mod}^{\equiv}(\vdash_2)$$

for some sets $\emptyset \neq G \subsetneq F \subsetneq B$, an \mathcal{L}_{\vdash_1} -algebra \mathbf{B}_1 , and an \mathcal{L}_{\vdash_2} -algebra \mathbf{B}_2 such that $B_1 = B_2 = B$.

Now, let \mathbf{B} be the common expansion of \mathbf{B}_1 and \mathbf{B}_2 with all finitary operations on B . Moreover, let \vdash be the logic, formulated in $\kappa := \max\{|Fm(\vdash_1)|, |Fm(\vdash_2)|\}$ variables induced by the set of matrices

$$\mathbf{K} := \{\langle \mathbf{B}, F \rangle, \langle \mathbf{B}, G \rangle\}.$$

As in the proof of Theorem 5.8, we obtain $\mathbf{K} \subseteq \text{Mod}^{\equiv}(\vdash)$ and $\vdash_1, \vdash_2 \leq \vdash$. Finally, applying Theorem 5.9 to the fact that $\mathbf{K} \subseteq \text{Mod}^{\equiv}(\vdash)$ and $F \neq G$, we conclude that \vdash is not truth-equational. \square

5.3. Assertional logics.

Theorem 5.11. *The Leibniz class of assertional logics is meet-prime among logics with theorems.*

Proof. Consider two logics \vdash_1 and \vdash_2 with theorems that are not assertional. As usual, it will be enough to construct a logic \vdash that is not assertional and in which \vdash_1 and \vdash_2 are interpretable.

We claim that for every $i = 1, 2$ there are $\langle A_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $|F_i| \geq 2$ and $F_i \subsetneq A_i$. To prove this, consider $i = 1, 2$, and observe that the fact that \vdash_i is not assertional implies that there is $\langle A_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that F_i is not a singleton. Since \vdash_i has theorems and $\langle A_i, F_i \rangle$ is a model of \vdash_i we know that $F_i \neq \emptyset$ and, therefore, $|F_i| \geq 2$. This fact, together with Lemma 5.5, ensures that $F_i \subsetneq A_i$, establishing the claim.

By the claim we can find a cardinal κ large enough to guarantee that

$$|A_1^\kappa| = |F_1^\kappa| = |A_1^\kappa \setminus F_1^\kappa| = |A_2^\kappa| = |F_2^\kappa| = |A_2^\kappa \setminus F_2^\kappa|.$$

Thus there is a set B , a set $F \subseteq B$, an \mathcal{L}_{\vdash_1} -algebra \mathbf{B}_1 , and an \mathcal{L}_{\vdash_2} -algebra \mathbf{B}_2 such that $B_1 = B_2 = B$ and

$$\langle \mathbf{B}_1, F \rangle \cong \langle A_1^\kappa, F_1^\kappa \rangle \text{ and } \langle \mathbf{B}_2, F \rangle \cong \langle A_2^\kappa, F_2^\kappa \rangle.$$

Since $\text{Mod}^{\equiv}(\vdash_i)$ is closed under the formation of direct powers for every $i = 1, 2$ [23, Lem. 2.3], we obtain $\langle \mathbf{B}_1, F \rangle \in \text{Mod}^{\equiv}(\vdash_1)$ and $\langle \mathbf{B}_2, F \rangle \in \text{Mod}^{\equiv}(\vdash_2)$.

Now, let \mathbf{B} be the common expansion of \mathbf{B}_1 and \mathbf{B}_2 with all finitary operations on B . Moreover, let \vdash be the logic, formulated in $\kappa := \max\{|Fm(\vdash_1)|, |Fm(\vdash_2)|\}$ variables induced by the matrix $\langle \mathbf{B}, F \rangle$. As in the proof of Theorem 5.8, we obtain $\langle \mathbf{B}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ and $\vdash_1, \vdash_2 \leq \vdash$. Finally, since $\langle \mathbf{B}, F \rangle$ and F is not a singleton, we conclude that \vdash is not assertional. \square

Remark 5.12. In Theorems 5.10 and 5.11 it is shown that the Leibniz classes of truth-equational and assertional logics are meet-prime among logics with theorems. As we mentioned, in the light of Proposition 4.3 this restriction cannot be dropped. \square

5.4. Logics with theorems.

Theorem 5.13. *The Leibniz class of logics with theorems is meet-prime.*

Proof. Let \vdash_{AI} be an almost inconsistent logic, and recall that if \vdash is a logic without theorems, then $\vdash \leq \vdash_{\text{AI}}$ [23, Thm. 7.3]. Together with the fact that \vdash_{IA} lacks theorems, this immediately implies that the Leibniz class of logics with theorems is meet-prime. \square

6. MEET-REDUCIBLE LEIBNIZ CLASSES

Recall that protoalgebraic, equivalential, and order algebraizable logics have theorems. In the light of Proposition 4.3 the corresponding Leibniz classes are trivially meet-reducible. In this section we strengthen this result by proving that they remain meet-reducible even among the restricted setting of logics with theorems. In addition, it is shown that the Leibniz class of parametrically truth-equational logics is meet-reducible in the absolute sense.³ Remarkably, the latter result cannot be inferred directly from Proposition 4.3, since parametrically truth-equational logics need not have theorems [26, Sec. 4]. In what follows we rely on the next technical observation:

Proposition 6.1. *Let \vdash_1, \vdash_2 , and \vdash be logics, and τ an interpretation of $\vdash_1 \otimes \vdash_2$ into \vdash . Then for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ there are $\langle A_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_1)$, $\langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_2)$, and an isomorphism*

$$f: \langle A_1 \otimes A_2, F_1 \times F_2 \rangle \rightarrow \langle A^\tau, F \rangle.$$

Moreover, for every submatrix $\langle B, F \cap B \rangle \subseteq \langle A, F \rangle$ and $i = 1, 2$, there is a submatrix $\langle B_i, F_i \cap B_i \rangle \subseteq \langle A_i, F_i \rangle$ such that:

(i) the following restriction of f is a well-defined isomorphism:

$$f: \langle B_1 \otimes B_2, (F_1 \cap B_1) \times (F_2 \cap B_2) \rangle \rightarrow \langle B^\tau, F \cap B \rangle;$$

(ii) if $F \cap B \neq \emptyset$, then for every $\theta \in \text{Con}B$ compatible with $F \cap B$ and every $i = 1, 2$, there is $\theta_i \in \text{Con}B_i$ compatible with $F_i \cap B_i$ such that

$$\theta = \{ \langle f \langle a, b \rangle, f \langle c, d \rangle \rangle : \langle a, c \rangle \in \theta_1 \text{ and } \langle b, d \rangle \in \theta_2 \}.$$

Proof. The fact that there are $\langle A_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_1)$, $\langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_2)$, and an isomorphism $f: \langle A_1 \otimes A_2, F_1 \times F_2 \rangle \rightarrow \langle A^\tau, F \rangle$ is a direct consequence of [23, Cor. 4.14]. From now on we assume without loss of generality that f is the identity map and, therefore, that

$$\langle A^\tau, F \rangle = \langle A_1 \otimes A_2, F_1 \times F_2 \rangle.$$

(i): Consider a submatrix $\langle B, F \cap B \rangle \subseteq \langle A, F \rangle$. Therefore, $B^\tau \subseteq A^\tau = A_1 \otimes A_2$. As a consequence, for every $i = 1, 2$ there is $B_i \subseteq A_i$ such that $B^\tau = B_1 \otimes B_2$ [36, Lem. 1.10]. Moreover, since $B = B_1 \times B_2$ and $F = F_1 \times F_2$,

$$F \cap B = (F_1 \times F_2) \cap (B_1 \times B_2) = (F_1 \cap B_1) \times (F_2 \cap B_2).$$

As a consequence, we obtain

$$\langle B^\tau, F \cap B \rangle = \langle B_1 \otimes B_2, (F_1 \cap B_1) \times (F_2 \cap B_2) \rangle.$$

(ii): Consider $\theta \in \text{Con}B$ compatible with $F \cap B$. Clearly $\theta \in \text{Con}B^\tau = \text{Con}(B_1 \otimes B_2)$. As shown in [36, Lem. 1.12], for every $i = 1, 2$ there is $\theta_i \in \text{Con}B_i$ such that

$$\theta = \{ \langle \langle a, b \rangle, \langle c, d \rangle \rangle : \langle a, c \rangle \in \theta_1 \text{ and } \langle b, d \rangle \in \theta_2 \}.$$

We turn to prove that θ_1 is compatible with $F_1 \cap B_1$. To this end, consider $a, c \in B_1$ such that $a \in F_1 \cap B_1$ and $\langle a, c \rangle \in \theta_1$. From the assumption we have $(F_1 \cap B_1) \times (F_2 \cap B_2) = F \cap B \neq \emptyset$. Then there is $b \in F_2 \cap B_2$. From the above display it follows that $\langle \langle a, b \rangle, \langle c, b \rangle \rangle \in \theta$. Since θ is compatible with F and $\langle a, b \rangle \in F_1 \times F_2 = F$, we get $\langle c, b \rangle \in F = F_1 \times F_2$. In

³These negative results are compensated by some positive ones in Section 7 at least for what concerns protoalgebraic and equivalential logics.

particular, this guarantees that $c \in F_1 \cap B_1$. As a consequence we conclude that θ_1 is compatible with $F_1 \cap B_1$. A similar argument shows that θ_2 is compatible with $F_2 \cap B_2$. \square

6.1. Protoalgebraic logics. Our aim is to show that the Leibniz class of protoalgebraic logics is meet-reducible among logics with theorems. To this end, it is useful to recall a few concepts. An algebra A is said to be *congruence uniform* [3, Sec. 7.1] if $|a/\theta| = |b/\theta|$, for every $a, b \in A$ and $\theta \in \text{Con}A$. It is well-known that Boolean algebras are congruence uniform.

We denote by BA the variety of Boolean algebras, and by \vdash_{BA}^* the logic formulated in countably many variables induced by the following class of matrices:

$$\{\langle A, F \rangle : A \in \text{BA} \text{ and } 1 \in F\}.$$

Lemma 6.2. *The logic \vdash_{BA}^* has theorems, but is not protoalgebraic. Moreover, the algebraic reducts of the matrices in $\text{Mod}^{\equiv}(\vdash_{\text{BA}}^*)$ belong to BA .*

Proof. Clearly 1 is a theorem of \vdash_{BA}^* . Moreover, the fact that the algebraic reducts of the matrices in $\text{Mod}^{\equiv}(\vdash_{\text{BA}}^*)$ belong to BA is an immediate consequence of [23, Cor. 2.6].

It only remains to prove that \vdash_{BA}^* is not protoalgebraic. Suppose the contrary, with a view to contradiction. Then there is a set $\Delta(x, y, \vec{z})$ of congruence formulas with parameters for \vdash_{BA}^* . We consider the four-element Boolean algebra A with universe $\{a, b, 0, 1\}$, where 0 and 1 are respectively the bottom and the top element of the lattice order. Then we set $F := \{1, a\}$ and $G := \{1, a, b\}$. From the definition of \vdash_{BA}^* it follows that $\langle A, F \rangle, \langle A, G \rangle \in \text{Mod}(\vdash_{\text{BA}}^*)$. Together with the fact that Δ is a set of congruence formulas with parameters for \vdash_{BA}^* and that $F \subseteq G$, this yields that for every $p, q \in A$,

$$\begin{aligned} \langle p, q \rangle \in \Omega^A F &\iff \Delta^A(p, q, \vec{c}) \subseteq F, \text{ for every } \vec{c} \in A \\ &\implies \Delta^A(p, q, \vec{c}) \subseteq G, \text{ for every } \vec{c} \in A \\ &\iff \langle p, q \rangle \in \Omega^A G. \end{aligned}$$

Hence we conclude that $\Omega^A F \subseteq \Omega^A G$. On the other hand, it is easy to see that $\Omega^A G$ is the identity relation, while $\Omega^A F$ is the congruence with blocks $\{1, a\}$ and $\{0, b\}$. But this contradicts the fact that $\Omega^A F \subseteq \Omega^A G$, as desired. \square

We denote by Proto the Leibniz class of protoalgebraic logics, and by Asrt that of assertional logics. Bearing this in mind, we obtain the following result, in which suprema are taken in the Leibniz hierarchy.

Theorem 6.3. *The Leibniz class of protoalgebraic logics is meet-reducible among logics with theorems, and can be described as follows:*

$$(\text{Proto} \vee \text{Asrt}) \cap (\text{Proto} \vee \text{Log}(\vdash_{\text{BA}}^*)).$$

Proof. We set

$$\mathbb{K}_1 := \text{Proto} \vee \text{Asrt} \quad \text{and} \quad \mathbb{K}_2 := \text{Proto} \vee \text{Log}(\vdash_{\text{BA}}^*).$$

Observe that \mathbb{K}_1 and \mathbb{K}_2 are Leibniz classes by Theorems 3.4 and 3.11. Moreover, they comprise only logics with theorems by Propositions 2.6 and 3.15. We have

$$\text{Proto} \subsetneq \mathbb{K}_1 \quad \text{and} \quad \text{Equiv} \subsetneq \mathbb{K}_2. \quad (15)$$

The validity of the inclusion $\text{Proto} \subseteq \mathbb{K}_1 \cap \mathbb{K}_2$ is straightforward. The fact that $\text{Proto} \subsetneq \mathbb{K}_1$ is witnessed by the existence of assertional logics that are not protoalgebraic [32, Ex. 7]. Finally, the fact that the inclusion $\text{Equiv} \subseteq \mathbb{K}_2$ is strict follows from Lemma 6.2.

In the light of (15), it only remains to prove that $\mathbb{K}_1 \cap \mathbb{K}_2 \subseteq \text{Proto}$. Suppose the contrary, with a view to contradiction. Then there is a logic $\vdash \in \mathbb{K}_1 \cap \mathbb{K}_2$ that is not protoalgebraic. By Proposition 2.6, and considering that the non-indexed product of two protoalgebraic logics is protoalgebraic and interpretable in each of them, there are a protoalgebraic logic \vdash_{pr} and an assertional logic \vdash_{as} such that

$$\vdash_{pr} \otimes \vdash_{as} \leq \vdash \quad \text{and} \quad \vdash_{pr} \otimes \vdash_{BA}^* \leq \vdash. \quad (16)$$

Since \vdash has theorems and is not protoalgebraic, we can apply Theorem 3.1(ii) obtaining that $\text{Mod}^\equiv(\vdash) \neq \mathbb{R}(\text{Mod}^\equiv(\vdash))$, i.e., that there is a matrix $\langle A, F \rangle \in \text{Mod}^\equiv(\vdash)$ such that $\Omega^A F$ is not the identity relation.

We claim that for every pair of different $a, c \in A$, if $\langle a, c \rangle \in \Omega^A F$, then $a, c \notin F$. To prove this, consider different $a, c \in A$ such that $\langle a, c \rangle \in \Omega^A F$. By (16) there is an interpretation τ of $\vdash_{pr} \otimes \vdash_{as}$ into \vdash . By Proposition 6.1 we obtain without loss of generality that⁴

$$\langle A^\tau, F \rangle = \langle A_1 \otimes A_2, F_1 \times F_2 \rangle$$

for some $\langle A_1, F_1 \rangle \in \text{Mod}^\equiv(\vdash_{pr})$ and $\langle A_2, F_2 \rangle \in \text{Mod}^\equiv(\vdash_{as})$. Since $a, c \in A = A_1 \times A_2$, there are $a_1, c_1 \in A_1$ and $a_2, c_2 \in A_2$ such that

$$a = \langle a_1, a_2 \rangle \quad \text{and} \quad c = \langle c_1, c_2 \rangle.$$

From Proposition 6.1(ii) it follows that there are $\theta_1 \in \text{Con}A_1$ and $\theta_2 \in \text{Con}A_2$ compatible with F_1 and F_2 , respectively, such that

$$\Omega^A F = \{ \langle \langle p, q \rangle, \langle r, s \rangle \rangle : \langle p, r \rangle \in \theta_1 \text{ and } \langle q, s \rangle \in \theta_2 \}. \quad (17)$$

Since \vdash_{pr} is protoalgebraic and $\langle A_1, F_1 \rangle \in \text{Mod}^\equiv(\vdash_{pr})$, we can apply Theorem 3.1(ii) obtaining that $\Omega^{A_1} F_1$ is the identity relation. Together with the fact that $\theta_1 \subseteq \Omega^{A_1} F_1$ and (17), this yields

$$\Omega^A F = \{ \langle \langle p, q \rangle, \langle p, s \rangle \rangle : p \in A_1 \text{ and } \langle q, s \rangle \in \theta_2 \}. \quad (18)$$

Since $\langle a, c \rangle \in \Omega^A F$ and $a \neq c$, we conclude that $a_2 \neq c_2$ and $\langle a_2, c_2 \rangle \in \theta_2$.

Now, from the fact that θ_2 is compatible with F_2 and $\langle a_2, c_2 \rangle \in \theta_2$, it follows

$$\text{either } a_2, c_2 \in F_2 \text{ or } a_2, c_2 \notin F_2. \quad (19)$$

Since \vdash_{as} is assertional and $\langle A_2, F_2 \rangle \in \text{Mod}^\equiv(\vdash_{as})$, we know that F_2 is a singleton. Together with $a_2 \neq c_2$ and (19), this yields that $a_2, c_2 \notin F_2$. As a consequence, we obtain $\langle a, c \rangle \notin F_1 \times F_2 = F$, establishing the claim.

Now, by (16) there is an interpretation τ of $\vdash_{pr} \otimes \vdash_{BA}^*$ into \vdash . By Proposition 6.1 we obtain without loss of generality that

$$\langle A^\tau, F \rangle = \langle A_1 \otimes A_2, F_1 \times F_2 \rangle$$

for some $\langle A_1, F_1 \rangle \in \text{Mod}^\equiv(\vdash_{pr})$ and $\langle A_2, F_2 \rangle \in \text{Mod}^\equiv(\vdash_{BA}^*)$.

Recall that the matrix $\langle A, F \rangle$ is not reduced. Then there are different $a, c \in A$ such that $\langle a, c \rangle \in \Omega^A F$. Since $A = A_1 \times A_2$, there are $a_1, c_1 \in A_1$ and $a_2, c_2 \in A_2$ such that

$$a = \langle a_1, a_2 \rangle \quad \text{and} \quad c = \langle c_1, c_2 \rangle.$$

⁴For the sake of simplicity we assume that the map f in the statement of Proposition 6.1 is the identity. This assumption will be used systematically in this section without further notice.

As in the proof of the claim, we obtain

$$\Omega^A F = \{ \langle \langle p, q \rangle, \langle p, s \rangle \rangle : p \in A_1 \text{ and } \langle q, s \rangle \in \theta \} \quad (20)$$

for some $\theta \in \text{Con}A_2$ compatible with F_2 . Then we choose elements $e_1 \in F_1$ and $e_2 \in F_2$ (the fact that the logics \vdash_{pr} and \vdash_{BA}^* have theorems guarantees $F_1, F_2 \neq \emptyset$). Clearly we have

$$\langle e_1, e_2 \rangle \in F_1 \times F_2 = F. \quad (21)$$

Together with $a \neq c$ and $\langle a, c \rangle \in \Omega^A F$, the display (20) implies $a_2 \neq c_2$ and $\langle a_2, c_2 \rangle \in \theta$. Since $\langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_{BA}^*)$, we can apply Lemma 6.2 obtaining that A_2 is a Boolean algebra. Since A_2 is congruence uniform and θ identifies two distinct elements of A_2 (namely a_2 and c_2), then there is $b \in A_2 \setminus \{e_2\}$ such that $\langle e_2, b \rangle \in \theta$. By (20) we conclude

$$\langle \langle e_1, e_2 \rangle, \langle e_1, b \rangle \rangle \in \Omega^A F \text{ and } \langle e_1, e_2 \rangle \neq \langle e_1, b \rangle.$$

Together with (21) this contradicts the claim. This produces the desired contradiction. \square

6.2. Equivalential logics. To prove that the Leibniz class of equivalential logics is meet-reducible among logics with theorems, we need to introduce a new concept:

Definition 6.4. A formula $\varphi(x)$ is an *injective theorem* of a logic \vdash if $\varphi(x)$ is a theorem of \vdash and for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ the term-function $\varphi^A: A \rightarrow A$ is injective.

We will rely on the following:

Proposition 6.5. *Logics with an injective theorem form a Leibniz class, comprising a protoalgebraic non-equivalential logic.*

The proof of the above result proceeds through a series of technical observations.

Fact 1. Logics with an injective theorem form a Leibniz class.

Proof. It is not hard to show that logics with an injective theorem are closed under term-equivalence, compatible expansions, and non-indexed products of sets. In the light of Theorem 2.2 this implies that they form a Leibniz class. \square

Consider the logic \vdash_{∇} , formulated in countably many variables and in the language consisting of a single binary connective \rightarrow , axiomatized by the following rules:

$$\emptyset \triangleright x \rightarrow x \quad x, x \rightarrow y \triangleright y.$$

The logic \vdash_{∇} has been studied in depth in [15, 16, 17]. We set $\nabla(x, y) := \{x \rightarrow y\}$.

Fact 2. The logic \vdash_{∇} is protoalgebraic with set of congruence formulas with parameters

$$\hat{\nabla}(x, y, \vec{z}) := \{\varphi(x, \vec{z}) \rightarrow \varphi(y, \vec{z}) : \varphi(x, \vec{z}) \in \text{Fm}(\vdash)\}.$$

Proof. The logic \vdash_{∇} is protoalgebraic, since the set $\nabla(x, y)$ satisfies the requirements of Theorem 3.1(iv). By the same theorem, $\hat{\nabla}$ is a set of congruence formulas for \vdash_{∇} . \square

Fact 3. For every formula $\varphi \in \text{Fm}(\vdash_{\nabla})$,

$$\emptyset \vdash_{\nabla} \varphi \iff \varphi = \psi \rightarrow \psi, \text{ for some formula } \psi.$$

Proof. See [16, Prop. 2.1]. \square

Now, consider the extension \vdash_{Δ} of \vdash_{∇} , obtained adding for each $\psi \in \hat{\nabla}(x, y, \vec{z})$ the rule

$$\hat{\nabla}(x \rightarrow x, y \rightarrow y, \vec{z}) \triangleright \psi. \quad (22)$$

Fact 4. The logic \vdash_{Δ} is protoalgebraic.

Proof. This is a consequence of Fact 2, together with the fact that protoalgebraicity is preserved by extensions. \square

Fact 5. The formula $x \rightarrow x$ is an injective theorem of \vdash_{Δ} .

Proof. Clearly $x \rightarrow x$ is a theorem of \vdash_{Δ} . Then consider $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_{\Delta})$ and $a, b \in A$ such that $a \rightarrow^A a = b \rightarrow^A b$. We need to show that $a = b$. To this end, recall by Fact 4 that \vdash_{Δ} is protoalgebraic. Then we can apply Theorem 3.1(iii), obtaining that the matrix $\langle A, F \rangle$ is reduced, i.e., that the congruence $\Omega^A F$ is the identity relation. Therefore, to conclude the proof, it will be enough to show that $\langle a, b \rangle \in \Omega^A F$. By [23, Prop. 2.2(i)] this amounts to establishing that for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash_{\Delta})$ and every $\vec{c} \in A$,

$$\varphi^A(a, \vec{c}) \in F \iff \varphi^A(b, \vec{c}) \in F. \quad (23)$$

To prove the implication from left to right, consider $\varphi(x, \vec{z}) \in \text{Fm}(\vdash_{\Delta})$ and $\vec{c} \in A$ such that $\varphi^A(a, \vec{c}) \in F$. Consider also an arbitrary formula $\psi(x, \vec{z}) \in \text{Fm}(\vdash_{\nabla})$. We have that $\psi(x, \vec{z}) \rightarrow \psi(x, \vec{z})$ is a theorem of \vdash_{Δ} . Together with the fact that $a \rightarrow^A a = b \rightarrow^A b$, this yields

$$\psi^A(a \rightarrow a, \vec{c}) \rightarrow \psi^A(b \rightarrow b, \vec{c}) = \psi^A(a \rightarrow a, \vec{c}) \rightarrow \psi^A(a \rightarrow a, \vec{c}) \in F.$$

Hence we conclude that $\hat{\nabla}(a \rightarrow^A a, b \rightarrow^A b, \vec{c}) \subseteq F$. Since $\langle A, F \rangle$ is a model of (22), this implies $\hat{\nabla}(a, b, \vec{c}) \subseteq F$. As a consequence, we obtain $\varphi^A(a, \vec{c}) \rightarrow \varphi^A(b, \vec{c}) \in F$. Together with the assumption that $\varphi^A(a, \vec{c}) \in F$, and the fact that $\langle A, F \rangle$ is a model of the rule $x, x \rightarrow y \triangleright y$, this implies $\varphi^A(b, \vec{c}) \in F$. This concludes the proof of the left to right implication in (23). The proof of the other implication is analogous. From (23) it follows that $\langle a, b \rangle \in \Omega^A F$ and, therefore, $a = b$. \square

Now, recall that a rule $\Gamma \triangleright \varphi$ is *admissible* [35] in a logic \vdash , if its addition to \vdash does not produce new theorems. Equivalently, this means that $\emptyset \vdash \sigma\varphi$, for every substitution σ such that $\emptyset \vdash \sigma[\Gamma]$.

Fact 6. The rules in (22) are admissible in \vdash_{∇} .

Proof. To prove this, consider a substitution σ such that $\emptyset \vdash_{\nabla} \sigma[\hat{\nabla}(x \rightarrow x, y \rightarrow y, \vec{z})]$. Since $x \rightarrow y \in \hat{\nabla}(x, y, \vec{z})$, we have $(x \rightarrow x) \rightarrow (y \rightarrow y) \in \hat{\nabla}(x \rightarrow x, y \rightarrow y, \vec{z})$. As a consequence, we obtain $\emptyset \vdash_{\nabla} (\sigma x \rightarrow \sigma x) \rightarrow (\sigma y \rightarrow \sigma y)$. By Fact 3 this implies $\sigma x \rightarrow \sigma x = \sigma y \rightarrow \sigma y$ and, therefore, $\sigma x = \sigma y$. Since $\emptyset \vdash_{\nabla} x \rightarrow x$ and $\sigma x = \sigma y$, we obtain $\emptyset \vdash_{\nabla} \sigma\varphi(x, \vec{z}) \rightarrow \sigma\varphi(y, \vec{z})$, for every formula $\varphi(x, \vec{z}) \in \text{Fm}(\vdash_{\nabla})$. But this amounts to the fact that $\emptyset \vdash_{\nabla} \sigma[\hat{\nabla}(x, y, \vec{z})]$. Hence we conclude that the rules in (22) are admissible in \vdash_{∇} . \square

Given a formula φ , we denote by $\text{Var}(\varphi)$ the set of variables occurring in φ .

Fact 7. The logic \vdash_{Δ} is not equivalential.

Proof. Suppose, with a view to contradiction, that \vdash_{Δ} is equivalential. Then \vdash_{Δ} has a set of congruence formulas $\Delta(x, y)$.

We claim that $\emptyset \vdash_{\Delta} \varphi$, for every formula φ such that $\Delta(x, y) \vdash_{\Delta} \varphi$ and $\text{Var}(\varphi) \not\subseteq \{x, y\}$. To demonstrate this, we reason by complete induction on the length of proofs in \vdash_{Δ} . Consider an ordinal α , and suppose that $\emptyset \vdash_{\Delta} \psi$, for every formula ψ such that $\text{Var}(\psi) \not\subseteq \{x, y\}$, and of which there exists a proof indexed by an ordinal $< \alpha$ from $\Delta(x, y)$. Now, let

$\pi := \{\gamma_\beta: \beta < \alpha\}$ be a proof of a formula φ such that $\text{Var}(\varphi) \not\subseteq \{x, y\}$ from $\Delta(x, y)$. If φ is a substitution instance of the axiom $x \rightarrow x$, then it is a theorem and we are done. Moreover, observe that $\varphi \notin \Delta(x, y)$, since $\text{Var}(\varphi) \not\subseteq \{x, y\}$. Therefore, φ is obtained by the application of one of the inference rules of \vdash_Δ to a proper initial segment of π .

First consider the case where φ is obtained by an application of the rule $x, x \rightarrow y \triangleright y$. Then there is some formula ψ such that ψ and $\psi \rightarrow \varphi$ appear in a proper initial segment of π . Since $\text{Var}(\varphi) \not\subseteq \{x, y\}$, then $\text{Var}(\psi \rightarrow \varphi) \subseteq \{x, y\}$. Therefore we can apply the induction hypothesis obtaining $\emptyset \vdash_\Delta \psi \rightarrow \varphi$. By Fact 6 we get $\emptyset \vdash_\nabla \psi \rightarrow \varphi$. Moreover, by Fact 3 this yields $\psi = \varphi$. In particular, this implies $\text{Var}(\psi) \not\subseteq \{x, y\}$. Therefore we can apply the induction hypothesis, obtaining $\emptyset \vdash_\Delta \psi$. Since $\varphi = \psi$, we conclude that $\emptyset \vdash_\Delta \varphi$, as desired.

Then we consider the case where φ is obtained by an application of one of the rules in (22). Then there is a substitution σ and a formula $\psi(x, y, \vec{z}) \in \hat{\nabla}(x, y, \vec{z})$ such that $\varphi = \sigma\psi$, and each element of $\sigma[\hat{\nabla}(x \rightarrow x, y \rightarrow y, \vec{z})]$ appears in a proper initial segment of π . Suppose, with a view to contradiction, that $\emptyset \not\vdash_\Delta \sigma\psi$. Since $\psi \in \hat{\nabla}(x, y, \vec{z})$, there is a formula $\gamma(x, y, \vec{z}) \in \text{Fm}(\vdash_\nabla)$ such that $\psi = \gamma(x, y, \vec{z}) \rightarrow \gamma(y, y, \vec{z})$. Together with the fact that $\emptyset \vdash_\Delta x \rightarrow x$ and $\emptyset \not\vdash_\Delta \sigma\psi$, this yields $\sigma(\gamma(x, y, \vec{z})) \neq \sigma(\gamma(y, y, \vec{z}))$. But this easily implies $\sigma x \neq \sigma y$ and $x \in \text{Var}(\gamma(x, y, \vec{z}))$. As a consequence we obtain $\sigma(\gamma(x \rightarrow x, y \rightarrow y, \vec{z})) \neq \sigma(\gamma(y \rightarrow y, y \rightarrow y, \vec{z}))$. Moreover, from Facts 6 and 3, this yields

$$\emptyset \not\vdash_\Delta \sigma(\gamma(x \rightarrow x, y \rightarrow y, \vec{z})) \rightarrow \sigma(\gamma(y \rightarrow y, y \rightarrow y, \vec{z})),$$

which amounts to

$$\emptyset \not\vdash_\Delta \sigma\psi(x \rightarrow x, y \rightarrow y, \vec{z}), \text{ where } \psi = \psi(x, y, \vec{z}). \quad (24)$$

Now, from the fact that $\text{Var}(\sigma\psi) = \text{Var}(\varphi) \not\subseteq \{x, y\}$ it follows

$$\text{Var}(\sigma\psi(x \rightarrow x, y \rightarrow y, \vec{z})) \not\subseteq \{x, y\}.$$

The above display and the fact that $\sigma\psi(x \rightarrow x, y \rightarrow y, \vec{z}) \in \sigma[\hat{\nabla}(x \rightarrow x, y \rightarrow y, \vec{z})]$ allow us to apply the induction hypothesis, obtaining

$$\emptyset \vdash_\Delta \sigma\psi(x \rightarrow x, y \rightarrow y, \vec{z}).$$

But this contradicts (24). Hence we conclude that $\emptyset \vdash_\Delta \sigma\psi$, establishing the claim.

Now we move back to the main proof. First observe that

$$\Delta(x, y) \vdash_\Delta (x \rightarrow z) \rightarrow (y \rightarrow z). \quad (25)$$

To prove this, consider $\langle A, F \rangle \in \text{Mod}(\vdash_\Delta)$ and $a, b, c \in A$ such that $\Delta^A(a, b) \subseteq F$. Since Δ is a set of congruence formulas for \vdash_Δ , this yields $\langle a, b \rangle \in \Omega^A F$ and, therefore,

$$\langle (a \rightarrow c) \rightarrow (a \rightarrow c), (a \rightarrow c) \rightarrow (b \rightarrow c) \rangle \in \Omega^A F.$$

Since $x \rightarrow x$ is a theorem of \vdash_Δ , we have that $(a \rightarrow c) \rightarrow (a \rightarrow c) \in F$. As $\Omega^A F$ is compatible with F , this implies $(a \rightarrow c) \rightarrow (b \rightarrow c) \in F$, establishing (25). From (25) and the claim it follows that $\emptyset \vdash_\Delta (x \rightarrow z) \rightarrow (y \rightarrow z)$. By Facts 6 and 3 this yields $x \rightarrow z = y \rightarrow z$, which is false. \square

We also rely on the following result [18, Thm. 6.73]:

Theorem 6.6. *A protoalgebraic logic \vdash is equivalential if and only if $\text{Mod}^{\equiv}(\vdash)$ is closed under \mathbb{S} .*

We denote by Equiv and Injctv the Leibniz classes of equivalential logics and of logics with an injective theorem, respectively. Bearing this in mind, the main result of this part takes the following form:

Theorem 6.7. *The Leibniz class of equivalential logics is meet-reducible among logics with theorems, and can be described as follows:*

$$(\text{Equiv} \vee (\text{Proto} \cap \text{Injctv})) \cap (\text{Equiv} \vee (\text{Proto} \cap \text{Asrt})).$$

Proof. We begin by setting

$$\mathbb{K}_1 := \text{Equiv} \vee (\text{Proto} \cap \text{Injctv}) \text{ and } \mathbb{K}_2 := \text{Equiv} \vee (\text{Proto} \cap \text{Asrt}).$$

The fact that \mathbb{K}_1 and \mathbb{K}_2 are Leibniz classes is a consequence of Theorems 3.4, 3.7, and 3.11, and Proposition 6.5. Moreover, they comprise only logics with theorems by Propositions 2.6 and 3.15. We have

$$\text{Equiv} \subsetneq \mathbb{K}_1 \text{ and } \text{Equiv} \subsetneq \mathbb{K}_2. \quad (26)$$

The validity of the inclusion $\text{Equiv} \subseteq \mathbb{K}_1 \cap \mathbb{K}_2$ is straightforward. Moreover, Proposition 6.5 guarantees that $\text{Equiv} \subsetneq \mathbb{K}_1$. Finally, the fact that the inclusion $\text{Equiv} \subseteq \mathbb{K}_2$ is strict follows from the following observation [13, Prop. 6.1 and Thm. 6.4]:

$$(\text{Proto} \cap \text{Asrt}) \setminus \text{Equiv} \neq \emptyset.$$

In the light of (26), it only remains to prove $\mathbb{K}_1 \cap \mathbb{K}_2 \subseteq \text{Equiv}$. Suppose the contrary, with a view to contradiction. Then there is a logic $\vdash \in \mathbb{K}_1 \cap \mathbb{K}_2$ that is not equivalential. By Proposition 2.6 there are an equivalential logic \vdash_{eq} , a protoalgebraic logic with an injective theorem \vdash_{pin} , and an assertional protoalgebraic logic \vdash_{ap} such that

$$\vdash_{eq} \otimes \vdash_{pin} \leq \vdash \text{ and } \vdash_{eq} \otimes \vdash_{ap} \leq \vdash. \quad (27)$$

Together with [23, Prop. 3.8] this implies that \vdash is term-equivalent to a compatible expansion of the non-indexed product of a pair of protoalgebraic logics. Since protoalgebraic logics form a Leibniz class, by Theorem 2.2(ii) we conclude that \vdash is protoalgebraic. Moreover, since \vdash is not equivalential, we can apply Theorem 6.6 obtaining that $\text{Mod}^{\equiv}(\vdash)$ is not closed under \mathbb{S} , i.e., there is a matrix $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ with a submatrix $\langle \mathbf{B}, F \cap B \rangle \subseteq \langle \mathbf{A}, F \rangle$ such that $\langle \mathbf{B}, F \cap B \rangle \notin \text{Mod}^{\equiv}(\vdash)$.

We claim that for every pair of different $a, c \in B$, if $\langle a, c \rangle \in \Omega^B(F \cap B)$, then $a, b \notin F$. To prove this, consider different $a, c \in B$ such that $\langle a, c \rangle \in \Omega^B(F \cap B)$. By (28) there is an interpretation τ of $\vdash_{eq} \otimes \vdash_{ap}$ into \vdash . By Proposition 6.1(i) we obtain without loss of generality that

$$\begin{aligned} \langle \mathbf{A}^\tau, F \rangle &= \langle \mathbf{A}_1 \otimes \mathbf{A}_2, F_1 \times F_2 \rangle \\ \langle \mathbf{B}^\tau, F \cap B \rangle &= \langle \mathbf{B}_1 \otimes \mathbf{B}_2, (F_1 \cap B_1) \times (F_2 \cap B_2) \rangle \end{aligned}$$

for some $\langle \mathbf{B}_1, F_1 \rangle \subseteq \langle \mathbf{A}_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_{eq})$ and $\langle \mathbf{B}_2, F_2 \rangle \subseteq \langle \mathbf{A}_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_{pa})$. Since $a, c \in B = B_1 \times B_2$, there are $a_1, c_1 \in B_1$ and $a_2, c_2 \in B_2$ such that

$$a = \langle a_1, a_2 \rangle \text{ and } c = \langle c_1, c_2 \rangle.$$

Again from Proposition 6.1(ii) it follows that there are $\theta_1 \in \text{Con}B_1$ and $\theta_2 \in \text{Con}B_2$ compatible with $F_1 \cap B_1$ and $F_2 \cap B_2$, respectively, such that

$$\Omega^B(F \cap B) = \{ \langle \langle p, q \rangle, \langle r, s \rangle \rangle : \langle p, r \rangle \in \theta_1 \text{ and } \langle q, s \rangle \in \theta_2 \}.$$

Together with the fact that $\langle a, c \rangle \in \Omega^B(F \cap B)$, this yields

$$\langle a_i, c_i \rangle \in \theta_i \subseteq \Omega^{B_i}(F_i \cap B_i), \text{ for every } i = 1, 2. \quad (28)$$

Since \vdash_{eq} is equivalential, we can apply Theorem 6.6 obtaining that $\text{Mod}^{\equiv}(\vdash_{eq})$ is closed under \mathbb{S} . Together with $\langle A_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_{eq})$, this implies $\langle B_1, F_1 \cap B_1 \rangle \in \text{Mod}^{\equiv}(\vdash_{eq})$. Now, recall from Theorem 3.1(ii) that $\text{Mod}^{\equiv}(\vdash_{eq}) = \mathbb{R}(\text{Mod}(\vdash_{eq}))$. As a consequence, the congruence $\Omega^{B_1}(F_1 \cap B_1)$ is the identity relation. Hence by (28) we get $a_1 = c_1$. Since $a \neq c$, we conclude that $a_2 \neq c_2$.

Now, from (28) and the fact that $\Omega^{B_2}(F_2 \cap B_2)$ is compatible with $F_2 \cap B_2$ it follows that

$$\text{either } a_2, c_2 \in F_2 \cap B_2 \text{ or } a_2, c_2 \notin F_2 \cap B_2. \quad (29)$$

Since \vdash_{pa} is assertional and $\langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_{pa})$, we know that F_2 is a singleton. Together with $a_2 \neq c_2$ and (29), this yields $a_2, c_2 \notin F_2 \cap B_2$. As a consequence, we obtain $\langle a, c \rangle \notin F_1 \times F_2 = F$, establishing the claim.

Now, by (28) there is an interpretation τ of $\vdash_{eq} \otimes \vdash_{pin}$ into \vdash . By Proposition 6.1(i) we obtain without loss of generality that

$$\begin{aligned} \langle A^\tau, F \rangle &= \langle A_1 \otimes A_2, F_1 \times F_2 \rangle \\ \langle B^\tau, F \cap B \rangle &= \langle B_1 \otimes B_2, (F_1 \cap B_1) \times (F_2 \cap B_2) \rangle \end{aligned}$$

for some $\langle B_1, F_1 \rangle \subseteq \langle A_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_{eq})$ and $\langle B_2, F_2 \rangle \subseteq \langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_{pin})$.

Recall that the matrix $\langle B, F \cap B \rangle$ is not reduced. Then there are different $a, c \in B$ such that $\langle a, c \rangle \in \Omega^B(F \cap B)$. Since $B = B_1 \times B_2$, there are $a_1, c_1 \in B_1$ and $a_2, c_2 \in B_2$ such that

$$a = \langle a_1, a_2 \rangle \text{ and } c = \langle c_1, c_2 \rangle.$$

As in the proof of the claim, we obtain $a_2 \neq c_2$ and $\langle a_2, c_2 \rangle \in \Omega^{B_2}(F_2 \cap B_2)$.

Now, let $\varphi(x)$ be an injective theorem of \vdash_{pin} . Since $\langle A_2, F_2 \rangle \in \text{Mod}^{\equiv}(\vdash_{pin})$ and $a_2 \neq c_2$, we have

$$\varphi^{A_2}(a_2) \neq \varphi^{A_1}(a_2) \text{ and } \varphi^{A_2}(a_2), \varphi^{A_2}(c_2) \in F_2. \quad (30)$$

Let also $\top(x)$ be an arbitrary theorem of \vdash_{eq} . Since $\langle A_1, F_1 \rangle \in \text{Mod}(\vdash_{eq})$, we have

$$\top^{A_1}(a_1), \top^{A_1}(c_1) \in F_1. \quad (31)$$

Observe that the pair $\langle \top, \varphi \rangle$ is a unary connective of $\vdash_{eq} \otimes \vdash_{pld}$. Together with (30, 31), this yields

$$\begin{aligned} \langle \top, \varphi \rangle^{B_1 \otimes B_2}(a) &= \langle \top^{A_1}(a_1), \varphi^{A_2}(a_2) \rangle \in F_1 \times F_2 = F \\ \langle \top, \varphi \rangle^{B_1 \otimes B_2}(c) &= \langle \top^{A_1}(c_1), \varphi^{A_2}(c_2) \rangle \in F_1 \times F_2 = F \end{aligned}$$

and

$$\langle \top, \varphi \rangle^{B_1 \otimes B_2}(a) \neq \langle \top, \varphi \rangle^{B_1 \otimes B_2}(c).$$

Since $B_1 \otimes B_2 = B^\tau$, we know that $\langle \top, \varphi \rangle^{B_1 \otimes B_2}$ is a term-function of B . Together with the above displays and the fact that $\langle a, c \rangle \in \Omega^B(F \cap B)$, this implies that $\Omega^B(F \cap B)$ identifies two different elements of F , i.e., $\langle \top, \varphi \rangle^{B_1 \otimes B_2}(a)$ and $\langle \top, \varphi \rangle^{B_1 \otimes B_2}(c)$. But this contradicts the claim. Hence we reached a contradiction, as desired. \square

6.3. Order algebraizable logics. We denote by Order and Truth the Leibniz classes of order algebraizable and truth-equational logics, respectively.

Theorem 6.8. *The Leibniz class of order algebraizable logics is meet-reducible among logics with theorems, and can be described as follows:*

$$(\text{Order} \vee \text{Truth}) \cap \text{Equiv}.$$

Proof. First we set $\mathbb{K} := \text{Order} \vee \text{Truth}$. Observe that \mathbb{K} and Equiv are Leibniz classes by Theorems 3.7, 3.12, and 3.13. The fact that they comprise only logics with theorems is a consequence of Propositions 2.6 and 3.15. Moreover, we have

$$\text{Order} \subsetneq \mathbb{K} \quad \text{and} \quad \text{Order} \subsetneq \text{Equiv}. \quad (32)$$

To prove this, recall that every order algebraizable logic is equivalential [33, Prop. 7.11(iii)]. In particular, this implies that $\text{Order} \subseteq \mathbb{K} \cap \text{Equiv}$. The fact that the inclusion $\text{Order} \subseteq \mathbb{K}$ is strict is an immediate consequence of the fact that so is the inclusion $\text{Equiv} \subseteq \mathbb{K}_2$ in the proof of Theorem 6.7. On the other hand, the fact that $\text{Order} \subsetneq \text{Equiv}$ is witnessed by the existence of equivalential logics that are not order algebraizable [33, p. 267].

In the light of (32), it only remains to prove that $\mathbb{K} \cap \text{Equiv} \subseteq \text{Order}$. To this end, consider a logic $\vdash \in \mathbb{K} \cap \text{Equiv}$. Clearly, \vdash is equivalential and, therefore, there is a set of formulas $\Delta(x, y)$ of \vdash such that for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ and $a, c \in A$,

$$a = c \iff \Delta^A(a, c) \subseteq F. \quad (33)$$

Moreover, since $\vdash \in \mathbb{K}$, there are an order algebraizable logic \vdash_{or} , a truth-equational logic \vdash_{tr} , and an interpretation τ of $\vdash_{or} \otimes \vdash_{tr}$ into \vdash . Then there are a set of formulas $\nabla(x, y)$ and a set of inequalities $I(x)$ of \vdash_{or} such that for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_{or})$,

(i) the relation \preceq_F^A on A defined for every $a, c \in A$ by

$$a \preceq_F^A c \iff \nabla(a, c)^A \subseteq F$$

is a partial order; and

(ii) for every $a \in A$, $a \in F$ if and only if $\langle A, \preceq_F^A \rangle \models I(a)$.

Finally, since \vdash_{tr} is truth-equational, there is a set of equations $E(x)$ of \vdash_{tr} such that for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_{tr})$ and $a \in A$

$$a \in F \iff A \models E(a). \quad (34)$$

Now, we choose a theorem \top of \vdash_{tr} , and for every formula $\varphi \in \nabla(x, y)$ we consider the following basic operations of $\vdash_{or} \otimes \vdash_{tr}$, in which π_i is the projection map on the i -th coordinate:

$$\begin{aligned} \hat{\varphi}(x, y) &:= \langle \varphi(x_1, x_2), \top(x_1) \rangle \\ x \multimap y &:= \langle \pi_1(x_1, x_2), \pi_2(x_1, x_2) \rangle. \end{aligned}$$

Observe that for every $\mathcal{L}_{\vdash_{or}}$ -algebra A_1 , $\mathcal{L}_{\vdash_{tr}}$ -algebra A_2 , and elements $\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle \in A_1 \times A_2$, we have

$$\begin{aligned} \hat{\varphi}^{A_1 \otimes A_2}(\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle) &= \langle \varphi^{A_1}(a_1, c_1), \top^{A_2}(a_2) \rangle \\ \langle a_1, a_2 \rangle \multimap^{A_1 \otimes A_2} \langle c_1, c_2 \rangle &= \langle a_1, c_2 \rangle. \end{aligned} \quad (35)$$

Then we define the following sets of formulas and inequalities of \vdash :

$$\begin{aligned}\nabla^*(x, y) &:= \{\tau(\hat{\varphi}) : \varphi \in \nabla\} \cup \Delta(x, \tau(x \multimap y)) \\ I^*(x) &:= \{\tau(\langle \varepsilon(x_1), \zeta(x_1) \rangle) \preceq \tau(\langle \delta(x_1), \gamma(x_1) \rangle) : \varepsilon \preceq \delta \in I \text{ and } \zeta \approx \gamma \in E\}.\end{aligned}$$

To conclude the proof, it will be enough to show that the sets ∇^* and I^* witness the order algebraizability of \vdash . To this end, consider a matrix $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash)$, and let \preceq_F^A be the relation on A defined for every $a, c \in A$ as

$$a \preceq_F^A c \iff \nabla^*(a, c)^A \subseteq F.$$

We need to show that \preceq_F^A is a partial order on A and that for every $a \in A$,

$$a \in F \iff \langle A, \preceq_F^A \rangle \models I^*(a). \quad (36)$$

We claim that for every $\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle \in A_1 \times A_2 = A$,

$$\langle a_1, a_2 \rangle \preceq_F^A \langle c_1, c_2 \rangle \iff a_1 \preceq_{F_1}^{A_1} c_1 \text{ and } a_2 = c_2.$$

To prove this, observe that

$$\begin{aligned}\langle a_1, a_2 \rangle \preceq_F^A \langle c_1, c_2 \rangle &\iff \nabla^*(\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle)^A \subseteq F \\ &\iff \{\tau(\hat{\varphi})^A(\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle) : \varphi \in \nabla\} \subseteq F \text{ and} \\ &\quad \Delta^A(\langle a_1, a_2 \rangle, \tau(\langle a_1, a_2 \rangle \multimap \langle c_1, c_2 \rangle)) \subseteq F \\ &\iff \{\hat{\varphi}^{A_1 \otimes A_2}(\langle a_1, a_2 \rangle, \langle c_1, c_2 \rangle) : \varphi \in \nabla\} \subseteq F \text{ and} \\ &\quad \Delta^A(\langle a_1, a_2 \rangle, \langle a_1, a_2 \rangle \multimap^{A_1 \otimes A_2} \langle c_1, c_2 \rangle) \subseteq F \\ &\iff \{\langle \varphi^{A_1}(a_1, c_1), \top^{A_2}(a_2) \rangle : \varphi \in \nabla\} \subseteq F \text{ and} \\ &\quad \Delta^A(\langle a_1, a_2 \rangle, \langle a_1, c_2 \rangle) \subseteq F \\ &\iff \nabla^{A_1}(a_1, c_1) \subseteq F_1 \text{ and } \langle a_1, a_2 \rangle = \langle a_1, c_2 \rangle \\ &\iff a_1 \preceq_{F_1}^{A_1} c_1 \text{ and } a_2 = c_2.\end{aligned}$$

The above equivalences are justified as follows: the first, the second, and the sixth are straightforward, the third follows from the fact that $A^\tau = A_1 \otimes A_2$, the fourth is a consequence of (35), and the fifth follows from (33) and the observation that $F = F_1 \times F_2$ and $\top^{A_2}(a_2) \in F_2$. This establishes the claim.

Recall that $\preceq_{F_1}^{A_1}$ is a partial order on A_1 by (i). Together with the claim, this implies that \preceq_F^A is a partial order on A . Then consider an element $\langle a_1, a_2 \rangle \in A_1 \times A_2 = A$. We have

$$\begin{aligned}\langle a_1, a_2 \rangle \in F &\iff a_1 \in F_1 \text{ and } a_2 \in F_2 \\ &\iff \langle A_1^\tau, \preceq_{F_1}^{A_1} \rangle \models I(a_1) \text{ and } A_2^\tau \models E(a_2) \\ &\iff \langle A, \preceq_F^A \rangle \models I^*(\langle a_1, a_2 \rangle).\end{aligned}$$

The above equivalences are justified as follows: the first is a consequence of the equality $F = F_1 \times F_2$, the second follows from (ii) and (34), and the third from the claim. This establishes (36). Hence we conclude that \vdash is order algebraizable. \square

6.4. Parametrically truth-equational logics. As we mentioned, parametrically truth-equational logics need not have theorems in general [26, Sec. 4]. In particular, they lie outside the scope of Proposition 4.3 and we cannot immediately infer that their Leibniz class is meet-reducible in the absolute sense. We proceed to prove that this is indeed the case. To this end, we need the following observation:

Proposition 6.9. *Every parametrically truth-equational logic is truth-minimal, but the converse does not hold in general.*

Proof. Let \vdash be a parametrically truth-equational logic. Then consider two matrices $\langle A, F \rangle, \langle A, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \neq G \subseteq F$. Let also $E(x, \vec{y})$ be the set of equations that witnesses the fact that \vdash is parametrically truth-equational. Since $\langle A, F \rangle, \langle A, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ and $F, G \neq \emptyset$, for every $a \in A$ we have

$$a \in F \iff A \models E(a, \vec{c}) \text{ for every } \vec{c} \in A \iff a \in G.$$

As a consequence we obtain $F = G$ and, therefore, that \vdash is truth-minimal. This establishes that every parametrically truth-equational logic is truth-minimal.

To conclude the proof, we need to exhibit a truth-minimal logic that is not parametrically truth-equational. To this end, let \vdash be the logic formulated in countably many variables induced by the set of matrices $\{\langle B_2, \{1\} \rangle, \langle B_2, \{0\} \rangle\}$, where B_2 is the two-element Boolean algebra with universe $\{0, 1\}$. By Proposition 5.2 the logic \vdash is truth-minimal. Now, since B_2 is a two-element algebra, it is immediate that the matrices $\langle B_2, \{1\} \rangle$ and $\langle B_2, \{0\} \rangle$ are reduced. As a consequence, we obtain

$$\langle B_2, \{1\} \rangle, \langle B_2, \{0\} \rangle \in \mathbb{R}(\text{Mod}(\vdash)) \subseteq \text{Mod}^{\equiv}(\vdash).$$

Suppose, with a view to contradiction, that \vdash is parametrically truth-equational, and let $E(x, \vec{y})$ be the set of equations witnessing this fact. We have

$$\begin{aligned} 1 \in \{1\} &\implies B_2 \models E(1, \vec{c}), \text{ for every } \vec{c} \in B_2 \\ &\implies 1 \in \{0\} \\ &\implies 0 = 1. \end{aligned}$$

The first implication above follows from the fact that $\langle B_2, \{1\} \rangle \in \text{Mod}^{\equiv}(\vdash)$, the second from $\langle B_2, \{0\} \rangle \in \text{Mod}^{\equiv}(\vdash)$, and the third is straightforward. Since $0 \neq 1$, this produces a contradiction. Hence we conclude that \vdash is not parametrically truth-equational. \square

Given a logic \vdash and an \mathcal{L}_\vdash -algebra A , we denote by $\mathcal{F}_{i_\vdash} A$ the set of deductive filters of \vdash on A . We build on the following characterization result.

Theorem 6.10. *A logic \vdash is parametrically truth-equational if and only if for every \mathcal{L}_\vdash -algebra A and every family $X \cup \{F\} \subseteq \mathcal{F}_{i_\vdash} A \setminus \{\emptyset\}$,*

$$\text{if } \bigcap \{\Omega^A G : G \in X\} \subseteq \Omega^A F, \text{ then } \bigcap X \subseteq F.$$

Proof. The result is essentially a re-working of an analogous characterization of truth-equational logics in [32]. For the details, see [26, Thm. 3.9]. \square

We denote by Thrms , ParTruth , and Mnml the Leibniz classes of logics with theorems, parametrically truth-equational logics, and truth-minimal logics respectively.

Theorem 6.11. *The Leibniz class of parametrically truth-equational logics is meet-reducible, and can be described as follows:*

$$(\text{Thrms} \vee \text{ParTruth}) \cap \text{Mnml}.$$

Proof. First we set $\mathbb{K} := \text{Thrms} \vee \text{ParTruth}$, and observe that \mathbb{K} and Mnml are Leibniz classes by Theorems 3.12, 3.13, and 5.4. Moreover, we have

$$\text{ParTruth} \subsetneq \mathbb{K} \quad \text{and} \quad \text{ParTruth} \subsetneq \text{Mnml}. \quad (37)$$

The validity of the inclusion $\text{ParTruth} \subseteq \mathbb{K}$ is straightforward. The fact that it is strict is witnessed by the existence of logics with theorems that are not parametrically truth-equational, e.g., [26, Ex. 7.5]. On the other hand, from Proposition 6.9 it follows that $\text{ParTruth} \subsetneq \text{Mnml}$.

In the light of (37), it only remains to prove that $\mathbb{K} \cap \text{Mnml} \subseteq \text{ParTruth}$. To this end, consider a logic $\vdash \in \mathbb{K} \cap \text{Mnml}$. Clearly, \vdash is truth-minimal, and there are a parametrically truth-equational logic \vdash_{pt} , and a logic with theorems \vdash_{thm} such that $\vdash_{pt} \otimes \vdash_{thm} \leq \vdash$. Then let τ be an interpretation of $\vdash_{pt} \otimes \vdash_{thm}$ into \vdash .

In order to establish that \vdash is parametrically truth-equational, we rely on Theorem 6.10. Consider an \mathcal{L}_\vdash -algebra on A , and a family $X \cup \{F\} \subseteq \mathcal{F}i_\vdash A \setminus \{\emptyset\}$ such that

$$\bigcap \{\Omega^A G : G \in X\} \subseteq \Omega^A F. \quad (38)$$

To conclude the proof it suffices to show that $\bigcap X \subseteq F$. If $\bigcap X = \emptyset$, we are done. Then we consider the case where $\bigcap X$ is non-empty. For the sake of readability, we set $H := \bigcap X$ and $\theta := \tilde{\Omega}_\vdash^A \bigcap X$.

From the fact that $\langle A/\theta, H/\theta \rangle \in \text{Mod}^\equiv(\vdash)$ and [1, Prop. 4.12], it follows that

$$\langle (A/\theta)^\tau, H/\theta \rangle = \langle A_1, H_1 \rangle \otimes \langle A_2, H_2 \rangle \quad (39)$$

for some $\langle A_1, H_1 \rangle \in \text{Mod}^\equiv(\vdash_{pt})$ and $\langle A_2, H_2 \rangle \in \text{Mod}^\equiv(\vdash_{thm})$ with $H_1, H_2 \neq \emptyset$. Moreover, by (38) we obtain

$$\theta \subseteq \bigcap \{\Omega^A G : G \in X\} \subseteq \Omega^A F.$$

As a consequence, θ is compatible with F and, therefore, $F/\theta \in \mathcal{F}i_\vdash(A/\theta)$. Together with the fact that τ is an interpretation of $\vdash_{pt} \otimes \vdash_{thm}$ into \vdash and [23, Prop. 3.3], this yields

$$F/\theta \in \mathcal{F}i_{\vdash_{pt} \otimes \vdash_{thm}}((A/\theta)^\tau) \text{ and } F/\theta \neq \emptyset.$$

With an application of Lemma 5.3 to the above display and (39), we conclude that

$$F/\theta = F_1 \times F_2 \text{ for some } F_1 \in \mathcal{F}i_{\vdash_{pt}} A_1 \setminus \{\emptyset\}, \text{ and } F_2 \in \mathcal{F}i_{\vdash_{thm}} A_2. \quad (40)$$

We claim that $H/\theta \cap F/\theta \neq \emptyset$. To prove this, consider the family $Y := \{G \in \mathcal{F}i_{\vdash_{pt}} A_1 : H_1 \subseteq G\}$. We have

$$\bigcap \{\Omega^{A_1} G : G \in Y\} = \tilde{\Omega}_{\vdash_{pt}}^{A_1} H_1 \subseteq \Omega^{A_1} F_1,$$

where the inclusion $\tilde{\Omega}_{\vdash_{pt}}^{A_1} H_1 \subseteq \Omega^{A_1} F_1$ follows from the fact that $\tilde{\Omega}_{\vdash_{pt}}^{A_1} H_1$ is the identity relation, because $\langle A_1, H_1 \rangle \in \text{Mod}^\equiv(\vdash_{pt})$. Since $Y \cup \{F_1\} \subseteq \mathcal{F}i_{\vdash_{pt}} A_1 \setminus \{\emptyset\}$, we apply the fact that \vdash_{pt} is parametrically truth-equational and Theorem 6.10, obtaining $H_1 = \bigcap Y \subseteq F_1$. In particular, this guarantees that $H_1 \cap F_1 \neq \emptyset$, since $H_1 \neq \emptyset$. Similarly, the fact that

\vdash_{thm} has theorems and $H_2, F_2 \in \mathcal{F}_{\vdash_{thm}}^i \mathbf{A}_2$ implies $H_2 \cap F_2 \neq \emptyset$. Thus for every $i = 1, 2$ there is $a_i \in H_i \cap F_i$. By (39, 40) we conclude

$$\langle a_1, a_2 \rangle \in (H_1 \times H_2) \cap (F_1 \times F_2) = H/\theta \cap F/\theta,$$

establishing the claim.

Now, observe that the intersection $Z := H/\theta \cap F/\theta$ is clearly a deductive filter of \vdash on \mathbf{A}/θ . Together with the fact that $Z \subseteq H/\theta$, this implies $\tilde{\Omega}_{\vdash}^{\mathbf{A}/\theta} Z \subseteq \tilde{\Omega}_{\vdash}^{\mathbf{A}/\theta}(H/\theta)$. Since the latter congruence is the identity, so is $\tilde{\Omega}_{\vdash}^{\mathbf{A}} Z$ and, therefore, $\langle \mathbf{A}/\theta, Z \rangle \in \text{Mod}^{\equiv}(\vdash)$. This fact, together with the claim and $Z \subseteq H/\theta$, implies

$$H/\theta = Z = H/\theta \cap F/\theta.$$

Since θ is compatible both with H and F , we conclude that $\bigcap X = H \subseteq F$. \square

7. MEET-PRIME LOGICS

Traditional abstract algebraic logic tends to attribute the status of fundamental concepts both to protoalgebraic and equivalential logics. Unfortunately, this intuition does not match the fact that, when regarded as Leibniz classes, protoalgebraic and equivalential logics happen to be meet-reducible in the Leibniz hierarchy (Theorems 6.3 and 6.7). With an eye towards softening this apparent incoherence, we shall explore a different sense in which a Leibniz class can be considered to capture a primitive or fundamental concept.

Definition 7.1. A logic \vdash is *meet-prime* when $\llbracket \vdash \rrbracket$ is meet-prime in Log .

A Leibniz class can then also be considered primitive or fundamental when it is induced by a Leibniz condition whose members are meet-prime logics. In this section we show that this is indeed the case for protoalgebraic, equivalential, and assertional logics.

It is convenient to start with the case of protoalgebraic logics. Recall that for every infinite cardinal κ , the basic protoalgebraic logic of rank κ is denoted by $\vdash_{\mathfrak{p}}^{\kappa}$ (Definition 3.2). Our aim is to prove the following:

Theorem 7.2. *For every infinite cardinal $\kappa > 0$, the logic $\vdash_{\mathfrak{p}}^{\kappa}$ is meet-prime.*

As a consequence, we obtain the desired result.

Corollary 7.3. *The class of protoalgebraic logics has the form $\text{Log}(\Phi)$ for some Leibniz condition Φ consisting of meet-prime logics.*

Proof. Immediate from Theorems 3.4 and 7.2. \square

The proof of Theorem 7.2 proceeds through a series of technical observations. Given a pair of infinite cardinals κ and ν , we let

$$\{\langle \mathbf{A}_j, F_j \rangle : j \in J_{\kappa\nu}\}$$

be the set of ν -generated matrices in $\text{Mod}^{\equiv}(\vdash_{\mathfrak{p}}^{\kappa})$ up to isomorphism. We can assume without loss of generality that the various algebras \mathbf{A}_j have disjoint universes. Then let $\lambda_{\kappa\nu} := \max\{\omega, |\bigcup_{j \in J_{\kappa\nu}} \mathbf{A}_j|\}$ and consider the sets

$$\begin{aligned} A_{\kappa\nu} &:= \{\top, \perp\} \cup \{p_{\alpha} : \alpha < \lambda_{\kappa\nu}^+\} \cup \bigcup_{j \in J_{\kappa\nu}} A_j \\ F_{\kappa\nu} &:= \{\top\} \cup \{p_{\alpha} : \alpha < \lambda_{\kappa\nu}^+\} \cup \bigcup_{j \in J_{\kappa\nu}} F_j \end{aligned}$$

where \perp, \top, p_α are new distinct elements. We endow $A_{\kappa\nu}$ with the structure of an \mathcal{L}_P^κ -algebra $A_{\kappa\nu}$ stipulating that for every $\alpha < \kappa, 0 < n \in \omega$, and $a, b, c_1, \dots, c_n \in A_{\kappa\nu}$,

$$a \multimap_\alpha^{A_{\kappa\nu}} b := \begin{cases} a \multimap_\alpha^{A_j} b & \text{if } a, b \in A_j \text{ for some } j \in J_{\kappa\nu} \\ \top & \text{if } a = b \text{ and } \{a, b\} \not\subseteq \bigcup_{j \in J_{\kappa\nu}} A_j \\ \perp & \text{if } a \neq b \text{ and } \{a, b\} \not\subseteq \bigcup_{j \in J_{\kappa\nu}} A_j \end{cases}$$

and

$$*_{n\alpha}^{A_{\kappa\nu}}(c_1, \dots, c_n) := \begin{cases} *_{n\alpha}^{A_j}(c_1, \dots, c_n) & \text{if } c_1, \dots, c_n \in A_j \text{ for some } j \in J_{\kappa\nu} \\ \perp & \text{otherwise.} \end{cases}$$

Observe that $A_{\kappa\nu}$ is well defined, since the various A_j have disjoint universes.

Fact 8. For every pair of infinite cardinals κ and ν , we have $|F_{\kappa\nu}| > |A_{\kappa\nu} \setminus F_{\kappa\nu}|$.

Proof. From the definition of $A_{\kappa\nu}$ and $F_{\kappa\nu}$ it follows that

$$|F_{\kappa\nu}| \geq \lambda_{\kappa\nu}^+ > \lambda_{\kappa\nu} = \max\{\omega, \bigcup_{j \in J_{\kappa\nu}} |A_j|\} \geq |A_{\kappa\nu} \setminus F_{\kappa\nu}|. \quad \square$$

Fact 9. For every pair of infinite cardinals κ and ν , and $j \in J_{\kappa\nu}$,

$$\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle \in \text{Mod}^{\equiv}(\vdash_P^\kappa) \quad \text{and} \quad \langle A_j, F_j \rangle \subseteq \langle A_{\kappa\nu}, F_{\kappa\nu} \rangle.$$

Proof. The fact that $\langle A_j, F_j \rangle \subseteq \langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$ is clear.

To establish that $\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle \in \text{Mod}^{\equiv}(\vdash_P^\kappa)$, it suffices to prove that the matrix $\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$ is a reduced model of \vdash_P^κ . The fact that it is reduced is justified as follows. Consider two distinct elements $a, b \in A_{\kappa\nu}$. We have to prove that $\langle a, b \rangle \notin \Omega^{A_{\kappa\nu}} F_{\kappa\nu}$. First we consider the case where there is $j \in J_{\kappa\nu}$ such that $a, b \in A_j$. Since \vdash_P^κ is protoalgebraic and $\langle A_j, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_P^\kappa)$, we can apply Theorem 3.1(ii), obtaining that the matrix $\langle A_j, F_j \rangle$ is reduced. In particular, we can assume without loss of generality that there is a unary polynomial function p of A_j such that $p(a) \in F_j$ and $p(b) \notin F_j$ [23, Prop. 2.2(i)]. Since $\langle A_j, F_j \rangle \subseteq \langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$, the map p is also a unary polynomial function of $A_{\kappa\nu}$ such that $p(a) \in F_{\kappa\nu}$ and $p(b) \notin F_{\kappa\nu}$. As a consequence, $\langle a, b \rangle \notin \Omega^{A_{\kappa\nu}} F_{\kappa\nu}$ by [23, Prop. 2.2(i)]. Then we consider the case where there is no $j \in J_{\kappa\nu}$ such that $a, b \in A_j$. Choose an arbitrary $\alpha < \kappa$, and consider the unary polynomial function $p(x) := x \multimap_\alpha^{A_{\kappa\nu}} a$ of $A_{\kappa\nu}$. Observe that

$$p(b) = a \multimap_\alpha^{A_{\kappa\nu}} b = \perp \notin F_{\kappa\nu}.$$

On the other hand, we shall see that $p(a) \in F_{\kappa\nu}$. If $a \in A_j$ for some $j \in J_{\kappa\nu}$, then we have

$$p(a) = a \multimap_\alpha^{A_{\kappa\nu}} a \in F_j \subseteq F_{\kappa\nu},$$

since $\langle A_j, F_j \rangle \in \text{Mod}(\vdash_P^\kappa)$ and $\emptyset \vdash_P^\kappa x \multimap_\alpha x$. Moreover, if $a \notin \bigcup_{j \in J_{\kappa\nu}} A_j$, then

$$p(a) = a \multimap_\alpha^{A_{\kappa\nu}} a = \top \in F_{\kappa\nu}.$$

Hence we conclude that $p(a) \in F_{\kappa\nu}$. Together with the fact that $p(b) \notin F_{\kappa\nu}$ and [23, Prop. 2.2(i)], this implies $\langle a, b \rangle \notin \Omega^{A_{\kappa\nu}} F_{\kappa\nu}$. We conclude that $\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$ is reduced.

It only remains to show that $\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle \in \text{Mod}(\vdash_P^\kappa)$, i.e., that $\langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$ is a model of the rules $\emptyset \triangleright \Delta_\kappa(x, x)$ and $x, \Delta_\kappa(x, y) \triangleright y$. We detail only the case of $x, \Delta_\kappa(x, y) \triangleright y$, since the other one is similar. Consider $a, b \in A_{\kappa\nu}$ such that $\{a\} \cup \Delta^{A_{\kappa\nu}}(a, b) \subseteq F_{\kappa\nu}$. First we consider the case where $b \in A_j$ for some $j \in J_{\kappa\nu}$. Looking at the definition of $A_{\kappa\nu}$, it is not hard to see that the fact that $b \in A_j$ and $\Delta^{A_{\kappa\nu}}(a, b) \subseteq F_{\kappa\nu}$ implies $a, b \in A_j$. In particular, this

guarantees that $\Delta^{A_{\kappa\nu}}(a, b) = \Delta^{A_j}(a, b)$, whence $\Delta^{A_j}(a, b) \subseteq A_j \cap F_{\kappa\nu} = F_j$. Together with the fact that $\langle A_j, F_j \rangle$ is a model of the rule $x, \Delta_\kappa(x, y) \triangleright y$, this yields $b \in F_j \subseteq F_{\kappa\nu}$. Then we consider the case where $b \notin \bigcup_{j \in J_{\kappa\nu}} A_j$. Again looking at the definition of $A_{\kappa\nu}$, it is not difficult to see that the fact that $b \notin \bigcup_{j \in J_{\kappa\nu}} A_j$ and $\Delta^{A_{\kappa\nu}}(a, b) \subseteq F_{\kappa\nu}$ implies $a = b$, whence $b = a \in F_{\kappa\nu}$. \square

Fact 10. Let κ and ν be infinite cardinals, \vdash_1 and \vdash_2 logics, and τ an interpretation of $\vdash_1 \otimes \vdash_2$ into $\vdash_{\mathfrak{p}}^\kappa$. Then for every $i = 1, 2$ there is $\langle B_i, G_i \rangle \in \text{Mod}^\equiv(\vdash_i)$ such that

$$\langle A_{\kappa\nu}^\tau, F_{\kappa\nu} \rangle \cong \langle B_1 \otimes B_2, G_1 \times G_2 \rangle.$$

Moreover, either $\langle B_1, G_1 \rangle$ or $\langle B_2, G_2 \rangle$ is trivial.

Proof. Since τ is an interpretation of $\vdash_1 \otimes \vdash_2$ into $\vdash_{\mathfrak{p}}^\kappa$, we can apply Fact 9 obtaining $\langle A_{\kappa\nu}^\tau, F_{\kappa\nu} \rangle \in \text{Mod}^\equiv(\vdash_1 \otimes \vdash_2)$. By [23, Cor. 4.14] for every $i = 1, 2$ there is $\langle B_i, G_i \rangle \in \text{Mod}^\equiv(\vdash_i)$ such that $\langle A_{\kappa\nu}^\tau, F_{\kappa\nu} \rangle \cong \langle B_1 \otimes B_2, G_1 \times G_2 \rangle$. For the sake of simplicity, we assume without loss of generality that

$$\langle A_{\kappa\nu}^\tau, F_{\kappa\nu} \rangle = \langle B_1 \otimes B_2, G_1 \times G_2 \rangle. \quad (41)$$

It only remains to prove that either $\langle B_1, G_1 \rangle$ or $\langle B_2, G_2 \rangle$ is trivial. Suppose the contrary, with a view to contradiction. We have

$$\begin{aligned} |B_1| + |B_2| &\leq |(B_1 \times (B_2 \setminus G_2)) \cup ((B_1 \setminus G_1) \times B_2)| \\ &= |(B_1 \times B_2) \setminus (G_1 \times G_2)| \\ &= |A_{\kappa\nu} \setminus F_{\kappa\nu}|. \end{aligned}$$

The first inequality above follows from the fact that, by Lemma 5.5, $G_i \subsetneq B_i$ for every $i = 1, 2$. The second one is obvious, and the third one follows from (41).

Now, recall that $|A_{\kappa\nu}| \geq \lambda_{\kappa\nu}^+ \geq \omega$. Thus the set $A_{\kappa\nu} = B_1 \times B_2$ is infinite, whence so is either B_1 or B_2 . In particular, this implies $|B_1| + |B_2| = |B_1 \times B_2| = |A_{\kappa\nu}|$. Together with the above display, this yields

$$|F_{\kappa\nu}| \leq |A_{\kappa\nu}| \leq |A_{\kappa\nu} \setminus F_{\kappa\nu}|.$$

But this is in contradiction with Fact 8. \square

Proof of Theorem 7.2. Consider an infinite cardinal κ . Our aim is to show that the logic $\vdash_{\mathfrak{p}}^\kappa$ is meet-prime. To this end, consider two logics \vdash_1 and \vdash_2 with an interpretation τ of $\vdash_1 \otimes \vdash_2$ into $\vdash_{\mathfrak{p}}^\kappa$. Then let τ_1 be the translation of \mathcal{L}_{\vdash_1} into $\mathcal{L}_{\mathfrak{p}}^\kappa$ defined for every n -ary $* \in \mathcal{L}_{\vdash_1}$ as

$$\tau_1(*) := \tau(\langle *(x_1, \dots, x_n), x_1 \rangle).$$

The above definition is sound, since the pair $\langle *(x_1, \dots, x_n), x_1 \rangle$ can be regarded as a basic n -ary operation of $\vdash_1 \otimes \vdash_2$. Let also τ_2 be the translation of \mathcal{L}_{\vdash_2} into $\mathcal{L}_{\mathfrak{p}}^\kappa$ defined analogously.

We claim that for every infinite cardinal ν , there is $i = 1, 2$ such that

$$\langle A_j^{\tau_i}, F_j \rangle \in \text{Mod}^\equiv(\vdash_i) \text{ for every } j \in J_{\kappa\nu}.$$

To prove this, consider an infinite cardinal ν . By Fact 10 we can assume without loss of generality that

$$\langle A_{\kappa\nu}^\tau, F_{\kappa\nu} \rangle = \langle A \otimes \mathbf{1}, F \times \{1\} \rangle$$

for some $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_1)$. We shall prove that $\langle A_j^{\tau_1}, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_1)$ for all $j \in J_{\kappa\nu}$. To this end, consider $j \in J_{\kappa\nu}$. By Fact 9 we obtain $\langle A_j, F_j \rangle \subseteq \langle A_{\kappa\nu}, F_{\kappa\nu} \rangle$, whence

$$\langle A_j^{\tau}, F_j \rangle \subseteq \langle A_{\kappa\nu}^{\tau}, F_{\kappa\nu} \rangle = \langle A \otimes \mathbf{1}, F \times \{1\} \rangle.$$

Thus there is $B \subseteq A$ such that

$$\langle A_j^{\tau}, F_j \rangle = \langle B \otimes \mathbf{1}, (F \cap B) \times \{1\} \rangle. \quad (42)$$

Observe that

$$\langle B, F \cap B \rangle \in \text{Mod}^{\equiv}(\vdash_1). \quad (43)$$

To prove this, observe that $\langle B, F \cap B \rangle \in \text{Mod}(\vdash_1)$, since $\langle B, F \cap B \rangle \subseteq \langle A, F \rangle$ and $\langle A, F \rangle \in \text{Mod}(\vdash_1)$. Hence it only remains to show that $\widetilde{\Omega}_{\vdash_1}^B F \cap B$ is the identity relation. Consider two distinct elements $a, c \in B$. Since $\langle A_j, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_{\mathfrak{p}}^{\kappa})$ and τ is an interpretation of $\vdash_1 \otimes \vdash_2$ into $\vdash_{\mathfrak{p}}^{\kappa}$, we have

$$\langle B \otimes \mathbf{1}, (F \cap B) \times \{1\} \rangle \in \text{Mod}^{\equiv}(\vdash_1 \otimes \vdash_2).$$

Since the elements $\langle a, 1 \rangle, \langle c, 1 \rangle \in B \times \{1\}$ are distinct, we can apply [23, Prop. 2.2(ii)] to the above display obtaining without loss of generality a set $F \cap B \subseteq G \subseteq B$ such that $G \times \{1\}$ is a deductive filter of $\vdash_1 \otimes \vdash_2$ on $B \otimes \mathbf{1}$, a pair $\langle \varphi(x, y_1, \dots, y_n), \psi(x, y_1, \dots, y_n) \rangle$ such that $\varphi \in \text{Fm}(\vdash_1)$ and $\psi \in \text{Fm}(\vdash_2)$, and elements $b_1, \dots, b_n \in B$ such that

$$\begin{aligned} \langle \varphi, \psi \rangle^{B \otimes \mathbf{1}}(\langle a, 1 \rangle, \langle b_1, 1 \rangle, \dots, \langle b_n, 1 \rangle) &\in G \times \{1\} \\ \langle \varphi, \psi \rangle^{B \otimes \mathbf{1}}(\langle c, 1 \rangle, \langle b_1, 1 \rangle, \dots, \langle b_n, 1 \rangle) &\notin G \times \{1\}. \end{aligned}$$

In particular, we have

$$\varphi^B(a, b_1, \dots, b_n) \in G \quad \text{and} \quad \varphi^B(c, b_1, \dots, b_n) \notin G.$$

Observe that G is a deductive filter of \vdash_1 on B by Lemma 5.3. Together with the fact that $F \cap B \subseteq G$ and the above display, this allows us to apply [23, Prop. 2.2(ii)] yielding $\langle a, c \rangle \notin \widetilde{\Omega}_{\vdash_1}^B(F \cap B)$. This concludes the proof that $\widetilde{\Omega}_{\vdash_1}^B(F \cap B)$ is the identity relation and establishes (43).

Now, recall from (42) that $A_j = B \times \{1\}$. Then let $\pi: A_j \rightarrow B$ be the projection on the first coordinate. The fact that π a bijection between A_j and B such that $\pi[F_j] = F \cap B$ and $F_j = \pi^{-1}[F \cap B]$ is a direct consequence of (42). Moreover, it is easy to show that π is a homomorphism. Therefore, π is an isomorphism from $\langle A_j^{\tau_1}, F_j \rangle$ to $\langle B, F \cap B \rangle$. Together with (43) this yields $\langle A_j^{\tau_1}, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_1)$, establishing the claim.

From the claim it follows that there is $i = 1, 2$ such that for every infinite cardinal ν there is a cardinal $\mu \geq \nu$ such that $\langle A_j^{\tau_i}, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ for all $j \in J_{\kappa\mu}$. Bearing in mind that if ν and μ are infinite cardinals such that $\nu \leq \mu$, then $\{\langle A_j, F_j \rangle : j \in J_{\kappa\nu}\} \subseteq \{\langle A_j, F_j \rangle : j \in J_{\kappa\mu}\}$, this yields $\langle A_j^{\tau_i}, F_j \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ for every infinite cardinal ν and $j \in J_{\kappa\nu}$. Hence we have $\langle A^{\tau_i}, F \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ for every $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_{\mathfrak{p}}^{\kappa})$, whence τ_i is a translation of \vdash_i into $\vdash_{\mathfrak{p}}^{\kappa}$. We conclude that $\vdash_i \leq \vdash_{\mathfrak{p}}^{\kappa}$ and, therefore, that $\vdash_{\mathfrak{p}}^{\kappa}$ is a meet-prime logic. \square

Recall that for every infinite cardinal $\kappa > 0$, the basic equivalential logic of rank κ is denoted by $\vdash_{\mathfrak{F}}^{\kappa}$ (Definition 3.5). An argument, similar to the one described above, yields the following conclusion:

Theorem 7.4. *For every infinite cardinal $\kappa > 0$, the logic $\vdash_{\mathbb{E}}^{\kappa}$ is meet-prime. As a consequence, the class of equivoential logics has the form $\text{Log}(\Phi)$ for some Leibniz condition Φ consisting of meet-prime logics.*

Recall that the basic assertional logic is denoted by $\vdash_{\mathbb{A}}$ (Definition 3.9). We have the following:

Theorem 7.5. *The logic $\vdash_{\mathbb{A}}$ is meet-prime. As a consequence, the class of assertional logics has the form $\text{Log}(\Phi)$ for a strong Leibniz condition Φ consisting of a meet-prime logic.*

Proof. In the light to Theorem 3.11 it will be enough to show that $\vdash_{\mathbb{A}}$ is meet-prime. To this end, consider two logics \vdash_1 and \vdash_2 , and an interpretation τ of $\vdash_1 \otimes \vdash_2$ into $\vdash_{\mathbb{A}}$. Moreover, let $\mathbf{2}$ be the two-element pointed set. By Proposition 3.10 we have $\langle \mathbf{2}, \{\top_*^2\} \rangle \in \text{Mod}^{\equiv}(\vdash_{\mathbb{A}})$, whence $\langle \mathbf{2}^{\tau}, \{\top_*^2\} \rangle \in \text{Mod}^{\equiv}(\vdash_1 \otimes \vdash_2)$. Together with [23, Cor. 4.14], this implies that for every $i = 1, 2$ there is $\langle A_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i)$ such that $\langle \mathbf{2}^{\tau}, \{\top_*^2\} \rangle = \langle A_1 \otimes A_2, F_1 \times F_2 \rangle$.

Now, observe that $A_1 \times A_2$ is a two-element set. As a consequence, either A_1 or A_2 is a singleton. We can assume without loss of generality that so is A_2 . Together with the fact that $F_2 \neq \emptyset$, this implies that $\langle A_2, F_2 \rangle$ is the trivial matrix $\langle \mathbf{1}, \{1\} \rangle$. Thus

$$\langle \mathbf{2}^{\tau}, \{\top_*^2\} \rangle = \langle A_1 \otimes \mathbf{1}, F_1 \times \{1\} \rangle.$$

Consider the translation τ_1 of \mathcal{L}_{\vdash_1} into $\mathcal{L}_{\vdash_{\mathbb{A}}}$ defined as in the proof of Theorem 7.2. We shall prove that τ_1 is an interpretation of \vdash_1 into $\vdash_{\mathbb{A}}$. Making use of the above display, it is not hard to see that $\langle \mathbf{2}^{\tau_1}, \{\top_*^2\} \rangle \cong \langle A_1, F_1 \rangle$. Together with the fact that $\langle A_1, F_1 \rangle \in \text{Mod}^{\equiv}(\vdash_1)$, this yields $\langle \mathbf{2}^{\tau_1}, \{\top_*^2\} \rangle \in \text{Mod}^{\equiv}(\vdash_1)$. By [23, Lem. 2.3] we conclude

$$\mathbb{P}_{\text{SD}}(\langle \mathbf{2}^{\tau_1}, \{\top_*^2\} \rangle) \subseteq \text{Mod}^{\equiv}(\vdash_1). \quad (44)$$

To prove that τ_1 is an interpretation, consider a matrix $\langle A, F \rangle \in \text{Mod}^{\equiv}(\vdash_{\mathbb{A}})$. By Proposition 3.10 we know that A is a pointed set and $F = \{\top_*^A\}$. In particular, this easily implies $\langle A, F \rangle \in \mathbb{P}_{\text{SD}}(\langle \mathbf{2}, \{\top_*^2\} \rangle)$, whence $\langle A^{\tau_1}, F \rangle \in \mathbb{P}_{\text{SD}}(\langle \mathbf{2}^{\tau_1}, \{\top_*^2\} \rangle)$. By (44) this guarantees that $\langle A^{\tau_1}, F \rangle \in \text{Mod}^{\equiv}(\vdash_1)$. Hence we conclude that τ_1 is an interpretation of \vdash_1 into $\vdash_{\mathbb{A}}$, whence $\vdash_1 \leq \vdash_{\mathbb{A}}$. This shows that $\vdash_{\mathbb{A}}$ is meet-prime, as desired. \square

Acknowledgements. Thanks are due to James G. Raftery for rising the question about whether the theory of the Maltsev and Leibniz hierarchy could be, to some extent, unified. The authors were supported by the I+D+i research project PID2019-110843GA-I00 *La geometria de las logicas no-clasicas* funded by the Ministry of Science and Innovation of Spain. The second author was also supported by the *Beatriz Galindo* grant BEAGAL-18/00040 funded by the Ministry of Science and Innovation of Spain and by the grant CZ.02.2.69/0.0/0.0/17_050/0008361, OPVVV MŠMT, MSCA-IF *Lidské zdroje v teoretické informatice* funded by the Ministry of Education, Youth and Sports of the Czech Republic.

REFERENCES

- [1] H. Albuquerque, J. M. Font, R. Jansana, and T. Moraschini. Assertional logics, truth-equational logics, and the hierarchies of abstract algebraic logic. In J. Czelakowski, editor, *Don Pigozzi on Abstract Algebraic Logic and Universal Algebra*, volume 16 of *Outstanding Contributions*. Springer-Verlag, 2018.
- [2] W. Bentz and L. Sequeira. Taylor’s modularity conjecture holds for linear idempotent varieties. *Algebra Universalis*, 71(2):101–102, 2014.
- [3] C. Bergman. *Universal Algebra: Fundamentals and Selected Topics*. Chapman & Hall Pure and Applied Mathematics. Chapman and Hall/CRC, 2011.

- [4] G. Bezhanishvili, T. Moraschini, and J. G. Raftery. Epimorphisms in varieties of residuated structures. *Journal of Algebra*, 492:185–211, 2017.
- [5] W. J. Blok and E. Hoogland. The Beth property in Algebraic Logic. *Studia Logica*, 83(1–3):49–90, 2006.
- [6] W. J. Blok and P. Köhler. Algebraic semantics for quasi-classical modal logics. *The Journal of Symbolic Logic*, 48:941–964, 1983.
- [7] W. J. Blok and D. Pigozzi. Protoalgebraic logics. *Studia Logica*, 45:337–369, 1986.
- [8] W. J. Blok and D. Pigozzi. *Algebraizable logics*, volume 396 of *Mem. Amer. Math. Soc.* A.M.S., Providence, January 1989.
- [9] W. J. Blok and D. Pigozzi. Algebraic semantics for universal Horn logic without equality. In A. Romanowska and J. D. H. Smith, editors, *Universal Algebra and Quasigroup Theory*, pages 1–56. Heldermann, Berlin, 1992.
- [10] W. J. Blok and J. Rebagliato. Algebraic semantics for deductive systems. *Studia Logica, Special Issue on Abstract Algebraic Logic, Part II*, 74(5):153–180, 2003.
- [11] J. Czelakowski. Equivalential logics, I, II. *Studia Logica*, 40:227–236 and 355–372, 1981.
- [12] J. Czelakowski. *Protoalgebraic logics*, volume 10 of *Trends in Logic—Studia Logica Library*. Kluwer Academic Publishers, Dordrecht, 2001.
- [13] J. Czelakowski and R. Jansana. Weakly algebraizable logics. *The Journal of Symbolic Logic*, 65(2):641–668, 2000.
- [14] J. Czelakowski and D. Pigozzi. Amalgamation and interpolation in abstract algebraic logic. In X. Caicedo and Carlos H. Montenegro, editors, *Models, algebras and proofs*, number 203 in *Lecture Notes in Pure and Applied Mathematics Series*, pages 187–265. Marcel Dekker, New York and Basel, 1999.
- [15] J. M. Font. Atoms in the lattice of theories. *Bull. Sect. Logic*, 42(1–2):21–31, 2013.
- [16] J. M. Font. The simplest protoalgebraic logic. *Mathematical Logic Quarterly*, 59(6):435–451, 2013.
- [17] J. M. Font. Ordering protoalgebraic logics. *Journal of Logic and Computation*, 26(5):1395–1419, 2014.
- [18] J. M. Font. *Abstract Algebraic Logic - An Introductory Textbook*, volume 60 of *Studies in Logic - Mathematical Logic and Foundations*. College Publications, London, 2016.
- [19] J. M. Font and R. Jansana. *A general algebraic semantics for sentential logics*, volume 7 of *Lecture Notes in Logic*. A.S.L., second edition 2017 edition, 2009. First edition 1996. Electronic version freely available through Project Euclid at projecteuclid.org/euclid.lnl/1235416965.
- [20] J. M. Font, R. Jansana, and D. Pigozzi. A survey on abstract algebraic logic. *Studia Logica, Special Issue on Abstract Algebraic Logic, Part II*, 74(1–2):13–97, 2003. With an “Update” in 91 (2009), 125–130.
- [21] O. C. García and W. Taylor. *The lattice of interpretability types of varieties*, volume 50. *Mem. Amer. Math. Soc.*, 1984.
- [22] D. Hobby and R. McKenzie. *The structure of finite algebras*, volume 76 of *Contemporary Mathematics*. American Mathematical Society, Providence, RI, 1988.
- [23] R. Jansana and T. Moraschini. The poset of all logics I: interpretations and lattice structure. To appear in the *Journal of Symbolic Logic*, 2021.
- [24] R. Jansana and T. Moraschini. The poset of all logics III: finitely presentable logics. *Studia Logica*, published online, 2020.
- [25] K. A. Kearnes and E. W. Kiss. *The shape of congruences lattices*, volume 222 of *Mem. Amer. Math. Soc.* American Mathematical Society, 2013. Monograph.
- [26] T. Moraschini. A study of the truth predicates of matrix semantics. *Review of Symbolic Logic*, 11(4):780–804, 2018.
- [27] T. Moraschini. On equational completeness theorems. Submitted, available on the ArXiv, 2021.
- [28] T. Moraschini and J. J. Wannenburg. Epimorphisms in varieties of Heyting algebras. *Annals of Pure and Applied Logic*, 171(9), 2020.
- [29] W. Neumann. On Mal’cev conditions. *Journal of the Australian Mathematical Society*, 17:376–384, 1974.
- [30] J. Opršal. Taylor’s Modularity Conjecture and Related Problems for Idempotent Varieties. *Order*, 35(3):433–460, 2018.
- [31] A. F. Pixley. Local mal’cev conditions. *Canadian Mathematical Bulletin*, 15:559–568, 1972.
- [32] J. G. Raftery. The equational definability of truth predicates. *Reports on Mathematical Logic*, (41):95–149, 2006.
- [33] J. G. Raftery. Order algebraizable logics. *Annals of Pure and Applied Logic*, 164(3):251–283, 2013.
- [34] W. Rautenberg. On reduced matrices. *Studia Logica*, 52:63–72, 1993.

- [35] V. V. Rybakov. *Admissibility of logical inference rules*, volume 136 of *Studies in Logic*. Elsevier, Amsterdam, etc., 1997.
- [36] W. Taylor. Characterizing Mal'cev conditions. *Algebra Universalis*, 3:351–397, 1973.
- [37] W. Taylor. Varieties obeying homotopy laws. *Canadian Journal of Mathematics*, 29:498–527, 1977.
- [38] S. Tschantz. Congruence permutability is join prime. Unpublished manuscript, 1996.
- [39] R. Wille. *Kongruenzklassengeometrien*. Number 113 in Springer Lecture Notes. 1970.

DEPARTMENT OF PHILOSOPHY, UNIVERSITY OF BARCELONA, CARRER DE MONTALEGRE 6, 08001,
BARCELONA, SPAIN

Email address: jansana@ub.edu

Email address: tommaso.moraschini@ub.edu