

On interpretations between propositional logics

Tommaso Moraschini
joint with Ramon Jansana

Institute of Computer Science
Czech Academy of Sciences

BLAST 2019
University of Colorado, Boulder

Aim of the talk

To introduce a notion of **interpretability** between propositional logics and investigate the resulting “poset of **all** logics”.

Aim of the talk

To introduce a notion of **interpretability** between propositional logics and investigate the resulting “poset of **all** logics”.

Remark. In most results, propositional logics can be replaced by infinitary universal Horn theories without equality.

Aim of the talk

To introduce a notion of **interpretability** between propositional logics and investigate the resulting “poset of **all** logics”.

Remark. In most results, propositional logics can be replaced by infinitary universal Horn theories without equality.

Some sources of inspiration:

- ▶ **Matrix semantics** for logics (Łukasiewicz, Tarski, Łos, Suszko, Wójcicki . . .)
- ▶ Blok and Pigozzi’s seminal work on **algebraizable logics**
- ▶ **Leibniz hierarchy** of propositional logics (Czelakowski, Font, Herrmann, Jansana, Raftery . . .)
- ▶ **Maltsev conditions** (Day, Maltsev, Jónsson, Pixley, Kiss, Kearnes, McKenzie, Szendrei . . .)
- ▶ **Interpretations** between varieties (Taylor, Neumann, Garcia, Opršal, Tschantz . . .)

Interpretations between logics

Interpretations between logics

Basic questions:

Interpretations between logics

Basic questions:

- ▶ What do we mean by an **interpretation** between logics?

Interpretations between logics

Basic questions:

- ▶ What do we mean by an **interpretation** between logics?
- ▶ And what do we mean by **logic**?

Equality-free model theory

Let T be a first-order theory without equality.

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

$a \equiv c \iff a$ and c satisfies the same

equality-free types with constants

\iff for every non-equality atomic formula $\phi(x, y_1, \dots, y_n)$

and for every $b_1, \dots, b_n \in M$,

$\mathbf{M} \models \phi(a, b_1, \dots, b_n)$ iff $\mathbf{M} \models \phi(c, b_1, \dots, b_n)$.

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

$a \equiv c \iff a$ and c satisfies the same

equality-free types with constants

\iff for every non-equality atomic formula $\phi(x, y_1, \dots, y_n)$

and for every $b_1, \dots, b_n \in M$,

$\mathbf{M} \models \phi(a, b_1, \dots, b_n)$ iff $\mathbf{M} \models \phi(c, b_1, \dots, b_n)$.

- ▶ The indiscernibility relation is a congruence on \mathbf{M} , and the indiscernibility relation of the quotient \mathbf{M}/\equiv is the **identity**.

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

$a \equiv c \iff a$ and c satisfies the same

equality-free types with constants

\iff for every non-equality atomic formula $\phi(x, y_1, \dots, y_n)$

and for every $b_1, \dots, b_n \in M$,

$\mathbf{M} \models \phi(a, b_1, \dots, b_n)$ iff $\mathbf{M} \models \phi(c, b_1, \dots, b_n)$.

- ▶ The indiscernibility relation is a congruence on \mathbf{M} , and the indiscernibility relation of the quotient \mathbf{M}/\equiv is the **identity**.
- ▶ \mathbf{M}/\equiv satisfies the **same** sentences without equality than \mathbf{M} .

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

$a \equiv c \iff a$ and c satisfies the same

equality-free types with constants

\iff for every non-equality atomic formula $\phi(x, y_1, \dots, y_n)$

and for every $b_1, \dots, b_n \in M$,

$\mathbf{M} \models \phi(a, b_1, \dots, b_n)$ iff $\mathbf{M} \models \phi(c, b_1, \dots, b_n)$.

- ▶ The indiscernibility relation is a congruence on \mathbf{M} , and the indiscernibility relation of the quotient \mathbf{M}/\equiv is the **identity**.
- ▶ \mathbf{M}/\equiv satisfies the **same** sentences without equality than \mathbf{M} .
- ▶ Thus, the natural models of T are the ones whose indiscernibility relation is the identity relation.

Equality-free model theory

Let T be a first-order theory without equality.

- ▶ Every model \mathbf{M} of T is associated with an **indiscernibility** relation \equiv that mimics equality: for every $a, c \in M$,

$a \equiv c \iff a$ and c satisfies the same

equality-free types with constants

\iff for every non-equality atomic formula $\phi(x, y_1, \dots, y_n)$

and for every $b_1, \dots, b_n \in M$,

$\mathbf{M} \models \phi(a, b_1, \dots, b_n)$ iff $\mathbf{M} \models \phi(c, b_1, \dots, b_n)$.

- ▶ The indiscernibility relation is a congruence on \mathbf{M} , and the indiscernibility relation of the quotient \mathbf{M}/\equiv is the **identity**.
- ▶ \mathbf{M}/\equiv satisfies the **same** sentences without equality than \mathbf{M} .
- ▶ Thus, the natural models of T are the ones whose indiscernibility relation is the identity relation.
- ▶ This setting subsumes model theory **with** equality.

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics are infinitary Horn theories without equality.

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics are infinitary Horn theories without equality.

- ▶ Let \vdash be a logic and let $P(x)$ be a unary predicate symbol.

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics are infinitary Horn theories without equality.

- ▶ Let \vdash be a logic and let $P(x)$ be a unary predicate symbol.
- ▶ Let T_{\vdash} be the theory in the equality-free language obtained extending the algebraic language of \vdash with $P(x)$,

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics are infinitary Horn theories without equality.

- ▶ Let \vdash be a logic and let $P(x)$ be a unary predicate symbol.
- ▶ Let T_{\vdash} be the theory in the equality-free language obtained extending the algebraic language of \vdash with $P(x)$, axiomatized by the infinitary universal Horn sentences

$$\forall \vec{x} \bigwedge_{\gamma \in \Gamma} P(\gamma(\vec{x})) \rightarrow P(\varphi(\vec{x}))$$

for all valid inferences $\Gamma \vdash \varphi$ of \vdash .

- ▶ A **logic** is a consequence relation \vdash on the set \mathbf{Fm} of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$.

- ▶ A **logic** is a consequence relation \vdash on the set \mathbf{Fm} of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$.
A matrix $\langle \mathbf{A}, F \rangle$ is a

- ▶ A **logic** is a consequence relation \vdash on the set \mathbf{Fm} of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$. A matrix $\langle \mathbf{A}, F \rangle$ is a **model** of a logic \vdash (in the same language) if for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$,

- ▶ A **logic** is a consequence relation \vdash on the set \mathbf{Fm} of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$. A matrix $\langle \mathbf{A}, F \rangle$ is a **model** of a logic \vdash (in the same language) if for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$,

if $\Gamma \vdash \varphi$, then for every hom $v: \mathbf{Fm} \rightarrow \mathbf{A}$,

if $v[\Gamma] \subseteq F$, then $v(\varphi) \in F$.

- ▶ A **logic** is a consequence relation \vdash on the set **Fm** of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$. A matrix $\langle \mathbf{A}, F \rangle$ is a **model** of a logic \vdash (in the same language) if for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$,

$$\begin{aligned} &\text{if } \Gamma \vdash \varphi, \text{ then for every hom } v: \mathbf{Fm} \rightarrow \mathbf{A}, \\ &\quad \text{if } v[\Gamma] \subseteq F, \text{ then } v(\varphi) \in F. \end{aligned}$$

Intuitively, \mathbf{A} is an algebra of **truth-values** and F are the values representing **truth**.

- ▶ A **logic** is a consequence relation \vdash on the set \mathbf{Fm} of formulas of some algebraic language with **infinitely many** variables that, moreover, is **substitution invariant** in the sense that

$$\text{if } \Gamma \vdash \varphi, \text{ then } \sigma[\Gamma] \vdash \sigma(\varphi)$$

for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$ and every substitution σ .

Remark. Logics \vdash are infinitary Horn theories without equality T_{\vdash} .

- ▶ A **matrix** is a pair $\langle \mathbf{A}, F \rangle$ where \mathbf{A} is an algebra and $F \subseteq A$. A matrix $\langle \mathbf{A}, F \rangle$ is a **model** of a logic \vdash (in the same language) if for every $\Gamma \cup \{\varphi\} \subseteq \mathbf{Fm}$,

$$\begin{aligned} \text{if } \Gamma \vdash \varphi, \text{ then for every hom } v: \mathbf{Fm} \rightarrow \mathbf{A}, \\ \text{if } v[\Gamma] \subseteq F, \text{ then } v(\varphi) \in F. \end{aligned}$$

Intuitively, \mathbf{A} is an algebra of **truth-values** and F are the values representing **truth**.

- ▶ Observe that $\langle \mathbf{A}, F \rangle$ is a model of \vdash iff it is a model of T_{\vdash} in the standard sense.

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

Examples.

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

Examples.

- ▶ If \mathbf{A} is a Heyting algebra and F a lattice filter, then

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

Examples.

- ▶ If \mathbf{A} is a Heyting algebra and F a lattice filter, then

$$a \equiv c \iff \{a \rightarrow c, c \rightarrow a\} \subseteq F.$$

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

Examples.

- ▶ If \mathbf{A} is a Heyting algebra and F a lattice filter, then

$$a \equiv c \iff \{a \rightarrow c, c \rightarrow a\} \subseteq F.$$

- ▶ If \mathbf{A} is a modal algebra and F a lattice filter, then

- ▶ Every matrix $\langle \mathbf{A}, F \rangle$ is associated the an **indiscernibility** relation \equiv defined for every $a, c \in A$ as

$$a \equiv c \iff p(a) \in F \text{ iff } p(c) \in F,$$

for all unary polynomial functions p of \mathbf{A} .

Examples.

- ▶ If \mathbf{A} is a Heyting algebra and F a lattice filter, then

$$a \equiv c \iff \{a \rightarrow c, c \rightarrow a\} \subseteq F.$$

- ▶ If \mathbf{A} is a modal algebra and F a lattice filter, then

$$a \equiv c \iff \{\Box^n(a \rightarrow c), \Box^n(c \rightarrow a) : n \in \omega\} \subseteq F.$$

- ▶ Logics \vdash are associated with models **without** indiscernibles

- ▶ Logics \vdash are associated with models **without** indiscernibles

$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$

- ▶ Logics \vdash are associated with models **without** indiscernibles

$$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$$

Completeness.

- ▶ Logics \vdash are associated with models **without** indiscernibles

$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$

Completeness. \vdash is the logic induced by the class $\text{Mod}^{\equiv}(\vdash)$, i.e.

- ▶ Logics \vdash are associated with models **without** indiscernibles

$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$

Completeness. \vdash is the logic induced by the class $\text{Mod}^{\equiv}(\vdash)$, i.e.

$\Gamma \vdash \varphi \iff$ for every $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ and hom $v: \mathbf{Fm} \rightarrow \mathbf{A}$,
if $v[\Gamma] \subseteq F$, then $v(\varphi) \in F$.

- ▶ Logics \vdash are associated with models **without** indiscernibles

$$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$$

Completeness. \vdash is the logic induced by the class $\text{Mod}^{\equiv}(\vdash)$, i.e.

$$\Gamma \vdash \varphi \iff \text{for every } \langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash) \text{ and hom } v : \mathbf{Fm} \rightarrow \mathbf{A}, \\ \text{if } v[\Gamma] \subseteq F, \text{ then } v(\varphi) \in F.$$

Examples.

- ▶ Logics \vdash are associated with models **without** indiscernibles

$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and } \equiv \text{ is the identity relation}\}.$

Completeness. \vdash is the logic induced by the class $\text{Mod}^{\equiv}(\vdash)$, i.e.

$\Gamma \vdash \varphi \iff$ for every $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$ and hom $v: \mathbf{Fm} \rightarrow \mathbf{A}$,
if $v[\Gamma] \subseteq F$, then $v(\varphi) \in F$.

Examples.

$\text{Mod}^{\equiv}(\mathbf{CPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\}\}$

- ▶ Logics \vdash are associated with models **without** indiscernibles

$$\text{Mod}^{\equiv}(\vdash) := \mathbb{P}_{\text{sd}}\{\langle \mathbf{A}, F \rangle : \langle \mathbf{A}, F \rangle \text{ is a model of } \vdash \text{ and} \\ \equiv \text{ is the identity relation}\}.$$

Completeness. \vdash is the logic induced by the class $\text{Mod}^{\equiv}(\vdash)$, i.e.

$$\Gamma \vdash \varphi \iff \text{for every } \langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash) \text{ and hom } v : \mathbf{Fm} \rightarrow \mathbf{A}, \\ \text{if } v[\Gamma] \subseteq F, \text{ then } v(\varphi) \in F.$$

Examples.

$$\text{Mod}^{\equiv}(\text{CPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\}\}$$

$$\text{Mod}^{\equiv}(\text{IPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Heyting algebra and } F = \{1\}\}.$$

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

Example.

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{ \tau(f)^\mathbf{A} : f \in \mathcal{L} \} \rangle.$$

Example. Let $\mathcal{L}_{\wedge \vee}$ be the language of **lattices**, and \mathcal{L}_{BA} that of **Boolean algebras**.

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

Example. Let $\mathcal{L}_{\wedge\vee}$ be the language of **lattices**, and \mathcal{L}_{BA} that of **Boolean algebras**. If τ is the inclusion map from $\mathcal{L}_{\wedge\vee}$ to \mathcal{L}_{BA} , and \mathbf{A} a Boolean algebra, then \mathbf{A}^τ is its lattice reduct of \mathbf{A} .

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{ \tau(f)^\mathbf{A} : f \in \mathcal{L} \} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^\equiv(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^\equiv(\vdash).$$

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^\equiv(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^\equiv(\vdash).$$

Interpretations split in two halves.

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^{\equiv}(\vdash).$$

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^\equiv(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^\equiv(\vdash).$$

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^\equiv(\vdash)$, $\text{Mod}^\equiv(\vdash')$.

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^\equiv(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^\equiv(\vdash).$$

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^\equiv(\vdash)$, $\text{Mod}^\equiv(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_\vdash \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_\vdash -reducts of the matrices in $\text{Mod}^\equiv(\vdash')$ belong to $\text{Mod}^\equiv(\vdash)$.

- ▶ A **translation** of an algebraic language \mathcal{L} into another \mathcal{L}' is a map τ that assigns an n -ary term $\tau(f)(x_1, \dots, x_n)$ of \mathcal{L}' to every n -ary symbol $f(x_1, \dots, x_n)$ of \mathcal{L} .
- ▶ Given an \mathcal{L}' -algebra \mathbf{A} , we define an \mathcal{L} -algebra

$$\mathbf{A}^\tau := \langle \mathbf{A}; \{\tau(f)^\mathbf{A} : f \in \mathcal{L}\} \rangle.$$

- ▶ An **interpretation** of a logic \vdash into another \vdash' is a translation τ between their languages such that

$$\text{if } \langle \mathbf{A}, F \rangle \in \text{Mod}^\equiv(\vdash'), \text{ then } \langle \mathbf{A}^\tau, F \rangle \in \text{Mod}^\equiv(\vdash).$$

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^\equiv(\vdash)$, $\text{Mod}^\equiv(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_\vdash \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_\vdash -reducts of the matrices in $\text{Mod}^\equiv(\vdash')$ belong to $\text{Mod}^\equiv(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Example. Recall that

$$\text{Mod}^{\equiv}(\mathbf{CPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\}\}$$

$$\text{Mod}^{\equiv}(\mathbf{IPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Heyting algebra and } F = \{1\}\}.$$

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Example. Recall that

$$\text{Mod}^{\equiv}(\mathbf{CPC}) = \{ \langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\} \}$$

$$\text{Mod}^{\equiv}(\mathbf{IPC}) = \{ \langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Heyting algebra and } F = \{1\} \}.$$

- ▶ The **identity** map is an interpretation of **IPC** into **CPC**.

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Example. Recall that

$$\text{Mod}^{\equiv}(\mathbf{CPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\}\}$$

$$\text{Mod}^{\equiv}(\mathbf{IPC}) = \{\langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Heyting algebra and } F = \{1\}\}.$$

- ▶ The **identity** map is an interpretation of **IPC** into **CPC**.
- ▶ Is **CPC** interpretable into **IPC**?

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Example. Recall that

$$\text{Mod}^{\equiv}(\mathbf{CPC}) = \{ \langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Boolean algebra and } F = \{1\} \}$$

$$\text{Mod}^{\equiv}(\mathbf{IPC}) = \{ \langle \mathbf{A}, F \rangle : \mathbf{A} \text{ is a Heyting algebra and } F = \{1\} \}.$$

- ▶ The **identity** map is an interpretation of **IPC** into **CPC**.
- ▶ Is **CPC** interpretable into **IPC**? No, on **cardinality** grounds!

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Definition

Intepretability is a preorder on the proper class of all logics.

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Definition

Intepretability is a preorder on the proper class of all logics.
The associated partial order **Log** is the “**poset of all logics**”.

Interpretations split in two halves. Consider logics \vdash and \vdash' .

- ▶ \vdash and \vdash' are **term-equivalent** if so are $\text{Mod}^{\equiv}(\vdash)$, $\text{Mod}^{\equiv}(\vdash')$.
- ▶ \vdash' is a **compatible expansion** of \vdash if $\mathcal{L}_{\vdash} \subseteq \mathcal{L}_{\vdash'}$ and the \mathcal{L}_{\vdash} -reducts of the matrices in $\text{Mod}^{\equiv}(\vdash')$ belong to $\text{Mod}^{\equiv}(\vdash)$.

Lemma

\vdash is interpretable into \vdash' iff \vdash' is term-equivalent to a compatible expansion of \vdash .

Definition

Intepretability is a preorder on the proper class of all logics. The associated partial order **Log** is the “**poset of all logics**”.

- ▶ Elements of **Log** are classes $[[\vdash]]$ of **equi-interpretable** logics.

The structure of the poset of all logics

The structure of the poset of all logics

Basic question:

The structure of the poset of all logics

Basic question:

- ▶ Do **infima** and **suprema** exist?

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$,

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i: i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i: i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra,

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i: i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n): i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i: i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)): i \in I \rangle.$$

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i: i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n): i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i: i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)): i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i: i \in I\}$,

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i : i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i : i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\left\{ \left\langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \right\rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \right\}$$

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i : i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i : i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\left\{ \left\langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \right\rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \right\}$$

formulated with $\prod_{i \in I} |\mathbf{Fm}(\vdash_i)|$ variables.

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i : i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i : i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\left\{ \left\langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \right\rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \right\}.$$

$\bigotimes_{i \in I} \vdash_i$ is called the **non-indexed product** of the various \vdash_i .

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i : i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i : i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\{ \langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \}.$$

Theorem

- ▶ $\text{Mod}^{\equiv}(\bigotimes_{i \in I} \vdash_i)$ is the closure under \mathbb{P}_{sd} of the above display.

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i : i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i : i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i : i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\{ \langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \}.$$

Theorem

- ▶ $\text{Mod}^{\equiv}(\bigotimes_{i \in I} \vdash_i)$ is the closure under \mathbb{P}_{sd} of the above display.
- ▶ $\llbracket \bigotimes_{i \in I} \vdash_i \rrbracket$ is the infimum of $\{\llbracket \vdash_i \rrbracket : i \in I\}$ in Log.

Do infima exist?

- ▶ Given a set of algebraic languages $\{\mathcal{L}_i: i \in I\}$, let $\bigotimes_{i \in I} \mathcal{L}_i$ be the language whose n -ary operations f are the sequences

$$f = \langle \varphi_i(x_1, \dots, x_n) : i \in I \rangle$$

where φ_i is an n -ary term of \mathcal{L}_i .

- ▶ Given a set of algebras $\{\mathbf{A}_i: i \in I\}$ s.t. \mathbf{A}_i is an \mathcal{L}_i -algebra, let $\bigotimes_{i \in I} \mathbf{A}_i$ be the $\bigotimes_{i \in I} \mathcal{L}_i$ -algebra with universe $\prod_{i \in I} A_i$ s.t.

$$f \bigotimes_{i \in I} \mathbf{A}_i(\vec{a}_1, \dots, \vec{a}_n) := \langle \varphi_i^{\mathbf{A}_i}(\vec{a}_1(i), \dots, \vec{a}_n(i)) : i \in I \rangle.$$

- ▶ Given a set of logics $\{\vdash_i: i \in I\}$, let $\bigotimes_{i \in I} \vdash_i$ be the logic induced by the class of matrices

$$\{ \langle \bigotimes_{i \in I} \mathbf{A}_i, \prod_{i \in I} F_i \rangle : \langle \mathbf{A}_i, F_i \rangle \in \text{Mod}^{\equiv}(\vdash_i) \}.$$

Theorem

Log is a **set-complete** meet-semilattice.

Do suprema exist?

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let CPC_{\neg} be the **negation** fragment of classical logic,

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let CPC_{\neg} be the **negation** fragment of classical logic,

$$x \triangleright \neg\neg x \quad \neg\neg x \triangleright x \quad x, \neg x \triangleright y.$$

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let \mathbf{CPC}_{\neg} be the **negation** fragment of classical logic,

$$x \triangleright \neg\neg x \quad \neg\neg x \triangleright x \quad x, \neg x \triangleright y.$$


- ▶ Consider the algebra $\mathbf{A} = \langle A; \vee, \mathbf{a}, \mathbf{b}, \mathbf{0} \rangle$ depicted below.

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let \mathbf{CPC}_{\neg} be the **negation** fragment of classical logic,

$$x \triangleright \neg\neg x \quad \neg\neg x \triangleright x \quad x, \neg x \triangleright y.$$

- ▶ Consider the algebra $\mathbf{A} = \langle A; \vee, \mathbf{a}, \mathbf{b}, \mathbf{0} \rangle$ depicted below.




Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let \mathbf{CPC}_{\neg} be the **negation** fragment of classical logic,

$$x \triangleright \neg\neg x \quad \neg\neg x \triangleright x \quad x, \neg x \triangleright y.$$

- ▶ Consider the algebra $\mathbf{A} = \langle A; \vee, \mathbf{a}, \mathbf{b}, \mathbf{0} \rangle$ depicted below.



Then let \mathbf{L} be the logic induced by the pair of matrices


$$\{\langle \mathbf{A}, \{1\} \rangle, \langle \mathbf{A}, \{1, c\} \rangle\}.$$

Do suprema exist?

- ▶ **No**: even **binary** suprema may fail to exist.
- ▶ Let \mathbf{CPC}_{\neg} be the **negation** fragment of classical logic,

$$x \triangleright \neg\neg x \quad \neg\neg x \triangleright x \quad x, \neg x \triangleright y.$$

- ▶ Consider the algebra $\mathbf{A} = \langle A; \vee, \mathbf{a}, \mathbf{b}, \mathbf{0} \rangle$ depicted below.



Then let \mathbf{L} be the logic induced by the pair of matrices

$$\{\langle \mathbf{A}, \{1\} \rangle, \langle \mathbf{A}, \{1, c\} \rangle\}.$$

- ▶ The **supremum** of $\llbracket \mathbf{CPC}_{\neg} \rrbracket$ and $\llbracket \mathbf{L} \rrbracket$ does **not** exist in Log.

Leibniz classes and hierarchy

Leibniz classes and hierarchy

Basic question:

Leibniz classes and hierarchy

Basic question:

- ▶ What are **Leibniz classes** of logics?

What is the Leibniz hierarchy?

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ that **defines indiscernibility**

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Syntactic characterization.** A logic \vdash is **equivalential** iff there is a non-empty set $\Delta(x, y)$ of formulas s.t.

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Syntactic characterization.** A logic \vdash is **equivalential** iff there is a non-empty set $\Delta(x, y)$ of formulas s.t.

$$\begin{aligned} \emptyset \vdash \Delta(x, x) \quad & x, \Delta(x, y) \vdash y \\ \bigcup_{1 \leq i \leq n} \Delta(x_i, y_i) \vdash & \Delta(f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \end{aligned}$$

for every n -ary connective f .

What is the Leibniz hierarchy?

- ▶ A classification of logics in terms **syntactic principles** that govern the behaviour of the **indiscernibility** relation.

Example.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Syntactic characterization.** A logic \vdash is **equivalential** iff there is a non-empty set $\Delta(x, y)$ of formulas s.t.

$$\begin{aligned} \emptyset \vdash \Delta(x, x) \quad & x, \Delta(x, y) \vdash y \\ \bigcup_{1 \leq i \leq n} \Delta(x_i, y_i) \vdash & \Delta(f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \end{aligned}$$

for every n -ary connective f .

- ▶ Equivalential logics form a **Leibniz class**.

► A **Leibniz condition** is a sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$ of logics, indexed by all ordinals, s.t.

if $\alpha \leq \beta$, then \vdash_β is **interpretable** into \vdash_α .

- ▶ A **Leibniz condition** is a sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$ of logics, indexed by all ordinals, s.t.

if $\alpha \leq \beta$, then \vdash_β is **interpretable** into \vdash_α .

- ▶ A logic \vdash **satisfies** Φ if some \vdash_α is **intepretable** in \vdash .

- ▶ A **Leibniz condition** is a sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$ of logics, indexed by all ordinals, s.t.

if $\alpha \leq \beta$, then \vdash_β is **interpretable** into \vdash_α .

- ▶ A logic \vdash **satisfies** Φ if some \vdash_α is **interpretable** in \vdash . Let

$\text{Log}(\Phi) := \{\vdash : \vdash \text{ is a logic and } \text{satisfies } \Phi\}$.

- ▶ A **Leibniz condition** is a sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$ of logics, indexed by all ordinals, s.t.

if $\alpha \leq \beta$, then \vdash_β is **interpretable** into \vdash_α .

- ▶ A logic \vdash **satisfies** Φ if some \vdash_α is **interpretable** in \vdash . Let

$\text{Log}(\Phi) := \{\vdash : \vdash \text{ is a logic and } \text{satisfies } \Phi\}$.

- ▶ A **Leibniz class** is a class of logics of the form $\text{Log}(\Phi)$, for some Leibniz condition Φ .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of $3 \Rightarrow 1$.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of $3 \Rightarrow 1$.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ ,

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then some \vdash_α is interpretable into \vdash .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then \vdash is a term-equivalent to a compatible expansion of some \vdash_α .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then \vdash is a term-equivalent to a compatible expansion of a non-indexed product of a set of logics in K .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log.
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.
- ▶ If $\vdash \in K$,

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.
- ▶ If $\vdash \in K$, then $\vdash \in V_\alpha \cap K = K_\alpha$ for some α .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.
- ▶ If $\vdash \in K$, then \vdash_α is interpretable in \vdash for some α .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\otimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.
- ▶ If $\vdash \in K$, then \vdash satisfies Φ .

Theorem

Let K be a class of logics. TFAE:

1. K is a Leibniz class.
2. K is “essentially” a set-complete filter of Log .
3. K is closed under the formation of **term-equivalent** logics, **compatible expansions**, and **non-indexed products** indexed by arbitrarily large sets.

Proof sketch of 3 \Rightarrow 1.

- ▶ Consider the **cumulative hierarchy** of sets $\{V_\alpha : \alpha \in \text{OR}\}$.
- ▶ For every ordinal α , define the **set** $K_\alpha := K \cap V_\alpha$.
- ▶ Let \vdash_α be the non-indexed product $\bigotimes K_\alpha$.
- ▶ Then consider the sequence $\Phi = \{\vdash_\alpha : \alpha \in \text{OR}\}$.
- ▶ Φ is a Leibniz condition, as if $\alpha \leq \beta$, then $K_\alpha \subseteq K_\beta$.
- ▶ If \vdash satisfies Φ , then $\vdash \in K$.
- ▶ If $\vdash \in K$, then \vdash satisfies Φ .
- ▶ K is the class of logics satisfying Φ .

Indecomposable Leibniz classes

Indecomposable Leibniz classes

Basic question:

Indecomposable Leibniz classes

Basic question:

- ▶ Which of Leibniz classes are **primitive** or fundamental?

- ▶ When ordered under inclusion, Leibniz classes form a “lattice”.

Definition

A Leibniz class K is said to be

- ▶ When ordered under inclusion, Leibniz classes form a “lattice”.

Definition

A Leibniz class K is said to be

- ▶ **meet-irreducible** if for every pair K_1 and K_2 of Leibniz classes (of logics with some tautology),

if $K = K_1 \cap K_2$, then either $K = K_1$ or $K = K_2$.

- ▶ When ordered under inclusion, Leibniz classes form a “lattice”.

Definition

A Leibniz class K is said to be

- ▶ **meet-irreducible** if for every pair K_1 and K_2 of Leibniz classes (of logics with some tautology),

if $K = K_1 \cap K_2$, then either $K = K_1$ or $K = K_2$.

- ▶ **meet-prime** if for every pair of Leibniz classes K_1 and K_2 (of logics with some tautology),

if $K \subseteq K_1 \cap K_2$, then either $K \subseteq K_1$ or $K \subseteq K_2$.

- ▶ When ordered under inclusion, Leibniz classes form a “lattice”.

Definition

A Leibniz class K is said to be

- ▶ **meet-irreducible** if for every pair K_1 and K_2 of Leibniz classes (of logics with some tautology),

if $K = K_1 \cap K_2$, then either $K = K_1$ or $K = K_2$.

- ▶ **meet-prime** if for every pair of Leibniz classes K_1 and K_2 (of logics with some tautology),

if $K \subseteq K_1 \cap K_2$, then either $K \subseteq K_1$ or $K \subseteq K_2$.

- ▶ Intuitively, a Leibniz class is meet-prime (resp. irreducible) when it captures a **fundamental concept**.

- ▶ When ordered under inclusion, Leibniz classes form a “lattice”.

Definition

A Leibniz class K is said to be

- ▶ **meet-irreducible** if for every pair K_1 and K_2 of Leibniz classes (of logics with some tautology),

if $K = K_1 \cap K_2$, then either $K = K_1$ or $K = K_2$.

- ▶ **meet-prime** if for every pair of Leibniz classes K_1 and K_2 (of logics with some tautology),

if $K \subseteq K_1 \cap K_2$, then either $K \subseteq K_1$ or $K \subseteq K_2$.

- ▶ Intuitively, a Leibniz class is meet-prime (resp. irreducible) when it captures a **fundamental concept**.
- ▶ We shall apply this test to two conditions, i.e. the definability of truth-sets and of indiscernibility.

Definability of truth-sets.

- ▶ A logic \vdash is **truth-equational** if there is a set of equations $E(x)$ s.t. for every $\langle \mathbf{A}, F \rangle \in \text{Mod}^{\equiv}(\vdash)$

$$a \in F \iff \mathbf{A} \vDash E(a), \text{ for all } a \in A.$$

Definability of truth-sets.

- ▶ A logic \vdash with tautologies is **truth-equational** if there are no $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \subsetneq F \subsetneq G$.

Definability of truth-sets.

- ▶ A logic \vdash with tautologies is **truth-equational** if there are no $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \subsetneq F \subsetneq G$.

Theorem

Truth-equational logics form a **meet-prime** Leibniz class.

Proof sketch.

Definability of truth-sets.

- ▶ A logic \vdash with tautologies is **truth-equational** if there are no $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \subsetneq F \subsetneq G$.

Theorem

Truth-equational logics form a **meet-prime** Leibniz class.

Proof sketch.

- ▶ Let \vdash_1, \vdash_2 be non truth-equational logics (with tautologies).

Definability of truth-sets.

- ▶ A logic \vdash with tautologies is **truth-equational** if there are no $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \subsetneq F \subsetneq G$.

Theorem

Truth-equational logics form a **meet-prime** Leibniz class.

Proof sketch.

- ▶ Let \vdash_1, \vdash_2 be non truth-equational logics (with tautologies).
- ▶ **Goal:** find a **non truth-equational** logics in which \vdash_1 and \vdash_2 are **interpretable**.

Definability of truth-sets.

- ▶ A logic \vdash with tautologies is **truth-equational** if there are no $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ such that $\emptyset \subsetneq F \subsetneq G$.

Theorem


Truth-equational logics form a **meet-prime** Leibniz class.

Proof sketch.


- ▶ Let \vdash_1, \vdash_2 be non truth-equational logics (with tautologies).
- ▶ **Goal:** find a **non truth-equational** logics in which \vdash_1 and \vdash_2 are **interpretable**.
- ▶ As \vdash_1 and \vdash_2 are not truth-equational, there are matrices


$$\begin{aligned} \langle \mathbf{A}_1, F_1 \rangle, \langle \mathbf{A}_1, G_1 \rangle &\in \text{Mod}^{\equiv}(\vdash_1) \text{ s.t. } \emptyset \subsetneq F_1 \subsetneq G_1 \\ \langle \mathbf{A}_2, F_2 \rangle, \langle \mathbf{A}_2, G_2 \rangle &\in \text{Mod}^{\equiv}(\vdash_2) \text{ s.t. } \emptyset \subsetneq F_2 \subsetneq G_2. \end{aligned}$$

Algebra A_1




Algebra A_2






- ▶ We want to **merge** the two algebras into a single one.




- ▶ We want to **merge** the two algebras into a single one.
- ▶ The problem is that A_1 and A_2 have not the same universe.




- ▶ We want to **merge** the two algebras into a single one.
- ▶ The problem is that A_1 and A_2 have not the same universe.
- ▶ This is solved by “**adding points**” to A_1 and A_2 , taking sufficiently large direct powers.


Algebra A_1^k




Algebra A_2^k




- ▶ We want to **merge** the two algebras into a single one.
- ▶ The problem is that A_1 and A_2 have not the same universe.
- ▶ This is solved by “**adding points**” to A_1 and A_2 , taking sufficiently large direct powers.




- ▶ We want to **merge** the two algebras into a single one.
- ▶ The problem is that A_1 and A_2 have not the same universe.
- ▶ This is solved by “**adding points**” to A_1 and A_2 , taking sufficiently large direct powers.
- ▶ We assume w.l.o.g. that A_1 is A_1^K and A_2 is A_2^K .




- ▶ We **merge** A_1 and A_2 into an algebra A with universe $A = A_1 = A_2$ endowed with **all finitary operations**.




- ▶ We **merge** \mathbf{A}_1 and \mathbf{A}_2 into an algebra \mathbf{A} with universe $A = A_1 = A_2$ endowed with **all finitary operations**.
- ▶ Let \vdash be the logic induced by the matrices $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{A}, G \rangle$.




- ▶ We **merge** \mathbf{A}_1 and \mathbf{A}_2 into an algebra \mathbf{A} with universe $A = A_1 = A_2$ endowed with **all finitary operations**.
- ▶ Let \vdash be the logic induced by the matrices $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{A}, G \rangle$.
- ▶ **Goal:** to show that \vdash is not truth-equational and that \vdash_1 and \vdash_2 are interpretable in \vdash .



- ▶ We **merge** \mathbf{A}_1 and \mathbf{A}_2 into an algebra \mathbf{A} with universe $A = A_1 = A_2$ endowed with **all finitary operations**.
- ▶ Let \vdash be the logic induced by the matrices $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{A}, G \rangle$.
- ▶ \vdash is **not truth-equational**, since $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle \in \text{Mod}^{\equiv}(\vdash)$ and $\emptyset \subsetneq F \subsetneq G$.



- ▶ We **merge** \mathbf{A}_1 and \mathbf{A}_2 into an algebra \mathbf{A} with universe $A = A_1 = A_2$ endowed with **all finitary operations**.
- ▶ Let \vdash be the logic induced by the matrices $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{A}, G \rangle$.
- ▶ \vdash_i is **interpretable** into \vdash , since \vdash is induced by matrices $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle$ with a reduct in $\text{Mod}^{\equiv}(\vdash_i)$.



- ▶ We **merge** \mathbf{A}_1 and \mathbf{A}_2 into an algebra \mathbf{A} with universe $A = A_1 = A_2$ endowed with **all finitary operations**.
- ▶ Let \vdash be the logic induced by the matrices $\langle \mathbf{A}, F \rangle$ and $\langle \mathbf{A}, G \rangle$.
- ▶ \vdash_i is **interpretable** into \vdash , since \vdash is induced by matrices $\langle \mathbf{A}, F \rangle, \langle \mathbf{A}, G \rangle$ with a reduct in $\text{Mod}^{\equiv}(\vdash_i)$.
- ▶ The Leibniz class of truth-equational logics is a prime.

Definability of the indiscernibility relation.

Definability of the indiscernibility relation.

- ▶ A logic \vdash is **equivential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

Definability of the indiscernibility relation.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Problem.**

The class of equivalential logics is **not** meet-irreducible.

Definability of the indiscernibility relation.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Problem.**

The class of equivalential logics is **not** meet-irreducible.

- ▶ The class of equivalential logics is given by the Leibniz condition

$$\Phi = \{\vdash_{\alpha}^{\text{eq}} : \alpha \in \text{OR}\}$$

where $\vdash_{\alpha}^{\text{eq}}$ is the logic in the language with binary symbols $\{\neg_{\epsilon} : \epsilon < \max\{\omega, |\alpha|\}\}$ axiomatized by the rules

$$\begin{aligned} \emptyset \triangleright \Delta_{\alpha}(x, x) \quad x, \Delta_{\alpha}(x, y) \triangleright y \\ \Delta_{\alpha}(x_1, y_1) \cup \Delta_{\alpha}(x_2, y_2) \triangleright \Delta_{\alpha}(x_1 \neg_{\epsilon} x_2, y_1 \neg_{\epsilon} y_2) \end{aligned}$$

where $\Delta_{\alpha} := \{x \neg_{\epsilon} y : \epsilon < \max\{\omega, |\alpha|\}\}$.

Definability of the indiscernibility relation.

- ▶ A logic \vdash is **equivalential** if there is a non-empty set of formulas $\Delta(x, y)$ s.t. for all models $\langle \mathbf{A}, F \rangle$ of \vdash and $a, c \in A$,

$$a \equiv c \iff \Delta^{\mathbf{A}}(a, c) \subseteq F.$$

- ▶ **Problem.**

The class of equivalential logics is **not** meet-irreducible.

- ▶ The class of equivalential logics is given by the Leibniz condition

$$\Phi = \{\vdash_{\alpha}^{\text{eq}} : \alpha \in \text{OR}\}$$

where $\vdash_{\alpha}^{\text{eq}}$ is the logic axiomatized by the rules

$$\begin{aligned} \emptyset \triangleright \Delta_{\alpha}(x, x) \quad x, \Delta_{\alpha}(x, y) \triangleright y \\ \Delta_{\alpha}(x_1, y_1) \cup \Delta_{\alpha}(x_2, y_2) \triangleright \Delta_{\alpha}(x_1 \multimap_{\epsilon} x_2, y_1 \multimap_{\epsilon} y_2). \end{aligned}$$

Theorem

The logic $\vdash_{\alpha}^{\text{eq}}$ is **completely meet-prime** in Log. Thus equivalential logics are determined by a Leibniz condition consisting only of completely meet-prime logics.

Thank you for your attention!