

PROFINITENESS AND REPRESENTABILITY OF SPECTRA OF HEYTING ALGEBRAS

G. BEZHANISHVILI, N. BEZHANISHVILI, T. MORASCHINI AND M. STRONKOWSKI

ABSTRACT. We prove that there exist profinite Heyting algebras that are not isomorphic to the profinite completion of any Heyting algebra. This resolves an open problem from 2009. More generally, we characterize those varieties of Heyting algebras in which profinite algebras are isomorphic to profinite completions. It turns out that there exists largest such. We give different characterizations of this variety and show that it is finitely axiomatizable and locally finite. From this it follows that it is decidable whether in a finitely axiomatizable variety of Heyting algebras all profinite members are profinite completions. In addition, we introduce and characterize representable varieties of Heyting algebras, thus drawing connection to the classical problem of representing posets as prime spectra.

CONTENTS

1. Introduction	1
2. Heyting algebras and Esakia spaces	4
2.1. Esakia duality	4
2.2. Depth and width in Heyting algebras	6
2.3. Profinite algebras and completions	8
3. Cascade Heyting algebras	9
4. Diamond systems and algebras	11
5. Forbidden posets	15
6. Topologies for diamond systems	21
7. Consequences	28
References	30

1. INTRODUCTION

An algebra is *profinite* if it is isomorphic to the inverse limit of an inverse system of finite algebras. This concept has its origin in the study of profinite groups. One of the basic results states that a group G is profinite if and only if it is a topological group whose topology is compact, Hausdorff, and zero-dimensional (i.e., a Stone topology). This result generalizes to many settings, including semigroups, monoids, rings, modules, distributive lattices, etc. (see for instance [43, Sec. VI.2] or more recent [21, 61]).

The *profinite completion* \widehat{A} of an algebra A is the inverse limit of the inverse system of its finite homomorphic images. It follows from the definition that \widehat{A} is a profinite algebra. However, not every profinite algebra A is isomorphic to \widehat{B} for some algebra B . In lattice theory, the problem of determining whether a profinite distributive lattice A is isomorphic to \widehat{B} for some distributive lattice B is related to Grätzer's celebrated problem of representable posets. Grätzer himself gave two formulations of this problem, for distributive lattices as for bounded distributive lattices (see [35, Problems 33 and 34, p. 156]). In this paper we will mainly concentrate on the bounded case. A poset X is *representable* if it is isomorphic to the *prime spectrum* (the poset of prime filters) of a bounded distributive lattice. Grätzer's problem asks for an internal characterization of representable posets. The same problem for

2010 *Mathematics Subject Classification.* 06D20, 06F30, 06E15, 03B55, 54F05.

Key words and phrases. Heyting algebra, profinite algebra, profinite completion, prime spectrum, Representation Problem, Priestley space, Esakia space.

prime spectra of commutative rings was posed by Kaplansky [45, pp. 5–7]. In fact, bounded distributive lattices and commutative rings with unit give rise to the same prime spectra up to isomorphism (see, e.g., [58, Thm. 1.1]). Even more is true. It is a well-known result of Hochster [37] that the prime spectra of commutative rings with unit are precisely the spectral spaces. Cornish [22] showed that these spaces form a category isomorphic to that of Priestley spaces, i.e., prime spectra of bounded distributive lattices [56, 57].

It follows from Priestley duality that a poset X is representable if and only if there is a Stone topology on X such that the resulting ordered space is a Priestley space. Furthermore, in view of the work of Joyal [44] and Speed [64], a poset X is representable precisely when it is profinite. Thus, Grätzer’s problem asks for a description of profinite posets, or equivalently of the posets underlying Priestley spaces.

A solution of Grätzer’s problem for linearly ordered sets was given independently by Balbes [3, Thm. 9] (by lattice-theoretic means) and Lewis [48, Thm. 3.1] (by ring-theoretic means). It follows from their characterizations that representable chains are necessarily complete, whence a simple example of a chain that is not representable is given by the set of natural numbers with the usual order.

Grätzer’s problem is connected to the problem of determining when a profinite bounded distributive lattice A is isomorphic to the profinite completion of some bounded distributive lattice B , as we proceed to explain. Utilizing Priestley duality, it was shown in [7, Thm. 4.4] that A is a profinite bounded distributive lattice if and only if A is isomorphic to the lattice $\text{Up}(X)$ of all upsets (upward closed sets) of a poset X . On the other hand, by [10, Thm. 5.3], A is isomorphic to \widehat{B} for some bounded distributive lattice B if and only if A is isomorphic to $\text{Up}(X)$ for some representable poset X . Consequently, if the poset X is not representable and $A = \text{Up}(X)$, then A is not isomorphic to \widehat{B} for any bounded distributive lattice B [10, Cor. 5.4]. This allows us to construct easily profinite bounded distributive lattices A that are not isomorphic to profinite completions of bounded distributive lattices; for instance, take $A = \text{Up}(X)$ where X is any chain that is not complete.

In this paper we are interested in Heyting algebras. These are special bounded distributive lattices in which the meet operation has an adjoint, usually referred to as implication. Heyting algebras have been studied extensively as they have applications to different branches of mathematics, including:

- **Logic** (algebraic models of intuitionistic logic are Heyting algebras),
- **Topology** (the lattice of open sets of every topological space is a Heyting algebra),
- **Point-free Topology** (each locale is a Heyting algebra),
- **Domain Theory** (each continuous distributive lattice is a Heyting algebra),
- **Topos Theory** (the subobject classifier of every topos is a Heyting algebra).
- **Lattice Theory/Universal Algebra** (every algebraic distributive lattice and hence the congruence lattice of every algebra in a congruence-distributive variety is a Heyting algebra).

Grätzer’s problem of representability was reformulated by Esakia for Heyting algebras in [28, Appendix A.5]. We call a poset X *Esakia representable* if X is isomorphic to the prime spectrum of some Heyting algebra. A poset X is *image-finite* if the upset $\uparrow x := \{y \in X : x \leq y\}$ is finite for each $x \in X$. For a poset X , let

$$X_{\text{fin}} = \{x \in X : \uparrow x \text{ is finite}\}$$

be the image-finite subposet of X . By [7, Thm. 3.6], a Heyting algebra A is profinite if and only if A is isomorphic to $\text{Up}(X)$ for some image-finite poset X . By [10, Thm. 5.8], a Heyting algebra A is isomorphic to \widehat{B} for some Heyting algebra B if and only if A is isomorphic to $\text{Up}(X_{\text{fin}})$ for some Esakia representable poset X . This implies that if there is an image-finite poset X that is not isomorphic to Y_{fin} for any Esakia representable poset Y , then $\text{Up}(X)$ is an example of a profinite Heyting algebra that is not isomorphic to the profinite completion of any Heyting algebra [10, Cor. 5.9]. Finding such an image-finite poset is nontrivial, and was left as an open problem in [10].

We will resolve this problem by providing many such examples. In fact, we will provide a characterization of all varieties of Heyting algebras whose profinite members are profinite completions of some Heyting algebras. This we do by proving that there exists a largest such variety. We denote this variety by DHA and term its members *diamond Heyting algebras*. To explain the terminology, DHA is a subvariety of the variety CHA of *cascade Heyting algebras* of Esakia [28, Appendix A.9]. We recall that

CHA is generated by *Boolean cascades*, i.e., algebras whose prime spectrum is a linear sum of finitely many antichains. If each antichain has at most two elements and we do not allow two “back-to-back” two-element antichains, then the poset can be viewed as a linear sum of “diamonds,” and we refer to it as a *diamond sequence*. Let DHA be the subvariety of CHA generated by the Boolean cascades whose prime spectra are diamond sequences. This provides motivation for denoting this variety by DHA and calling its members diamond Heyting algebras. We call prime spectra of diamond Heyting algebras *diamond systems*. Thus, diamond sequences are special diamond systems.

Our main result states that the profinite members of a variety V of Heyting algebras are isomorphic to profinite completions of some Heyting algebras (which can always be chosen from V) if and only if V is a subvariety of DHA (Theorem 6.1). Consequently, DHA is the largest variety of Heyting algebras whose profinite members are profinite completions.

Among various characterizations of diamond Heyting algebras that we give, one is of special interest. Let P_1, P_2, P_3 , and P_4 be the posets shown in Figure 1.¹ The Heyting algebras $Up(P_i)$ of upsets of these posets are shown in Figure 2. We prove that the variety of diamond Heyting algebras is axiomatized by the Jankov formulas of $Up(P_i)$ for $i = 1, \dots, 4$. Consequently, the problem of determining whether a variety of Heyting algebras is such that its profinite members are profinite completions of some Heyting algebras is decidable, both for varieties presented by a finite set of equations and for varieties presented by a finite number of finite algebras (Theorem 7.1).




FIGURE 1. The posets P_1, P_2, P_3 , and P_4 .




FIGURE 2. The Heyting algebras of upsets of P_1, P_2, P_3 , and P_4 .

Our methods are based on a marriage of combinatorics of infinite posets, topology, and algebra. This is made possible by the use of Esakia duality [26, 28] that allows us to study Heyting algebras through the lenses of certain ordered topological spaces (known as Esakia spaces), which in turn are amenable to combinatorial principles such as Brouwer’s Fan Theorem (see, e.g., [24, Thm. 3.3.20]). Our main insight

¹In lattice theory it is customary to denote the posets P_1 and P_2 by M_3 and N_5 .

is a careful analysis of the structure of the diamond systems that can be endowed with a Stone topology so that the resulting structures are Esakia spaces.

Some of our results have purely logical formulation. We recall that an *intermediate logic* is a consistent axiomatic extension of intuitionistic propositional logic (see, e.g., [18, Sec. 2]). These are exactly the logics that are situated between intuitionistic and classical logics and are algebraized by nontrivial varieties of Heyting algebras [14]. Intermediate logics algebraized by varieties of diamond Heyting algebra share many properties with the well-known Gödel-Dummett logic (see, e.g., [23, Sec. 5.1] and [36, Sec. 4.2]). On the one hand, they are all finitely axiomatizable, locally tabular, and form a countable set (see Theorem 7.2). On the other hand, they are all structurally complete [60] and have the infinite Beth definability property [16] (see Theorem 7.4).

Finally, our results give a new insight on the representability problem as follows. Call a variety V of Heyting algebras *representable* if $\text{Up}(X) \in V$ implies that X is Esakia representable. We give a full characterization of representable varieties by proving that a variety V is representable if and only if $V \subseteq \text{DHA}$ and the depth of V is bounded by some positive integer (see Theorem 7.6). Consequently, every diamond system of bounded depth is Esakia representable. It follows that if a poset of finite depth validates the four axioms defining DHA, then it is Esakia representable. Since the axiomatization of DHA is given by Jankov formulas of the Heyting algebras of the posets P_1, \dots, P_4 (see Figures 1 and 2), we obtain that if a poset of finite depth does not contain any of the posets P_1, \dots, P_4 as a forbidden configuration (a p -morphic image of an upset), then it is Esakia representable. In particular, every *root system* (that is, a poset whose principal upsets are chains) [54] of bounded depth is Esakia representable. It remains an interesting open problem to give a full characterization of Esakia representable posets of finite depth. A solution of this problem would shed further new light on the difficult problem of Esakia representability.

2. HEYTING ALGEBRAS AND ESAKIA SPACES

We start by recalling that a bounded distributive lattice A is a *Heyting algebra* if \wedge has an adjoint \rightarrow given by

$$a \wedge b \leq c \iff a \leq b \rightarrow c$$

for all $a, b, c \in A$.

It is well known (see, e.g., [59, 4, 28]) that the class HA of Heyting algebras forms a variety (i.e., it is closed under homomorphic images, subalgebras, and direct products) and hence is equationally definable by Birkhoff's Theorem (see, e.g., [17, Thm. II.11.9]).

2.1. Esakia duality. The celebrated *Esakia duality* [26, 28] provides a representation of Heyting algebras by means of special ordered Stone spaces, and can be viewed as a restricted version of *Priestley duality* [56, 57] for bounded distributive lattices.

For a poset $X = \langle X, \leq \rangle$ and $U \subseteq X$, let $\uparrow U$ and $\downarrow U$ be the smallest upset and downset containing U , i.e.,

$$\begin{aligned} \uparrow U &= \{x \in X : \exists u \in U \text{ with } u \leq x\} \\ \downarrow U &= \{x \in X : \exists u \in U \text{ with } x \leq u\} \end{aligned}$$

We call U an *upset* if $U = \uparrow U$ and a *downset* if $U = \downarrow U$. If $U = \{x\}$, we simply write $\uparrow x$ and $\downarrow x$ instead of $\uparrow\{x\}$ and $\downarrow\{x\}$.

Definition 2.1. An *Esakia space* is a triple $X = \langle X, \tau, \leq \rangle$ where $\langle X, \tau \rangle$ is a Stone space (compact, Hausdorff, and zero-dimensional) and \leq is a partial order on X that is *continuous*; meaning that

- (i) $\uparrow x$ is closed for all $x \in X$;
- (ii) if $U \subseteq X$ is clopen, then $\downarrow U$ is clopen.

Given an Esakia space X , we denote its underlying poset by X as well.

Remark 2.2. The partial order \leq is continuous if and only if the corresponding map $\rho : X \rightarrow \mathcal{V}X$ from X to the Vietoris space $\mathcal{V}X$, given by $\rho(x) = \uparrow x$, is a well-defined continuous map (see [26, 1, 46]).

As usual, we denote by $\text{cl}(Y)$ the closure of a subset Y of a topological space. Also, we denote the set of maximal elements of a poset X by $\max X$. The following fundamental properties of Esakia spaces will be used repeatedly in the paper.

Proposition 2.3. *The following conditions hold for an Esakia space X .*

- (i) Priestley separation axiom: for all $x, y \in X$ such that $x \not\leq y$ there is a clopen upset U such that $x \in U$ and $y \notin U$.
- (ii) Dedekind completeness: Every nonempty chain $C \subseteq X$ has an infimum and a supremum in X .
- (iii) If C is a closed subset of X , then for every $x \in C$ there is $y \in \max C$ such that $x \leq y$.
- (iv) For every $Y \subseteq X$, $\uparrow \text{cl}(Y) = \text{cl}(\uparrow Y)$. Consequently, the closure of an upset is an upset and principal upsets are closed.
- (v) The downset of a closed set is closed. Consequently, principal downsets are closed.

Proof sketch. Conditions (i), (ii), and (iii) are respectively [28, Thms. 3.2.22(1), 3.2.19, and 3.2.1], while Conditions (iv) and (v) follow from [28, Thm. 3.1.2]. \square

In view of the above result, Esakia spaces satisfy Priestley separation axiom. In fact, Esakia spaces are those *Priestley spaces* (compact ordered spaces satisfying the Priestley separation axiom) in which the downset of each (cl)open is (cl)open [28, Thm. 3.1.2].

We recall that a *p-morphism* (or *bounded morphism*) between two posets X and Y is a map $f: X \rightarrow Y$ such that $\uparrow f(x) = f(\uparrow x)$ for each $x \in X$.

Definition 2.4. An *Esakia morphism* between Esakia spaces is a continuous p-morphism. Let ES be the category of Esakia spaces and Esakia morphisms between them.

Theorem 2.5 (Esakia duality). *HA is dually equivalent to ES.*

We briefly describe the contravariant functors $(-)_* : \text{HA} \rightarrow \text{ES}$ and $(-)^* : \text{ES} \rightarrow \text{HA}$ establishing Esakia duality. For this we first recall that for a poset X , the set $\text{Up}(X)$ of all upsets of X forms a Heyting algebra where join and meet are set-theoretic union and intersection, and \rightarrow is defined by

$$U \rightarrow V = X \setminus \downarrow(U \setminus V) = \{x \in X : \uparrow x \cap U \subseteq V\}.$$

For a Heyting algebra A , let X_A be the poset of prime filters of A ordered by inclusion. Define $\gamma_A : A \rightarrow \text{Up}(X_A)$ by

$$\gamma_A(a) = \{x \in X_A : a \in x\}.$$

Then $A_* = \langle X_A, \tau, \subseteq \rangle$ is an Esakia space, where τ is the topology on X_A given by the subbasis

$$\{\gamma_A(a) : a \in A\} \cup \{\gamma_A(a)^c : a \in A\}.$$

If $\alpha : A \rightarrow B$ is a homomorphism between Heyting algebras, define $\alpha_* : B_* \rightarrow A_*$ by setting $\alpha_*(U) = \alpha^{-1}(U)$ for all $U \subseteq B_*$. Then α_* is an Esakia morphism, and this defines the contravariant functor $(-)_* : \text{HA} \rightarrow \text{ES}$.

For an Esakia space X , let X^* be the subalgebra of $\text{Up}(X)$ consisting of clopen upsets of X . If $f : X \rightarrow Y$ is an Esakia morphism between Esakia spaces, define $f^* : Y^* \rightarrow X^*$ by $f^*(U) = f^{-1}(U)$ for all $U \in Y^*$. Then f^* is a homomorphism between the Heyting algebras Y^* and X^* , and this defines the contravariant functor $(-)^* : \text{ES} \rightarrow \text{HA}$.

The topology of a finite Esakia space is discrete (since it is Hausdorff). Therefore, the full subcategory of ES consisting of finite Esakia spaces is isomorphic to the category of finite posets and p-morphisms between them. Consequently, Esakia duality restricts to the following:

Theorem 2.6 (Finite Esakia duality). *The category of finite Heyting algebras is dually equivalent to the category of finite posets and p-morphisms between them.*

In addition, Stone spaces can be identified with the Esakia spaces whose underlying order is the identity. Under this identification, Esakia duality restricts to the famous *Stone duality* [65, 66] between the categories of Boolean algebras and Stone spaces.

To characterize homomorphic images and subalgebras of a Heyting algebra, we recall the notions of Esakia subspaces and Esakia quotients.

Definition 2.7. Let X be an Esakia space.

- (1) An *Esakia subspace* (*E-subspace* for short) is a closed upset of X equipped with the subspace topology and the restriction of the order.
- (2) An *Esakia equivalence* (or *E-partition* for short) is an equivalence relation R on X satisfying for all $x, y, z \in X$:
 - (i) if xRy and $y \leq z$, then $x \leq w$ and wRz for some $w \in X$;
 - (ii) if xRy , then there is an *R-saturated* clopen U (a union of equivalence classes of R) such that $x \in U$ and $y \notin U$.

Remark 2.8. E-partitions are sometimes called correct partitions or bisimulation equivalences [29, 12].

It is well known that E-subspaces of Esakia spaces are Esakia spaces [28, Lem. 3.4.11]. In particular, closed subspaces of Stone spaces are Stone spaces.

If R is an E-partition on X , we denote by X/R the Esakia space consisting of the quotient space of X with respect to R , equipped with the partial order \leq_R defined as follows for every $x, y \in X$:

$$\llbracket x \rrbracket \leq_R \llbracket y \rrbracket \iff \text{there are } x' \in \llbracket x \rrbracket \text{ and } y' \in \llbracket y \rrbracket \text{ such that } x' \leq y'.$$

The map $x \mapsto \llbracket x \rrbracket$ is an Esakia morphism from X to X/R , and for every Esakia morphism $f: X \rightarrow Y$, the kernel of f is an E-partition of X [12, Thm. 2.3.9].

An algebra A is *subdirectly irreducible* (SI for short) if the identity relation is completely meet-irreducible, and A is *finitely subdirectly irreducible* (FSI for short) if the identity relation is meet-irreducible in the congruence lattice of A (see, e.g., [5, 17]).

The following lemma collects some well-known consequences of Esakia duality.

Lemma 2.9. *Let A be a Heyting algebra.*

- (i) A is FSI iff its top element 1 is prime ($x \vee y = 1$ implies $x = 1$ or $y = 1$), which happens iff A_* is rooted (has a least element).
- (ii) The congruence lattice of A is dually isomorphic to the lattice of E-subspaces of A_* .
- (iii) The lattice of subalgebras of A is dually isomorphic to the lattice of E-partitions of A_* .

When Condition (i) holds, $\{1\}$ is the least prime filter of A [28, Prop. A.1.1]. For Conditions (ii) and (iii) see, e.g., [28, Thm. 3.4.16] and [12, Thm. 2.3.9].

2.2. Depth and width in Heyting algebras. We next recall how to define finite depth and width for Esakia spaces and Heyting algebras, and prove that the set of elements of depth $\leq n$ (resp. width $\leq n$) of an Esakia space forms an E-subspace.

Definition 2.10. Let X be a poset, n a positive integer, and $x \in X$.

- (1) X is said to have *depth at most n* if it does not contain any chain of $n + 1$ elements.
- (2) $x \in X$ is said to be of *depth $\leq n$* if the subposet $\uparrow x$ has depth $\leq n$.
- (3) $x \in X$ is said to be of *width $\leq n$* if $\uparrow x$ does not contain any antichain of $n + 1$ elements.
- (4) X is said to have *width at most n* if every $x \in X$ is of width $\leq n$.

Definition 2.11. Let A be a Heyting algebra and n a positive integer.

- (1) A has *depth at most n* if A_* has depth at most n . When this is the case, we write $d(A) \leq n$.
- (2) A has *depth exactly n* if $d(A) \leq n$ and $d(A) \not\leq n - 1$.
- (3) A has *width at most n* if A_* has width at most n . When this is the case, we write $w(A) \leq n$.
- (4) A has *width exactly n* if $w(A) \leq n$ and $w(A) \not\leq n - 1$.

The notion of depth originates in [38], while width was introduced for modal logics above K4 in [30], and was adapted to Heyting algebras in [63].

Given a Heyting algebra A and a formula φ , we write $A \models \varphi$ as a shorthand for $A \models \varphi \approx 1$. In this case, we say that φ is *valid* in A . The following result is well known.

Theorem 2.12. *Let A be a Heyting algebra and n a positive integer.*

(i) $d(\mathbf{A}) \leq n$ the formula d_n is valid in \mathbf{A} , where

$$\begin{aligned} d_1 &= p_1 \vee (p_1 \rightarrow 0) \\ d_{m+1} &= p_{m+1} \vee (p_{m+1} \rightarrow d_m), \text{ for all } m \geq 1. \end{aligned}$$

Consequently, the class \mathbf{D}_n of Heyting algebras of depth at most n is a variety.

(ii) $w(\mathbf{A}) \leq n$ iff the formula w_n is valid in \mathbf{A} , where

$$w_n = \bigvee_{i=0}^n (p_i \rightarrow \bigvee_{j \neq i} p_j).$$

Consequently, the class \mathbf{W}_n of Heyting algebras of width at most n is a variety.

Proof sketch. Condition (i) was essentially established in [55, 49] and Condition (ii) in [63]. \square

Heyting algebras of width at most one are called *Gödel algebras* [36, Sec. 4.2]. In view of the above result, they form a variety which, moreover, algebraizes the Gödel-Dummett logic [25]. Posets of width at most one have been called *root systems* and coincide with disjoint unions of posets whose order duals are trees [54].

Proposition 2.13. *Let X be an Esakia space and n a positive integer.*

- (i) *The set of points of X of depth $\leq n$ is an E-subspace of X .*
- (ii) *The set of points of X of width $\leq n$ is an E-subspace of X .*

Proof. Condition (i) originates in [6, Lem. 7]. Thus, we only give a proof of (ii). Let $X_{\leq n}$ be the set of points of X of width $\leq n$. Since $x \leq y$ implies $\uparrow y \subseteq \uparrow x$, it is clear that $x \in X_{\leq n}$ implies $y \in X_{\leq n}$. Therefore, $X_{\leq n}$ is an upset of X . Thus, it is sufficient to show that $X_{\leq n}$ is closed in X . Since $X_{\leq n}$ is an upset, so is its closure $\text{cl}(X_{\leq n})$ by Proposition 2.3(iv). Thus, $\text{cl}(X_{\leq n})$ is an E-subspace of X .

Let \mathbf{A} be the Heyting algebra of clopen upsets of $\text{cl}(X_{\leq n})$. For each $x \in X$, we have that $\uparrow x$ is an E-subspace of X by Proposition 2.3(iv). Let \mathbf{A}_x be the Heyting algebra of clopen upsets of $\uparrow x$. Since for each $x \in X_{\leq n}$ we have $\uparrow x \subseteq X_{\leq n}$, we see that $\mathbf{A}_x \in \mathbf{W}_n$. Moreover, since $X_{\leq n}$ is dense in $\text{cl}(X_{\leq n})$, the Heyting algebra \mathbf{A} is a subalgebra of the product of $\{\mathbf{A}_x : x \in X_{\leq n}\}$ [28, Appendix A.1]. Therefore, $\mathbf{A} \in \mathbf{W}_n$. Thus, the width of $\text{cl}(X_{\leq n})$ is $\leq n$ by Theorem 2.12(ii). Consequently, $\text{cl}(X_{\leq n}) = X_{\leq n}$, and hence $X_{\leq n}$ is closed in X .

Notice that, by interpreting $X_{\leq n}$ as the set of all elements of X of depth $\leq n$, the above argument becomes a proof of (i). \square

Remark 2.14. The above result fails for Priestley spaces. To see this, let X be the one-point compactification of an infinite discrete space Y . We endow X with a partial order by setting $x \leq y$ if and only if $x = y$ or x is the unique limit point of Y . It is well known and easy to check that the resulting ordered topological space is a Priestley space in which $\max X$ is dense. Thus, $\max X$ is the set of points of X of both depth and width 1. But its closure is X , whose depth is 2 and width is infinite.

We next recall the following definition from the introduction.

Definition 2.15. For an Esakia space X , let $X_{\text{fin}} = \langle X_{\text{fin}}, \leq \rangle$ be the subposet of X where

$$X_{\text{fin}} = \{x \in X : \uparrow x \text{ is finite}\}.$$

We call X_{fin} the *image-finite part* of X .

The next corollary to Proposition 2.13 will be used throughout the paper.


Corollary 2.16. *Let X be a poset of depth and width $\leq n$ for some positive integer n . If there exists an Esakia space whose image-finite part is X , then there is a Stone topology τ on X such that $\langle X, \tau \rangle$ is an Esakia space.*

Proof. Suppose that X is the image-finite part of an Esakia space Y . Let Z be the set of elements of Y of depth and width $\leq n$. By Proposition 2.13, Z is an E-subspace of Y . Therefore, it is sufficient to show that $X = Z$. By assumption, X has depth and width $\leq n$. Since X is an upset of Y , we see that $X \subseteq Z$. To prove the other inclusion, let $z \in Z$. Since z has depth and width $\leq n$ in Y , it belongs to the image-finite part of Y , so $z \in X$. Thus, $X = Z$. \square

2.3. Profinite algebras and completions. We recall that an *inverse system* of Heyting algebras is a family $\{A_i : i \in I\}$, indexed by a directed poset I , together with a family of homomorphisms $\alpha_{ij} : A_j \rightarrow A_i$ for $i \leq j$, satisfying:

- (i) $\alpha_{kj} = \alpha_{ki} \circ \alpha_{ij}$ for all $k \leq i \leq j$;
- (ii) α_{ii} is the identity homomorphism for each $i \in I$.

The *inverse limit* of such an inverse system is a Heyting algebra A together with a family $\alpha_i : A \rightarrow A_i$ of homomorphisms satisfying $\alpha_{ij} \circ \alpha_j = \alpha_i$ for each $i \leq j$, and having the following universal mapping property: For each Heyting algebra B and each family of homomorphisms $\beta_i : B \rightarrow A_i$ satisfying $\alpha_{ij} \circ \beta_j = \beta_i$ for each $i \leq j$, there is a unique homomorphism $\beta : B \rightarrow A$ such that $\alpha_i \circ \beta = \beta_i$ for each $i \in I$.



It is well known that A is isomorphic to the subalgebra of the product $\prod_{i \in I} A_i$ consisting of

$$\left\{ \vec{a} \in \prod_{i \in I} A_i : \alpha_{ij}(\vec{a}(j)) = \vec{a}(i) \text{ for each } j \geq i \right\}$$

For inverse limits see, e.g., [2, Ex. 13.13].

Definition 2.17. We call a Heyting algebra A *profinite* if it is isomorphic to the inverse limit of an inverse system of finite Heyting algebras.

Let A be a Heyting algebra and θ a congruence on A . We say θ has *finite index* if the quotient algebra A/θ is finite. Let I be the set of congruences of A of finite index. If $\theta \subseteq \phi$ are congruences of A , then there is a canonical homomorphism $\pi_{\phi\theta} : A/\theta \rightarrow A/\phi$ given by $\pi_{\phi\theta}(\llbracket a \rrbracket_{\theta}) = \llbracket a \rrbracket_{\phi}$. Furthermore, if θ and ϕ have finite index, then so does $\theta \cap \phi$ because $A/(\theta \cap \phi)$ embeds into $A/\theta \times A/\phi$ via the map given by

$$\llbracket a \rrbracket_{\theta \cap \phi} \mapsto \langle \llbracket a \rrbracket_{\theta}, \llbracket a \rrbracket_{\phi} \rangle,$$

see, e.g., [17, Ch. II, Lem. 8.2]. Therefore, (I, \supseteq) is a directed poset, and $\{A/\theta : \theta \in I\}$ endowed with the homomorphisms $\pi_{\phi\theta}$ is an inverse system.

Definition 2.18. The *profinite completion* \widehat{A} of a Heyting algebra A is the inverse limit of the inverse system described above.

Given a class K of similar algebras, we denote by $\mathbb{I}(K)$, $\mathbb{H}(K)$, $\mathbb{S}(K)$, and $\mathbb{P}(K)$ the classes of isomorphic copies, homomorphic images, subalgebras, and direct products of elements of K , respectively. When $K = \{A\}$, we write $\mathbb{I}(A)$ as a shorthand for $\mathbb{I}(\{A\})$. The same convention applies to \mathbb{H} , \mathbb{S} , and \mathbb{P} .

Remark 2.19. If a Heyting algebra A is a profinite completion of some Heyting algebra B , then A is also the profinite completion of some Heyting algebra C that belongs to the variety generated by A ; namely, $\mathbb{HSP}(A)$. This is because A is also the profinite completion of $C := B/\theta$, where θ is the intersection of all the congruences $\{\theta_i : i \in I\}$ of B of finite index. Since $B/\theta_j \in \mathbb{H}(A)$ for all $j \in I$ and $B/\theta \in \mathbb{ISP}(\{B/\theta_i : i \in I\})$, we obtain that $C \in \mathbb{ISP}\mathbb{H}(A) \subseteq \mathbb{HSP}(A)$.

By [9], A is a profinite Heyting algebra if and only if there is a compact Hausdorff topology on A under which the Heyting operations are continuous. We will rely on the following characterization of profinite Heyting algebras and profinite completions of Heyting algebras.

Theorem 2.20. *Let A be a Heyting algebra and X its Esakia dual.*

- (i) [7, Thm. 3.6] *A is a profinite Heyting algebra if and only if there is an image-finite poset Y such that $A \cong \text{Up}(Y)$.*
- (ii) [8, Thm. 4.7] *\hat{A} is isomorphic to $\text{Up}(X_{\text{fin}})$.*

Clearly each profinite completion is a profinite Heyting algebra. However, as we will see, there are profinite Heyting algebras that are not isomorphic to the profinite completion of any Heyting algebra.

3. CASCADE HEYTING ALGEBRAS

In this section we recall cascade Heyting algebras and their dual description. We also give their axiomatization via Jankov formulas.

Let X_1, \dots, X_n be Esakia spaces. We recall that the *linear sum* $X_1 \oplus \dots \oplus X_n$ is defined as the disjoint union of the spaces X_1, \dots, X_n , where the partial order is given by

$$x \leq y \iff (x, y \in X_i \text{ and } x \leq_i y \text{ for some } i \leq n) \text{ or } (x \in X_i \text{ and } y \in X_j \text{ for some } j < i \leq n).$$

Figuratively speaking, we are forming a tower by putting X_n, \dots, X_1 on top of each other. We emphasize that X_1 is at the top of the tower and X_n is at the bottom. It is well known that a linear sum of Esakia spaces is again an Esakia space [28, Prop. A.8.6]. If X is the linear sum of X_1, \dots, X_n and A_1, \dots, A_n are dual Heyting algebras to X_1, \dots, X_n , then the dual Heyting algebra A of X is obtained by putting the algebras A_1, \dots, A_n on top of each other and identifying the top element of A_i with the bottom element of A_{i+1} (see, e.g., [53, Lem. 5.1]).

If each X_i is a Stone space, then the linear sum $X = X_1 \oplus \dots \oplus X_n$ is a tower of antichains. Moreover, the dual of each X_i is a Boolean algebra. Consequently, Esakia called a Heyting algebra whose dual Esakia space is a linear sum of Stone spaces a *Boolean cascade* [28, Appendix A.9].

Definition 3.1. Let CHA be the variety of Heyting algebras generated by Boolean cascades.

Following Esakia [28, Appendix A.9], we call members of CHA *cascade Heyting algebras*.

Theorem 3.2. *A Heyting algebra A is a cascade Heyting algebra iff it validates the weak Pierce law:*

$$(p \rightarrow q) \vee (((q \rightarrow p) \rightarrow q) \rightarrow q).$$

Thus, the logic WPL of the weak Pierce law is algebraized by the variety CHA of cascade Heyting algebras.

Proof sketch. This result was proved in [39, 27]; see also [28, Prop. A.9.1(1)]. A slightly different axiomatization of cascade Heyting algebras is given in [51, Thm. 4]. □

An algebra is said to be *locally finite* if its finitely generated subalgebras are finite. Accordingly, a variety is called *locally finite* if all its members are locally finite. The following theorem originates with [50, 27]; see also [28, Prop. A.9.1(1)]. A stronger result was proved by Kuznetsov [47]; see, e.g., [20, Thm. 2].

Theorem 3.3. *CHA is locally finite.*

In view of the above result and the fact that the class of Boolean cascades is closed under subalgebras, CHA is generated by finite Boolean cascades. Hence, for many purposes, it is sufficient to work with linear sums of finite antichains.

Several useful characterizations of cascade Heyting algebras were given by Esakia [28] and Mardaev [51]. The Mardaev characterization utilizes the technique of Jankov formulas [40, 41, 42]. We recall that with each finite SI Heyting algebra A , we can associate the *Jankov formula* $\mathcal{J}(A)$ of A , which axiomatizes the largest variety of Heyting algebras omitting the algebra A . By Theorem 2.6 and Lemma 2.9(i), finite SI algebras correspond to finite rooted posets. Because of this, given a finite rooted poset X , we denote by $\mathcal{J}(X)$ the Jankov formula of the finite SI algebra $\text{Up}(X)$.

Lemma 3.4 (Jankov's Lemma). *Let A, B be Heyting algebras with A finite and SI. Then $B \models \mathcal{J}(A)$ if and only if $A \notin \text{SIH}(B)$.*

In view of Lemma 2.9, Jankov's Lemma can be rephrased as follows.

Lemma 3.5. *Let X, Y be Esakia spaces with X finite and rooted. Then $Y^* \models \mathcal{J}(X)$ if and only if X is not a p -morphic image of an E -subspace of Y .*

Definition 3.6 ([28, pg. 86]). A poset X is said to satisfy the *three point rule* provided for every distinct $x, y, z \in X$, if x and y are incomparable, then $x \leq z$ implies $y \leq z$.

In the following theorem, P_5 and P_6 are the posets depicted in Figure 3.




FIGURE 3. The posets P_5 and P_6 .

Theorem 3.7. *For a Heyting algebra A , the following are equivalent.*

- (i) A is a cascade Heyting algebra;
- (ii) $\uparrow x$ satisfies the three point rule for each $x \in A_*$;
- (iii) A validates the Jankov formulas $\mathcal{J}(P_2)$, $\mathcal{J}(P_5)$, and $\mathcal{J}(P_6)$.

Proof sketch. The equivalence of Conditions (i) and (ii) goes back to [27]; see also [28, Prop. A.9.2]. The equivalence of Conditions (i) and (iii) is a consequence of [51, Thm. 5] and the fact that CHA is generated by Boolean cascades. \square

Let $\text{CHA}_n = \text{CHA} \cap W_n$. In other words, CHA_n is the subvariety of CHA consisting of cascade Heyting algebras of width $\leq n$. We conclude this section by showing that CHA_n can be axiomatized by the Jankov formulas of the “ $(n+1)$ -fork” F_{n+1} and “ $(n+1)$ -diamond” D_{n+1} . For every positive integer m , let F_m and D_m be the finite rooted posets depicted in Figure 4.




FIGURE 4. The posets F_m and D_m .

Our proof is based on the following two lemmas.

Lemma 3.8. *If $A \in W_n$, then A validates $\mathcal{J}(F_{n+1})$ and $\mathcal{J}(D_{n+1})$.*

Proof. Since $A \in W_n$ and W_n is a variety, each $B \in \text{SIH}(A)$ is also in W_n . Therefore, each such B has width $\leq n$. On the other hand, the width of both $\text{Up}(F_{n+1})$ and $\text{Up}(D_{n+1})$ is $n+1$. Therefore, neither belong to $\text{SIH}(A)$. Thus, by Jankov's Lemma, A validates $\mathcal{J}(F_{n+1})$ and $\mathcal{J}(D_{n+1})$. \square

Lemma 3.9. *If A is a finite subdirectly irreducible cascade Heyting algebra and $A \notin W_n$, then A refutes either $\mathcal{J}(F_{n+1})$ or $\mathcal{J}(D_{n+1})$.*

Proof. By finite Esakia duality, since A is finite and subdirectly irreducible, we may assume without loss of generality that A is $\text{Up}(X)$ for some finite rooted poset X . In addition, since A is a finite cascade Heyting algebra and X is rooted, X satisfies the three point rule by Theorem 3.7. It follows that X is a linear sum of finite antichains. Since $A \notin W_n$ and X is rooted, there is an $(n+1)$ -element antichain C in X . Then C is a subset of one of the antichains whose linear sum is X . Let x be an immediate predecessor of the elements of C and let $Q = \uparrow x$. Then Q is an upset of X . If $C \subseteq \max X$, then Q is isomorphic to F_{n+1} , so $\text{Up}(F_{n+1}) \in \mathbf{H}(A)$. Therefore, by Jankov's Lemma, $A \not\equiv \mathcal{J}(F_{n+1})$. Otherwise, define an equivalence relation R on Q by identifying all $y \notin C \cup \{x\}$. It is straightforward to check that R is an E-partition of Q and that Q/R is isomorphic to D_{n+1} . Therefore, $\mathcal{J}(D_{n+1}) \in \mathbf{SIH}(A)$. Thus, by Jankov's Lemma, $A \not\equiv \mathcal{J}(D_{n+1})$. \square

Putting together Lemmas 3.8 and 3.9 yields the following:

Theorem 3.10. CHA_n is axiomatized over CHA by the Jankov formulas $\mathcal{J}(F_{n+1})$ and $\mathcal{J}(D_{n+1})$.

Proof. It follows from Lemmas 3.8 and 3.9 that CHA_n and the subvariety of CHA axiomatized by $\mathcal{J}(F_{n+1})$ and $\mathcal{J}(D_{n+1})$ have the same finite subdirectly irreducible members. The result follows since CHA is locally finite by Theorem 3.3. \square

Remark 3.11. The description of algebras of width $\leq n$ given in Theorem 3.10 cannot be extended to arbitrary Heyting algebras. For instance, the Heyting algebra of upsets of the poset P_7 depicted in Figure 5 validates $\mathcal{J}(F_3)$ and $\mathcal{J}(D_3)$ but is not of width ≤ 2 . Consequently, W_n is not axiomatized over HA by $\mathcal{J}(F_{n+1})$ and $\mathcal{J}(D_{n+1})$.




FIGURE 5. The poset P_7 .

4. DIAMOND SYSTEMS AND ALGEBRAS

In this section we introduce the key concept of diamond systems and the corresponding variety DHA of diamond Heyting algebras. We prove that DHA is finitely axiomatizable by Jankov formulas.

Definition 4.1. We call a poset X a *diamond system* if it satisfies the following conditions:

- (D1) $\uparrow x$ satisfies the three point rule for each $x \in X$;
- (D2) X has width at most two;
- (D3) Principal upsets are upward directed in X ;
- (D4) For every $\perp, x, y, z, v, \top \in X$, if $\perp \leq x, y \leq z, v \leq \top$, there is $w \in X$ such that

$$x, y \leq w \leq z, v.$$

If X is in addition downward directed, then we call it a *diamond sequence*.

Notice that finite diamond sequences are simply linear sums of antichains of size ≤ 2 in which we do not allow two “back-to-back” antichains. Typical examples of diamond sequences are displayed in Figure 6, thus justifying the name. An image-finite diamond system that is not a diamond sequence is depicted in Figure 7. Observe that root systems coincide with diamond systems of width at most one.

Image-finite diamond systems admit a simple characterization in terms of linear sums.

Proposition 4.2. An image-finite poset X is a diamond system if and only if for every $x \in X$ there are posets Y_1, \dots, Y_n such that $\uparrow x$ is order-isomorphic to $Y_1 \oplus \dots \oplus Y_n$, where Y_1 is a singleton, and for all $j > 1$, either Y_j is a singleton or isomorphic to P_3 .




FIGURE 6. Two examples of diamond sequences.




FIGURE 7. An image-finite diamond system.

Proof. The “if” part is straightforward. To prove the “only if” part, consider an image-finite diamond system X and an element $x \in X$. Then $\uparrow x$ is finite since X is image-finite. Therefore, by (D1), $\uparrow x$ can be partitioned into a finite family of disjoint nonempty sets X_1, \dots, X_m such that $X_1 = \{x\}$ and for every $y, z \geq x$,

$$y \leq z \iff \text{either } y = z \text{ or } (y \in X_i \text{ and } z \in X_j \text{ for some } i < j).$$

Notice that, as X has width at most two by (D2), each X_i has at most two elements. Furthermore, if X_i has two elements, then $i - 1$ is positive, since X_1 is a singleton. Together with Condition (D4), this implies that if X_i has two elements, then X_{i-1} exists and has one element. Thus, there are posets Y_1, \dots, Y_n such that $\uparrow x$ is order-isomorphic to the sum $Y_1 \oplus \dots \oplus Y_n$ where each Y_i is either a singleton or isomorphic to P_3 . To conclude the proof, it only remains to show that Y_1 is a singleton. But this is a consequence of the fact that X is upward directed by (D3). \square

Notice that the assumption that X is image-finite in the above result is essential since the chain of natural numbers (or the second poset in Figure 6) are diamond systems that do not satisfy Proposition 4.2.

Definition 4.3. Let DHA be the variety of Heyting algebras generated by the algebras of upsets of finite diamond sequences. The elements of DHA will be called *diamond Heyting algebras*.

Notice that, by definition, DHA is a subvariety of CHA. Furthermore, since the variety GA of Gödel algebra is generated by the algebras of upsets of finite chains, which are diamond sequences, we obtain that GA is a subvariety of DA.

Diamond systems and algebras are related by the following theorem, the hardest implication of which is (iv) \Rightarrow (ii).

Theorem 4.4. *The following conditions are equivalent for a Heyting algebra A :*

- (i) A is a diamond Heyting algebra;
- (ii) A_* is a diamond system;

- (iii) Every principal upset in A_* is a diamond sequence;
- (iv) A validates the Jankov formulas $\mathcal{J}(P_1), \mathcal{J}(P_2), \mathcal{J}(P_3)$, and $\mathcal{J}(P_4)$.

Proof. (i) \Rightarrow (iv): Notice that, by definition, diamond Heyting algebras are cascade Heyting algebras and have width ≤ 2 . Hence, by Theorem 3.7 and Lemma 3.10, they validate $\mathcal{J}(P_1)$ and $\mathcal{J}(P_2)$. It only remains to prove that diamond Heyting algebras validate $\mathcal{J}(P_3)$ and $\mathcal{J}(P_4)$. In view of Lemma 3.5, it suffices to show that P_3 and P_4 cannot be obtained as p-morphic images of upsets of finite diamond sequences. To this end, let X be a finite diamond sequence. If X is empty, we are done. Then suppose that $X \neq \emptyset$. In this case, X has a maximum, whence every nonempty p-morphic image of an upset of X has also a maximum. It follows that P_3 cannot be a p-morphic image of an upset of X . The fact that P_4 is not a p-morphic image of an upset of X follows from the fact that, by Proposition 4.2, X can be viewed as linear sum $Y_1 \oplus \cdots \oplus Y_n$, where Y_1 is a singleton, and for all $j > 1$, either Y_j is a singleton or isomorphic to P_3 .

(iv) \Rightarrow (ii): First we prove that A is a cascade Heyting algebra. To this end, observe that P_3 can be obtained as a p-morphic image of an upset of both P_5 and P_6 . Consequently, if a variety of Heyting algebras contains either $\text{Up}(P_5)$ or $\text{Up}(P_6)$, it also contains $\text{Up}(P_3)$. From the fact that A validates $\mathcal{J}(P_3)$ it follows that the variety generated by A omits $\text{Up}(P_3)$, whence it also omits $\text{Up}(P_5)$ and $\text{Up}(P_6)$. We obtain that the Jankov formulas $\mathcal{J}(P_5)$ and $\mathcal{J}(P_6)$ are valid in A . Recall also that, by assumption, A validates $\mathcal{J}(P_2)$. Hence, by Theorem 3.7, we conclude that A is a cascade Heyting algebra. From the same theorem it follows that principal upsets in A_* satisfy the three point rule. Thus, A_* satisfies Condition (D1).

We next prove that A_* satisfies Condition (D2), i.e., that it has width ≤ 2 . In view of Theorem 3.10 it suffices to prove that A validates $\mathcal{J}(F_3)$ and $\mathcal{J}(D_3)$. Observe that $D_3 = P_1$ and that A validates $\mathcal{J}(P_1)$ by assumption. On the other hand, P_3 is a p-morphic image of F_3 . It follows that if a variety of Heyting algebras contains $\text{Up}(F_3)$, then it also contains $\text{Up}(P_3)$. Since A validates $\mathcal{J}(P_3)$, the variety generated by A omits $\text{Up}(P_3)$. It follows that the variety generated by A also omits $\text{Up}(F_3)$, whence A validates $\mathcal{J}(F_3)$. Thus, A_* has width ≤ 2 .

The next two claims conclude the proof that A_* is a diamond system.

Claim 4.5. *Principal upsets are upward directed in A_* .*

Proof. Suppose not. Then there are $x, y, z \in A_*$ such that $x \leq y, z$ and $\uparrow y \cap \uparrow z = \emptyset$. As principal upsets are closed, we can apply Proposition 2.3(iii) to obtain that both $\uparrow y$ and $\uparrow z$ contain maximal elements. Since $\uparrow y \cap \uparrow z = \emptyset$, we may assume without loss of generality that these maximal elements are y and z . Similarly, as principal downsets are closed, Proposition 2.3(iii) implies that there is an element that is maximal in $\downarrow z \cap \downarrow y$. We may assume without loss of generality that this element is x . Thus, y and z are maximal and x is maximal among the lower bounds of y and z .

We show that

$$\uparrow x = \{x, y, z\}. \quad (1)$$

The inclusion from right to left is clear. Suppose, with a view to contradiction, that the other inclusion fails. Then there is $a > x$ different from y and z . Since A_* has width ≤ 2 and y and z are incomparable (since $\uparrow y \cap \uparrow z = \emptyset$), a is comparable either with y or z . By symmetry we may assume that a is comparable with y . Since y is maximal and $a \neq y$, we get $a < y$. Because $x < a < y$ and x is maximal among the lower bounds of y and z , we have $a \not\leq z$. As z is maximal, this implies that a and z are incomparable. Since $x \leq a, y, z$ and $a \leq y$ and a, y, z are different, the three point rule yields $a \leq z$, a contradiction.

Since principal upsets are closed in Esakia spaces, $\uparrow x$ is the universe of an E-subspace X of A_* . From (1) it follows that X is isomorphic to P_3 endowed with the discrete topology. Hence, by Lemma 2.9(ii), $\text{Up}(P_3) \in \mathbb{H}(A) \subseteq \text{DHA}$. But this contradicts the fact that A validates $\mathcal{J}(P_3)$. \square

Claim 4.6. *For every $\perp, x, y, z, v, \top \in A_*$, if $\perp \leq x, y \leq z, v \leq \top$, there is $w \in A_*$ such that $x, y \leq w \leq z, v$.*

Proof. Suppose, with a view to contradiction, that there are $\perp, x, y, z, v, \top \in A_*$ such that $\perp \leq x, y \leq z, v \leq \top$ and $\uparrow x \cap \uparrow y \cap \downarrow z \cap \downarrow v = \emptyset$. Notice that x and y must be incomparable, otherwise $\{x, y\} \cap \uparrow x \cap$

$\uparrow y \cap \downarrow z \cap \downarrow v \neq \emptyset$, a contradiction. Similarly, z is incomparable with v . Consequently, the elements \perp, x, y, z, v, \top are all different.

Recall that \perp is a lower bound of x and y . Therefore, since principal downsets are closed, Proposition 2.3(iii) implies the existence of a maximal lower bound of x and y . We may assume without loss of generality that this maximal lower bound is \perp . Bearing this in mind, we will prove that

$$\uparrow \perp = \{\perp\} \cup \uparrow x \cup \uparrow y. \quad (2)$$

The inclusion from right to left is clear. To prove the other inclusion, consider an element $w > \perp$ different from x and y . As A_* has width ≤ 2 , w must be comparable either with x or y . By symmetry we may assume that w is comparable with x . If $w \geq x$, then $w \in \{\perp\} \cup \uparrow x \cup \uparrow y$, as desired. Otherwise, $w < x$. We show this leads to a contradiction. Since \perp is a maximal lower bound of x and y and $\perp < w \leq x$, we get $w \not\leq y$. Moreover, $y \not\leq w$, since x and y are incomparable and $w \leq x$. Thus, x, y, w are different, y and w are incomparable and $w \leq x$. By the three point rule, $y \leq x$, a contradiction. This establishes (2).

Furthermore, observe that

$$\uparrow x \cup \uparrow y = \{x, y, z, v\} \cup (\uparrow z \cap \uparrow v). \quad (3)$$

The inclusion from right to left is clear. Suppose, with a view to contradiction, that the other inclusion fails. Then there is $w \in A_* \setminus \{x, y, z, v\}$ such that $w \in \uparrow x \cup \uparrow y$ and $w \notin \uparrow z \cap \uparrow v$. Since $w \in \uparrow x \cup \uparrow y$, by symmetry we may assume that $x \leq w$. Similarly, as $w \notin \uparrow z \cap \uparrow v$, by symmetry we may assume that $z \not\leq w$. Since A_* has width ≤ 2 , from $x \leq w, z, v$ and the fact that z and v are incomparable it follows that w is comparable either with z or v . We have two cases: either $w \in \downarrow z \cup \downarrow v$ or $w \notin \downarrow z \cup \downarrow v$. First suppose that $w \in \downarrow z \cup \downarrow v$. Since $w \notin \{x, z, v\}$, either $x < w < z$ or $x < w < v$. By symmetry, we may assume that $x < w < z$. Since x and y are incomparable, $\perp \leq x, y$, and $x < w$, we can apply the three point rule to obtain $y \leq w$. It follows that $w \in \uparrow x \cap \uparrow y \cap \downarrow z$. Together with the assumption that $\uparrow x \cap \uparrow y \cap \downarrow z \cap \downarrow v$ is empty, this yields $w \not\leq v$. Furthermore, from $w \leq z$ and the fact that z and v are incomparable it follows that $v \not\leq w$. Hence, w and v are also incomparable. Therefore, since $x \leq w, v$ and $w < z$, we can apply the three point rule to obtain $v < z$, a contradiction.

Next we consider the case where $w \notin \downarrow z \cup \downarrow v$. Together with the fact that w is comparable either with z or v and $z \not\leq w$, this implies $v < w$. Since z, v, w are distinct upper bounds of x such that z and v are incomparable and $v \leq w$, we can apply the three point rule to obtain $z \leq v$, a contradiction. This establishes (3).


From (2) and (3) it follows that

$$\uparrow \perp = \{\perp, x, y, z, v\} \cup (\uparrow z \cap \uparrow v). \quad (4)$$

Since $\uparrow \perp$ is a closed upset of A_* , it is the universe of an E-subspace X of A_* . Similarly, $\uparrow z \cap \uparrow v$ is closed in X . In view of (4), this implies that $\{\perp, x, y, z, v\}$ is open in X . As X is Hausdorff, we conclude that the points \perp, x, y, z, v are isolated in X . Furthermore, $\{\perp, x, y, z, v\}$ is closed because it is finite and X is Hausdorff. By (4) this yields that $\uparrow z \cap \uparrow v$ is open. Thus,

$$\{\perp\}, \{x\}, \{y\}, \{z\}, \{v\}, \uparrow z \cap \uparrow v \text{ are open in } X. \quad (5)$$

We denote the elements of the poset P_4 as follows:



Bearing this in mind, consider the map $f: X \rightarrow P_4$ defined by

$$f(a) = \begin{cases} k_a & \text{if } a \in \{\perp, x, y, z, v\} \\ k_{\top} & \text{if } a \in \uparrow z \cap \uparrow v. \end{cases}$$

Notice that f is a well-defined p-morphism by (4). Moreover, when P_4 is endowed with the discrete topology, f becomes an Esakia morphism. This is because $f^{-1}(k)$ is open in X for every $k \in P_4$ by (5). Thus, there is an E-partition R of X such that $X/R \cong P_4$. Together with the fact that X is an E-subspace of A_* , this implies that $\text{Up}(P_4) \in \mathbb{ISH}(A)$ by Lemma 2.9(ii, iii). But this contradicts the fact that A validates $\mathcal{J}(P_4)$. \square

(ii) \Rightarrow (iii): Immediate from the definitions.

(iii) \Rightarrow (i): For each $x \in A_*$, let A_x be the algebra dual to the E-subspace of A_* with the universe $\uparrow x$. From Esakia duality it follows that A is a subalgebra of the product of the family $\{A_x : x \in X\}$. Since varieties are closed under \mathbb{S} and \mathbb{P} , it suffices to prove that each A_x is a diamond Heyting algebra. To this end, take some $x \in A_*$. Since A_* is a diamond system, the dual $\uparrow x$ of A_* is a diamond sequence. Accordingly, it satisfies the three point rule, whence A_x is a cascade Heyting algebra by Theorem 3.7. Hence, A_x is locally finite by Theorem 3.3. It follows that A_x belongs to the variety generated by its finite subalgebras. Therefore, to conclude the proof, it suffices to show that all finite subalgebras of A_x are diamond Heyting algebras. Consider one such algebra B . The dual space B_* is a finite p-morphic image of the diamond sequence $\uparrow x$. Thus, B_* is a finite diamond sequence, whence B is a diamond Heyting algebra by definition. \square

Corollary 4.7. *DHA is axiomatized by $\mathcal{J}(P_1), \mathcal{J}(P_2), \mathcal{J}(P_3)$, and $\mathcal{J}(P_4)$.*

We close this section with the following corollary to Theorem 4.4 which will be used in Section 6.

Corollary 4.8. *Let X be an image-finite poset. If $\text{Up}(X)$ is a diamond algebra, then X is a diamond system.*

Proof. As $\text{Up}(X)$ is a diamond algebra, by Theorem 4.4 the poset underlying $\text{Up}(X)_*$ is a diamond system. It is well known that the map $\varepsilon: X \rightarrow \text{Up}(X)_*$, defined by

$$\varepsilon(x) = \{U \in \text{Up}(X) : x \in U\} \text{ for all } x \in X,$$

is a well-defined order embedding. Consequently, from the fact that $\text{Up}(X)_*$ is a diamond system it follows that X satisfies Conditions (D1) and (D2).

Suppose with a view to contradiction, that X does not satisfy Condition (D3). Then there are $x, y, z \in X$ such that $x \leq y, z$ and $\uparrow y \cap \uparrow z = \emptyset$. As X is image-finite, $\uparrow x$ is finite. Thus, we may assume without loss of generality that y and z are maximal and that x is maximal among the lower bounds of y and z . Then we can repeat the proof of Claim 4.5 and obtain the desired contradiction.

It only remains to prove that X satisfies Condition (D4). Suppose not. Then there are $\perp, x, y, z, v, \top \in X$ such that $\perp \leq x, y \leq z, v \leq \top$ and $\uparrow x \cap \uparrow y \cap \downarrow z \cap \downarrow v = \emptyset$. Since X is image-finite, the upset $\uparrow \perp$ is finite, whence we may assume without loss of generality that \perp is maximal among the lower bounds of x and y . Thus, repeating the proof of Claim 4.6 we obtain a contradiction, as desired. \square

5. FORBIDDEN POSETS

The aim of this section is to prove the following result.

Theorem 5.1. *Let \mathcal{V} be a variety of Heyting algebras. If the profinite members of \mathcal{V} are profinite completions, then $\mathcal{V} \subseteq \text{DHA}$.*

To this end, recall that a poset $\langle X, \leq \rangle$ is *Esakia representable* if there exists a topology τ on X such that $\langle X, \leq, \tau \rangle$ is an Esakia space. In order to establish Theorem 5.1, we require the following:

Proposition 5.2. *If $P \in \{P_1, P_2, P_3, P_4\}$, then there is a poset X which is a p-morphic image of a disjoint union of copies of P and is not Esakia representable.*

To derive Theorem 5.1 from Proposition 5.2 we recall the following well-known fact (the proof of which can essentially be found in [15, Thm. 5.47(ii) & 5.48]).

Lemma 5.3. *Let $\{X_i : i \in I\} \cup \{Y\}$ be a family of posets and \mathbb{V} a variety of Heyting algebras such that $\{\text{Up}(X_i) : i \in I\} \subseteq \mathbb{V}$. If Y is a p -morphic image of the disjoint union of the various X_i , then $\text{Up}(Y) \in \mathbb{V}$.*

Proof of Theorem 5.1. We reason by contraposition. Suppose \mathbb{V} is a variety of Heyting algebras such that $\mathbb{V} \not\subseteq \text{DHA}$. By Corollary 4.7, DHA is axiomatized by the Jankov formulas $\mathcal{J}(P_1), \mathcal{J}(P_2), \mathcal{J}(P_3)$, and $\mathcal{J}(P_4)$. Therefore, $\mathbb{V} \not\subseteq \text{DHA}$ implies

$$\mathbb{V} \cap \{\text{Up}(P_1), \text{Up}(P_2), \text{Up}(P_3), \text{Up}(P_4)\} \neq \emptyset.$$

Consequently, there is $P \in \{P_1, P_2, P_3, P_4\}$ such that $\text{Up}(P) \in \mathbb{V}$. By Proposition 5.2, there is a p -morphic image X of a disjoint union of copies of P that is not Esakia representable. We have $\text{Up}(X) \in \mathbb{V}$ by Lemma 5.3. Furthermore, as P has bounded depth and width, so does X . In particular, this guarantees that X is image-finite, and hence $\text{Up}(X)$ is a profinite member of \mathbb{V} by Theorem 2.20(i).

On the other hand, by Theorem 2.20(ii) and Corollary 2.16, the algebra $\text{Up}(X)$ is a profinite completion if and only if X is Esakia representable. As the latter is not the case, we conclude that $\text{Up}(X)$ is not a profinite completion. Thus, \mathbb{V} has a profinite member that is not a profinite completion. \square

The rest of this section is devoted to proving Proposition 5.2.

Definition 5.4. Let X be a poset and $D, E \subseteq X$. We say that (D, E) is a *surjective matching* if the following conditions hold:

- (M1) $D \cap E = \emptyset$;
- (M2) for every $x \in D$ there is $y \in E$ such that $y < x$;
- (M3) for every $y \in E$ there is $x \in D$ such that $y < x$;
- (M4) for every $x, z \in D$ and $y \in E$, if $y < x, z$ then $x = z$.

A surjective matching (D, E) is said to be a *bijective matching* if it satisfies the following additional conditions:

- (M5) for every $x, y \in D \cup E$, if $y < x$ then $x \in D$ and $y \in E$;
- (M6) for every $x \in D$ and $y, z \in E$, if $y, z < x$ then $y = z$.

In this case, $E \cup D$ becomes a disjoint union of two-element chains when endowed with the order inherited from X .

Lemma 5.5 (Matching Lemma). *Let X be an Esakia space and (D, E) a surjective matching in X . Let*

$$F = (\uparrow(D \cup E)) \setminus (D \cup E).$$

If E and F are compact, then so is D .

Proof. Suppose that E is compact and consider an open cover \mathcal{K} of D . For $U \in \mathcal{K}$, define

$$U' = \downarrow(U \setminus (E \cup F)) \text{ and } \mathcal{K}' = \{U' : U \in \mathcal{K}\}.$$

We will prove that \mathcal{K}' is an open cover of E .

Claim 5.6. *The following conditions hold:*

- (i) U' is open;
- (ii) $U \cap D = U' \cap D$;
- (iii) for every $x \in D$ and $y \in E$ with $y < x$ we have $x \in U'$ if and only if $y \in U'$.

Proof. (i): Since X is Hausdorff and E, F are compact, they are closed. Because U is open, this implies that so is $U \setminus (E \cup F)$. Lastly, since the downset of an open set is open in an Esakia space, we conclude that $U' = \downarrow(U \setminus (E \cup F))$ is open, as desired.

(ii): By Condition (M1) and the definition of F , the intersection $(E \cup F) \cap D$ is empty. This yields

$$U \cap D \subseteq U \setminus (E \cup F) \subseteq U'.$$

Therefore, it remains to show that $U' \cap D \subseteq U$.

For this we first show that

$$(\uparrow D) \setminus D \subseteq F. \quad (6)$$

Let $x \in (\uparrow D) \setminus D$. We show that $x \notin E$. From $x \in (\uparrow D) \setminus D$ it follows that there is $y \in D$ such that $y < x$. If $x \in E$, by Condition (M3) there is $z \in D$ such that $x < z$. Since $y \in D$, by Condition (M2), $v < y$ for some $v \in E$. Now, $v \in E$ and $v < y, z \in D$. Therefore, $y = z$ by Condition (M4), contradicting $y < x < z$. This establishes $x \notin E$. Consequently,

$$x \in (\uparrow D) \setminus (D \cup E) \subseteq (\uparrow(D \cup E)) \setminus (D \cup E) = F,$$

concluding the proof of (6).

Now suppose $x \in U' \cap D$. By (6),

$$x \in U' = \downarrow(U \setminus (E \cup F)) \subseteq \downarrow(U \setminus ((\uparrow D) \setminus D)).$$

Therefore, there is $y \in U \setminus ((\uparrow D) \setminus D)$ such that $x \leq y$. From $x \in D$ and $x \leq y$, it follows that $y \in \uparrow D$. Together with $y \in U \setminus ((\uparrow D) \setminus D)$, this implies $y \in D$. Finally, in view of $x \in D$, by Condition (M2) there is $z \in E$ such that $z < x$. Consequently, $z \leq x, y \in D$. By Condition (M4) we obtain $x = y$, whence $x = y \in U$. Hence, $U' \cap D \subseteq U$.

(iii): Suppose $x \in D$ and $y \in E$ with $y < x$. Since U' is a downset and $y < x$, if $x \in U'$, then $y \in U'$. To prove the converse, suppose that $y \in U'$. Then there is $z \in U \setminus (E \cup F)$ such that $y \leq z$. Notice that $z \in \uparrow E$ as $z \geq y \in E$. Together with $z \notin F$, this implies $z \in D \cup E$. Furthermore, since $z \notin E$, we obtain $z \in D$, whence $y \in E, x, z \in D$, and $y \leq x, z$. Thus, by Condition (M4), $x = z \in U'$. \square

Now, observe that \mathcal{K}' is a family of open sets by Condition (i) of the Claim. We prove that it is a cover of E . Let $y \in E$. By Condition (M3), there is $x \in D$ such that $y < x$. Since \mathcal{K} is a cover of D , by Condition (ii) of the Claim, the same holds for \mathcal{K}' . Thus, $x \in U'$ for some $U' \in \mathcal{K}'$. By Condition (iii) of the Claim, we conclude that $y \in U'$. Hence, \mathcal{K}' is an open cover of E .

Since E is compact, \mathcal{K}' has a finite subcover $\{U'_1, \dots, U'_n\}$. We prove that $\{U_1, \dots, U_n\}$ is a cover of D . Let $x \in D$. By Condition (M2), there is $y \in E$ such that $y < x$. Therefore, $y \in U'_i$ for some i . Furthermore, $x \in U'_i$ by Condition (iii) of the Claim. Thus, applying Condition (ii) of the Claim yields that $x \in U_i$. Consequently, D is compact. \square

We prove Proposition 5.2 by contradiction. Once we have a candidate $\langle X, \leq \rangle$ for a poset which is not Esakia representable, we suppose that there is a topology τ on X such that $\langle X, \leq, \tau \rangle$ is an Esakia space. Then we find a bijective matching (D, E) in X such that E is compact and D is not, a contradiction with the Matching Lemma.

For the compactness of E we use the fact that in Esakia spaces principal downsets are closed, and hence compact. In the part of the proof regarding P_1 and P_2 , the set E has the form $\downarrow x \cap \downarrow y$ for distinct elements x and y . When dealing with P_3 , we take E of the form $\downarrow x \setminus \{x\}$, where x is an isolated point. The case of P_4 will be reduced to that of P_3 . The non-compact set D will be constructed with the help of the following known result. We give a short proof for the sake of completeness.

Lemma 5.7. *Let X be a Hausdorff space and Y an infinite subset of it. There is $y \in Y$ such that for every finite subset Z of Y the set $Y \setminus (\{y\} \cup Z)$ is not compact in X .*

Proof. First suppose that Y is not compact. In this case, subtracting from Y any finite set gives a noncompact set. Thus, we can choose any $y \in Y$. Next suppose that Y is compact. Since Y is infinite, this implies that Y has a limit point y , i.e., an element $y \in Y$ such that every open neighbourhood U of y contains an element of $Y \setminus \{y\}$. To conclude the proof, it suffices to show that $Y \setminus \{y\}$ is not compact (as this implies that subtracting from $Y \setminus \{y\}$ any finite set produces a noncompact set). Since X is Hausdorff, for every $x \in Y \setminus \{y\}$, consider an open neighbourhood U_x of x and an open neighbourhood V_x of y such that $U_x \cap V_x = \emptyset$. Then $\{U_x : x \in Y \setminus \{y\}\}$ is an open cover of $Y \setminus \{y\}$. Since y is a limit point of Y , every intersection of finitely many sets of the form V_x contains a point in Y . Hence, $\{U_x : x \in Y \setminus \{y\}\}$ has no finite subcover of $Y \setminus \{y\}$, whence $Y \setminus \{y\}$ is not compact. \square

Lastly, we make use of the following observation.

Lemma 5.8. *Let X and Y be posets and $\min X$ the set of minimal elements of X . If $X = \uparrow \min X$ and $Y \cong \uparrow x$ for every $x \in \min X$, then X is a p-morphic image of a disjoint union of copies of Y .*

Proof. For each $x \in \min X$, let Y_x be a copy of Y and let $f_x: Y_x \rightarrow \uparrow x$ be an order-isomorphism. We let Z be the disjoint union of the various Y_x . Then the union of the maps f_x is a well-defined p-morphism $f: Z \rightarrow X$, which is onto since $X = \uparrow \min X$. Thus, X is a p-morphic image of a disjoint union of copies of Y . \square

Proof of Proposition 5.2. The proof is divided into four cases, corresponding to the posets P_1, P_2, P_3 , and P_4 .

The case of P_1 . We construct a p-morphic image X of a disjoint union of copies of P_1 that is not Esakia representable. For every three-element set $\{k, m, n\}$ of natural numbers, consider a new element $\perp_{\{k,m,n\}}$. Also, consider another element \top and define

$$X = \mathbb{N} \cup \{\top\} \cup \{\perp_{\{k,m,n\}} : k, m, n \text{ are distinct natural numbers}\};$$

for every $x, y \in X$,

$$\begin{aligned} x \leq y &\iff \text{either } x = y \text{ or } y = \top \\ &\text{or } (x = \perp_{\{n,m,k\}} \text{ and } y \in \{n, m, k\} \text{ for some } n, m, k \in \mathbb{N}). \end{aligned}$$

By Lemma 5.8, X is a p-morphic image of a disjoint union of copies of P_1 . Therefore, it only remains to prove that X is not Esakia representable. Suppose the contrary, with a view to contradiction. Then there is an Esakia space $\langle X, \leq, \tau \rangle$. Choose a natural number k and define $Y = \mathbb{N} \setminus \{k\}$. By Lemma 5.7, there is $m \in \mathbb{N} \setminus \{k\}$ such that the set

$$D := \mathbb{N} \setminus \{k, m\}$$

is not compact. On the other hand, the set

$$E := \{\perp_{\{k,m,n\}} : n \in D\} = \downarrow k \cap \downarrow m$$

is the intersection of two principal downsets, so it is closed, and hence compact.




FIGURE 8. The bijective matching (D, E) in the poset X , constructed in the case of P_1 .

Furthermore, the set

$$F := (\uparrow(D \cup E)) \setminus (D \cup E) = \{m, k, \top\}$$

is finite. Consequently, as X is a Hausdorff, F is closed. In order to obtain a contradiction with the Matching Lemma, it only remains to prove that (D, E) is a surjective matching in X . But this follows immediately from the construction of X and (D, E) , as illustrated in Figure 8.

The case of P_2 . We construct a p-morphic image X of a disjoint union of copies of P_2 that is not Esakia representable. Let $\mathbb{N}' = \{1', 2', 3', \dots\}$ be a disjoint copy of the set \mathbb{N} of natural numbers. For every pair of distinct natural numbers n and k , take a new element $\perp_{n,k}$. Consider two new elements \top and σ and define

$$X = \mathbb{N} \cup \mathbb{N}' \cup \{\top, \sigma\} \cup \{\perp_{n,k} : n \text{ and } k \text{ are distinct natural numbers}\};$$

for every $x, y \in X$,

$$\begin{aligned} x \leq y &\iff \text{either } x = y \text{ or } y = \top \\ &\text{or } (y = \sigma \text{ and } x \in X \setminus (\mathbb{N}' \cup \{\top\})) \\ &\text{or } (x = \perp_{n,k} \text{ and } y \in \{n, k'\} \text{ for some } n, k \in \mathbb{N}). \end{aligned}$$

By Lemma 5.8, X is a p-morphic image of a disjoint union of copies of P_2 .

Suppose, with a view to contradiction, that X is Esakia representable. Then there is an Esakia space $\langle X, \leq, \tau \rangle$. By Lemma 5.7, there exists a natural number k such that $\mathbb{N} \setminus \{k\}$ is not compact in X . Define

$$D = \mathbb{N} \setminus \{k\} \text{ and } E = \{\perp_{n,k} : n \in D\} = \downarrow k' \cap \downarrow \sigma.$$

Then E is closed, hence compact. Moreover, as illustrated in Figure 9, the set

$$F := (\uparrow(D \cup E)) \setminus (D \cup E) = \{\top, \sigma, k'\}$$

is finite. Consequently, as X is Hausdorff, F is closed. In order to obtain a contradiction with Matching Lemma, it only remains to prove that (D, E) is a surjective matching in X . But this is a direct consequence of the construction of X and (D, E) , as shown in Figure 9.




FIGURE 9. The bijective matching (D, E) in the poset X , constructed in the case of P_2 .

The case of P_3 . We follow the same pattern as in the previous two cases. First, we construct a poset X which is a p-morphic image of a disjoint union of copies of P_3 . For every pair of integers n and k such that $n + 1 < k$, consider a new element $\perp_{\{n,k\}}$ and define

$$X = \mathbb{Z} \cup \{\perp_{\{n,k\}} : n, k \in \mathbb{Z} \text{ and } n + 1 < k\};$$

for every $x, y \in X$,

$$x \leq y \iff \text{either } x = y \text{ or } (x = \perp_{\{n,k\}} \text{ and } y \in \{n, k\} \text{ for some } n, k \in \mathbb{Z}).$$

By Lemma 5.8, X is a p-morphic image of a disjoint union of copies of P_3 .

It only remains to show that X is not Esakia representable. Suppose the contrary, with a view to contradiction. Then there exists an Esakia space $\langle X, \leq, \tau \rangle$. As in in the previous cases, we need to find a surjective matching (D, E) in X such that E is compact, but D is not. To this end, we rely on the following:

Theorem 5.9 ([62, Thm. 8.5.4]). *Every countable compact Hausdorff space has an isolated point.*

We shall construct a bijective matching in X with the aid of the following:

Claim 5.10. *There exists an integer k that is isolated in X and such that $\mathbb{Z} \setminus \{k-1, k, k+1\}$ is not compact in X .*

Proof. Notice that \mathbb{Z} is the set of maximal elements of X . Therefore, by Proposition 2.13, \mathbb{Z} is closed in X . This means that the subspace \mathbb{Z} of X is a Stone space. Thus, we can apply Theorem 5.9, obtaining that \mathbb{Z} has an isolated point x , i.e., a point x such that $\{x\}$ is open in \mathbb{Z} . Since \mathbb{Z} is compact and infinite, it must have a limit point. Hence, \mathbb{Z} is the union of the two disjoint nonempty sets, respectively, of isolated points in \mathbb{Z} and of limit points in \mathbb{Z} . As a consequence, there exist two consecutive integers such that one of them is a limit point in \mathbb{Z} and the other is an isolated point in \mathbb{Z} (otherwise either the set of isolated or limit points in \mathbb{Z} would be empty).

Let k be the isolated point in this pair. Then either $k-1$ or $k+1$ is a limit point in \mathbb{Z} , whence the set $\mathbb{Z} \setminus \{k-1, k, k+1\}$ is not compact in \mathbb{Z} , and hence in X . To conclude the proof, it only remains to show that k is isolated in X too. Since k is isolated in \mathbb{Z} , there exists an open set U of X such that $U \cap \mathbb{Z} = \{k\}$. Let

$$V = X \setminus \downarrow(X \setminus U) = \{x \in X : \uparrow x \subseteq U\}.$$

As downsets of closed sets are closed in Esakia spaces, V is open in X . Furthermore, $V \setminus \mathbb{Z} = \emptyset$ as every element in $X \setminus \mathbb{Z}$ has exactly two successors in \mathbb{Z} and one of them should be different from k . Together with the fact that $U \cap \mathbb{Z} = \{k\}$, this yields that $V = \{k\}$. Consequently, k is isolated in X . \square

Let k be as in Claim 5.10 and define

$$D = \mathbb{Z} \setminus \{k-1, k, k+1\} \text{ and } E = (\downarrow k) \setminus \{k\}.$$

Then D is not compact and E is closed, hence compact. From the definition of the order relation on X it follows that

$$F := (\uparrow(D \cup E)) \setminus (D \cup E) = \{k\}$$

(see Figure 10). Thus, F is finite, and hence closed. As (D, E) is a bijective matching in X (see Figure 10), this contradicts the Matching Lemma.




FIGURE 10. The bijective matching (D, E) in the poset X , constructed in the case of P_3 .


The case of P_4 . Let X be the poset defined in the case of P_3 . We define a new poset Y in which X is a downset. Consider three new elements σ, τ , and \top and define

$$Y = X \cup \{\sigma, \tau, \top\};$$

for every $x, y \in Y$,

$$\begin{aligned} x \leq y \iff & \text{either } x = y \text{ or } y = \top \text{ or } (x \in X \text{ and } y \in \{\sigma, \tau\}) \\ & \text{or } (x = \perp_{\{n,k\}} \text{ and } y \in \{n, k\} \text{ for some } n, k \in \mathbb{N}). \end{aligned}$$

A portion of X is depicted in Figure 11.

FIGURE 11. A fragment of the poset Y , constructed in the case of P_4 .

By Lemma 5.8, X is a p-morphic image of a disjoint union of copies of P_4 . Therefore, to conclude the proof, it suffices to show that Y is not Esakia representable. As before, suppose with a view to contradiction, that there is an Esakia space $\langle Y, \leq, \tau \rangle$. We prove that the ordered subspace X of Y is an Esakia space. For this it suffices to show that X is a clopen of Y as, in this case, the conditions in the definition of an Esakia space follow [28, Thm. 3.2.6]. First notice that X is open in Y since Y is Hausdorff and $Y \setminus X$ is finite. Moreover, X is closed in Y since $X = \downarrow\sigma \cap \downarrow\tau$. Consequently, X is clopen in Y and, hence, X is an Esakia space. But this contradicts the fact that X is not Esakia representable, which was established in the case of P_3 . \square

6. TOPOLOGIES FOR DIAMOND SYSTEMS

The aim of this section is to establish our main result:

Theorem 6.1. *Let \mathcal{V} be a variety of Heyting algebras. The profinite members of \mathcal{V} are profinite completions if and only if \mathcal{V} is a subvariety of DHA.*

Notice that Theorem 5.1 provides one half of the above result. To prove the other half, we require the following:

Proposition 6.2. *Let X be an image-finite diamond system. Then there exists an Esakia space X^+ such that its underlying poset is also a diamond system and X is the image-finite part of X^+ .*

Proposition 6.2 implies Theorem 6.1 as follows.

Proof of Theorem 6.1. The “only if” part is Theorem 5.1. To prove the “if” part, suppose that $\mathcal{V} \subseteq \text{DHA}$ and consider a profinite algebra $A \in \mathcal{V}$. By Theorem 2.20(i), there is an image-finite poset X such that $A \cong \text{Up}(X)$. In particular, $\text{Up}(X)$ is a diamond Heyting algebra, whence X is a diamond system by Corollary 4.8. In view of Proposition 6.2, X is the image-finite part of some Esakia space. Consequently, $\text{Up}(X)$ is a profinite completion by Theorem 2.20(ii). Since $A \cong \text{Up}(X)$, we conclude that A is a profinite completion. \square

The rest of this section is devoted to proving Proposition 6.2.

Proof of Proposition 6.2. Recall from Definition 4.1 that diamond sequences are downward directed diamond systems. Let \mathcal{A}_∞ be the set of all infinite diamond sequences in X that are maximal with respect to inclusion. With every $A \in \mathcal{A}_\infty$ we associate a new element \perp_A (so that $A \neq B$ implies $\perp_A \neq \perp_B$). Let

$$X^+ = X \cup \{\perp_A : A \in \mathcal{A}_\infty\}$$

and extend the order on X to that on X^+ by setting for $x, y \in X^+$,

$$x \leq y \iff \text{either } x = y \text{ or } x \leq y \text{ in } X \\ \text{or } x = \perp_A \text{ and } y \in A \text{ for some } A \in \mathcal{A}_\infty.$$

The resulting poset X^+ is then obtained by adding a new lower bound \perp_A for each maximal diamond sequence A . It follows that $\uparrow\perp_A = \{\perp_A\} \cup A$. Notice that X^+ is a diamond system and X is the image-finite part of X^+ . In particular, if X has finite depth, then $X = X^+$.

To complete the proof, it suffices to show that X^+ can be endowed with a topology τ such that $\langle X^+, \tau, \leq \rangle$ is an Esakia space. Notice that such a topology is not unique. For example, let Y be an infinite poset of depth two with a maximum. We can turn Y into an Esakia space by making any of its elements the unique limit point of the space. This results in different Esakia spaces with the same underlying poset Y .

To define τ on X^+ , choose a maximal element \top of X (equiv. of X^+). Also, for every pair x, y of incomparable elements in X such that $\downarrow x \cap \downarrow y$ is nonempty, choose an immediate predecessor $\perp_{\{x,y\}}$ of x and y in X . The existence of $\perp_{\{x,y\}}$ follows from the fact that X is image-finite and Proposition 4.2.

We define a topology τ on X^+ as follows. A set $U \subseteq X^+$ is open in τ if and only if it satisfies the following conditions:

- (T1) If $\top \in U$, then there is a finite set $Z \subseteq X$ such that $X^+ \setminus \downarrow Z \subseteq U$;
- (T2) For every $x \in X$, if $x \in U$, then there is a finite set $Z \subseteq X \setminus \uparrow x$ such that $\downarrow x \setminus \downarrow Z \subseteq U$;
- (T3) For every pair of incomparable elements x, y of X such that $\downarrow x \cap \downarrow y \neq \emptyset$, if $\perp_{\{x,y\}} \in U$, then there is a finite set $Z \subseteq \downarrow x \cap \downarrow y$ such that $(\downarrow x \cap \downarrow y) \setminus \downarrow Z \subseteq U$;
- (T4) For every $A \in \mathcal{A}_\infty$, if $\perp_A \in U$, then there is $x \in A$ such that $\downarrow x \subseteq U$.

Remark 6.3. The following provides an intuitive meaning of τ . Postulate (T1) asserts that if \top belongs to an open set U , then *almost all* the space belongs to U . Similarly, (T2) should be read as stating that if an element $x \in X$ belongs to U , then *almost all* the downset $\downarrow x$ belongs to U . The remaining postulates can be explained along the same lines.

Claim 6.4. *The set τ is a topology on X^+ .*

Proof. That $\emptyset, X^+ \in \tau$ and that τ is closed under arbitrary unions follow from the definition. Thus, it is enough to consider two sets $U_1, U_2 \in \tau$ and verify that their intersection $U_1 \cap U_2$ satisfies (T1)–(T4).

First, suppose that $\top \in U_1 \cap U_2$. Then there are finite $Z_1, Z_2 \subseteq X$ such that $X^+ \setminus \downarrow Z_1 \subseteq U_1$ and $X^+ \setminus \downarrow Z_2 \subseteq U_2$. Therefore, $X^+ \setminus \downarrow(Z_1 \cup Z_2) \subseteq U_1 \cap U_2$. This shows that $U_1 \cap U_2$ satisfies (T1). A similar argument shows that $U_1 \cap U_2$ satisfies (T2) and (T3).

To show that $U_1 \cap U_2$ satisfies (T4), suppose that $\perp_A \in U_1 \cap U_2$ for some $A \in \mathcal{A}_\infty$. Then there are $x_1, x_2 \in A$ such that $\downarrow x_1 \subseteq U_1$ and $\downarrow x_2 \subseteq U_2$. Since A is a diamond sequence, there exists $x \in A$ such that $x \leq x_1, x_2$. Thus, $\downarrow x \subseteq U_1 \cap U_2$, as desired. \square

To prove that the ordered topological space $X^+ := \langle X^+, \leq, \tau \rangle$ is an Esakia space, we show three facts: that downsets of open sets are open in X^+ , that the space X^+ is compact, and that it satisfies the Priestley separation axiom. The verification of each of these is independent from the others.

Claim 6.5. *Let U be an open set in X^+ . Then its downset $\downarrow U$ is also open in X^+ .*

Proof. We need to prove that $\downarrow U$ satisfies (T1)–(T4). To prove (T1), suppose that $\top \in \downarrow U$. Since \top is maximal, this yields $\top \in U$. As U satisfies (T1), there is a finite set Z such that $X \setminus \downarrow Z \subseteq U$, whence also $X \setminus \downarrow Z \subseteq \downarrow U$.

Condition (T2) holds for every downset, and in particular for $\downarrow U$. To prove (T3), consider a pair of incomparable elements $x, y \in X$ such that $\downarrow x \cap \downarrow y \neq \emptyset$. Suppose that $\perp_{\{x,y\}} \in \downarrow U$. If $\perp_{\{x,y\}} \in U$, then the fact that $\downarrow U$ satisfies (T3) follows from the assumption that U satisfies (T3) and the inclusion $U \subseteq \downarrow U$. On the other hand, if $\perp_{\{x,y\}} \notin U$, we have $\perp_{\{x,y\}} < u$ for some $u \in U$. Since x and y are immediate successors of $\perp_{\{x,y\}}$, Proposition 4.2 implies that $x \leq u$ or $y \leq u$. Thus, $\downarrow x \cap \downarrow y \subseteq \downarrow u \subseteq \downarrow U$.

It only remains to prove (T4). Suppose that $\perp_A \in \downarrow U$ for some $A \in \mathcal{A}_\infty$. If $\perp_A \in U$, then the fact that $\downarrow U$ satisfies (T4) follows from the assumption that U satisfies (T4) and the inclusion $U \subseteq \downarrow U$. On the other hand, if $\perp_A \notin U$, there is $x \in U$ such that $\perp_A < x$. By the definition of the order on X^+ , we get $x \in A$. Therefore, $\downarrow x \subseteq \downarrow U$. \square

We next turn to proving that X^+ is a compact space. Consider an open cover \mathcal{K} of X^+ . Our aim is to find a finite subcover of \mathcal{K} . The proof consists of a series of technical claims.

Let U_\top be a member of \mathcal{K} containing \top . By (T1), there exists a finite subset Z of X such that

$$X^+ \setminus U_\top \subseteq \downarrow Z. \quad (7)$$

For every $z \in Z$, let z° be the unique maximal element in X such that $z \leq z^\circ$. The existence and uniqueness of z° is guaranteed by the fact that X is image-finite and Proposition 4.2. Define

$$Y_1 = \{z^\circ : z \in Z\}, \quad Y^+ = \downarrow Y_1, \quad \text{and} \quad Y = Y^+ \cap X, \quad (8)$$

where $\downarrow Y_1$ is computed in X^+ .

Claim 6.6. *For every $x \in X$ there is a finite subset \mathcal{K}_x of \mathcal{K} and a finite subset P_x of immediate predecessors of x such that*

$$\downarrow x \setminus \downarrow P_x \subseteq \bigcup \mathcal{K}_x.$$

Proof. There is $U_x \in \mathcal{K}$ such that $x \in U_x$. By (T2), there is a finite subset Z_x of $X \setminus \uparrow x$ such that $\downarrow x \setminus \downarrow Z_x \subseteq U_x$. Clearly we may assume that $\downarrow x \cap \downarrow z \neq \emptyset$ for every $z \in Z_x$. Define

$$Z_1^x = \{z \in Z : z < x\} \quad \text{and} \quad Z_2^x = \{z \in Z : x \text{ and } z \text{ are incomparable}\}.$$

Then Z_x is the union of the disjoint sets Z_1^x and Z_2^x . Since X is image-finite, we may assume without loss of generality that Z_1^x is a set of immediate predecessors of x .

Consider $z \in Z_2^x$. Since we assumed that $\downarrow x \cap \downarrow z \neq \emptyset$, there is an element in $\downarrow x \cap \downarrow z$ of the form $\perp_{\{x,z\}}$. Let $U_{\perp_{\{x,z\}}}$ be a member of \mathcal{K} containing $\perp_{\{x,z\}}$. By (T3), there exists a finite subset $Q_{x,z}$ of $\downarrow x \cap \downarrow z$ such that $(\downarrow x \cap \downarrow z) \setminus \downarrow Q_{x,z} \subseteq U_{\perp_{\{x,z\}}}$. Since X is image-finite, we may assume without loss of generality that $Q_{x,z}$ is a set of immediate predecessors of x .

Define

$$\mathcal{K}_x = \{U_x\} \cup \{U_{\perp_{\{x,z\}}} : z \in Z_2^x\} \quad \text{and} \quad P_x = Z_1^x \cup \bigcup \{Q_{x,z} : z \in Z_2^x\}. \quad (9)$$

Notice that \mathcal{K}_x is a finite subset of \mathcal{K} and P_x is a finite subset of the immediate predecessors of x . Therefore, it only remains to prove that $\downarrow x \setminus \downarrow P_x \subseteq \bigcup \mathcal{K}_x$, i.e., that $\downarrow x \setminus \bigcup \mathcal{K}_x \subseteq \downarrow P_x$. To this end, consider $y \in \downarrow x \setminus \bigcup \mathcal{K}_x$. Since $y \notin U_x$ and $y \leq x$, we have $y \in \downarrow Z_x = \downarrow(Z_1^x \cup Z_2^x)$. If $y \in \downarrow Z_1^x$, then $y \in \downarrow P_x$ as desired. So suppose that $y \leq z$ for some $z \in Z_2^x$. Then $y \in \downarrow x \cap \downarrow z$. Together with $y \notin U_{\perp_{\{x,z\}}}$ this yields $y \in \downarrow Q_{x,z} \subseteq \downarrow P_x$. \square

We define recursively a sequence of subsets S_n of Y as follows:

$$S_1 = Y_1 \quad \text{and} \quad S_{k+1} = \bigcup \{P_y : y \in S_k\} \quad \text{for every integer } k \geq 1,$$

where Y_1 is defined in (8) and P_y in (9). Then each S_k is a set of elements of depth k in X . For every positive integer k , set

$$S_{\leq k} = S_1 \cup \dots \cup S_k. \quad (10)$$

By Claim 6.6, the sets S_1, \dots, S_k are finite, whence we obtain:

Claim 6.7. *For every positive integer k , the set $S_{\leq k}$ is finite.*

Define

$$S = \bigcup \{S_k : k \text{ is a positive integer}\}, \quad (11)$$

$$S^+ = S \cup \{\perp_A : A \in \mathcal{A}_\infty \text{ and } A \cap S \text{ is infinite}\}.$$

For every $A \in \mathcal{A}_\infty$ such that the set $A \cap S$ is infinite, choose an element $U_{\perp_A} \in \mathcal{K}$ containing \perp_A . By (T4), there exists $x \in A$ such that $\downarrow x \subseteq U_{\perp_A}$. As $\langle A, \leq \rangle$ is an image-finite diamond sequence and $x \in A$, by Proposition 4.2 the set $A \setminus \downarrow x$ is finite. Since $A \cap S$ is infinite, this implies that $A \cap S \cap \downarrow x \neq \emptyset$. Therefore, there is $x_A \in S \cap A$ such that $x_A \leq x$. Thus,

$$\downarrow x_A \subseteq U_{\perp_A}. \quad (12)$$

Bearing this in mind, set

$$B = \{x_A : A \in \mathcal{A}_\infty \text{ and } A \cap S \text{ is infinite}\} \text{ and } S^- = S \setminus \downarrow B. \quad (13)$$

We show that S^- is finite. For this we use the classical Brouwer's fan theorem; see for instance [24, Thm. 3.3.20]. We recall that a poset is a *fan* if it is rooted, its principal downsets are finite chains, and every element has only finitely many immediate successors.

Theorem 6.8 (Brouwer's Fan Theorem). *If all chains in a fan F are finite, then their length is bounded by some positive integer.*

We turn $\langle S^-, \leq \rangle$ into a tree by means of a variant of the standard technique in modal logic known as *unravelling*; see for instance [18, Thm. 2.19]. Consider the following set of finite sequences

$$T = \{\langle x_1, x_2, \dots, x_n \rangle : n \text{ is a nonnegative integer, } x_1 \in S_1 \cap S^-, \dots, x_n \in S_n \cap S^-, \\ \text{and } x_n < \dots < x_2 < x_1\}.$$

We assume that the empty sequence, denoted by $\langle \rangle$, belongs to T .²

Notice that since the elements of each S_k have depth exactly k , the assumptions that $x_n < \dots < x_2 < x_1$ and $x_1 \in S_1, \dots, x_n \in S_n$ imply that x_i is an immediate successor of x_{i+1} for every $i < n$. We define an order relation \sqsubseteq on T by setting for every $s, t \in T$ that

$$s \sqsubseteq t \iff s \text{ is an initial segment of } t.$$

Claim 6.9. *The poset $T := \langle T, \sqsubseteq \rangle$ is a fan.*

Proof. It is well known that T is a tree with root $\langle \rangle$ (see, e.g., [18, Thm. 2.19]). To see that every element of T has only finitely many immediate successors, consider a sequence $\langle x_1, \dots, x_n \rangle \in T$. The immediate successors of $\langle x_1, \dots, x_n \rangle$ have the form $\langle x_1, \dots, x_n, y \rangle$ where $y \in S_{n+1}$ and $y < x_n$. Since the set S_{n+1} is finite, we conclude that $\langle x_1, \dots, x_n \rangle$ has only finitely many immediate successors. \square

Claim 6.10. *The set S^- is finite.*

Proof. Since $|S^-| \leq |T|$, it is enough to show that T is finite. In view of Brouwer's Fan Theorem and Claim 6.9, it suffices to prove that T does not have infinite chains.

Suppose, with a view to contradiction, that there is an infinite chain C in T . Define

$$D = \{x \in S^- : \text{there exists } \langle x_1, \dots, x_n \rangle \in C \text{ such that } x = x_n\}.$$

Observe that C is a set of finite sequences without repetitions of elements of D . Since C is infinite, so is D . We show that D is a chain in $\langle S^-, \leq \rangle$. Let $y, z \in D$. Then there are sequences $s = (s_1, \dots, s_n)$ and $t = (t_1, \dots, t_k)$ in C such that $y = s_n$ and $z = t_k$. Since C is a chain, either $s \sqsubseteq t$ or $t \sqsubseteq s$ in T . By symmetry we may assume that $s \sqsubseteq t$. Then $n \leq k$ and $s = (t_1, \dots, t_n)$. Consequently, $y = s_n = t_n \geq \dots \geq t_k = z$, as desired. Thus, D is an infinite chain in $\langle S^-, \leq \rangle$.

Define

$$A = \bigcup \{\uparrow y : y \in D\},$$

where the upsets are computed in X . We show that A is an infinite maximal diamond sequence in X . First, A is infinite since $D \subseteq A$. Because A is a subset of X which is a diamond system, it is a diamond system itself. Moreover, A is downward directed since D is a chain. Hence, A is a diamond sequence in X .

To prove that A is maximal, consider a diamond sequence A' in X such that $A \subseteq A'$. Let $x \in A'$. Suppose the depth of x is n and let y be an element of A of depth $n+1$ (its existence follows from the fact that D is infinite). Since $x, y \in A'$ and A' is downward directed, there is $z \in A'$ such that $z \leq x, y$. Thus, there is $v \in X$ of depth $n+1$ such that $z \leq v \leq x$. If $v = y$, then $y \leq x$, whence $x \in A$ (as $y \in A$ and A is an upset of X). Otherwise, having the same depth, v and y are incomparable. Therefore, applying the three point rule to $z \leq v, y, x$ and $v \leq x$, we get $y \leq x$. Thus, $x \in A$, whence $A' \subseteq A$. We

²Notice that our definition differs from the standard one in that our order is $x_n < \dots < x_1$ as opposed to the standard order $x_1 < \dots < x_n$.

conclude that A is an infinite maximal diamond sequence in X , i.e., $A \in \mathcal{A}_\infty$. In particular, this implies that X^+ contains an element of the form \perp_A .

Now, since $D \subseteq A$ is infinite and $D \subseteq S$, also $A \cap S$ is infinite. Then the set B , defined in (13), contains an element of the form x_A . By the definition of A , there exists $y \in D$ such that $y \leq x_A$. Therefore, $y \in \downarrow B$. But this contradicts the fact that $D \subseteq S^- = S \setminus \downarrow B$. \square

Let B' be the set of maximal elements of the set $\langle B, \leq \rangle$, defined in (13). Since X is image-finite, $\downarrow B' = \downarrow B$.

Claim 6.11. *The set B' is finite.*

Proof. Recall from (13) that $B \subseteq S$ and $S^- = S \setminus \downarrow B$. We show that every element of B' is either in S_1 or has an immediate successor in S^- . Let $x \in B' \setminus S_1$. Since $B' \subseteq S$, we get $x \in S_k$ for some integer $k > 1$. By definition of S_k , there is an immediate successor $y \in S_{k-1}$ of x . Notice that x is maximal in B and $x < y$. Consequently, $y \notin \downarrow B$, whence $y \in S_{k-1} \setminus \downarrow B \subseteq S^-$. Thus, x has an immediate successor in S^- .

By Claim 6.10, there is a positive integer k such that $S^- \subseteq S_{\leq k}$. Recall that S_n is a set of elements of depth n in X , for every positive integer n . Therefore, the set of elements of S with an immediate successor in $S^- \subseteq S_{\leq k}$ is contained in $S_{\leq k+1}$. As every element of $B' \subseteq S$ is either in S_1 or has an immediate successor in S^- , we conclude that $B' \subseteq S_{\leq k+1}$. By Claim 6.7, the set $S_{\leq k+1}$ is finite. Thus, so is B' . \square

Claim 6.12. *The space X^+ is compact.*

Proof. Recalling that \mathcal{K} is an open cover of X^+ , define

$$\mathcal{K}' = \{U_\top\} \cup \bigcup \{ \mathcal{K}_x : x \in S^- \} \cup \{U_{\perp_A} : x_A \in B'\},$$

where U_\top is defined before Claim 6.6, each \mathcal{K}_x is defined in (9), and each U_{\perp_A} is defined in the paragraph preceding (12). It follows from the definition that $\mathcal{K}' \subseteq \mathcal{K}$. By Claims 6.6, 6.10 and 6.11, the set \mathcal{K}' is finite. Thus, it only remains to prove that \mathcal{K}' covers X^+ . Let $x \in X^+$. We have two cases: either $x \in S^+$ or $x \notin S^+$.

First suppose that $x \in S^+$. If $x \in S^-$, then $x \in U_x$ by the definition of \mathcal{K}_x in (9). Thus, $x \in U_x \in \mathcal{K}_x \subseteq \mathcal{K}'$. Next suppose that $x \in S^+ \setminus S^-$. Then

$$x \in S^+ \setminus S^- \subseteq \{ \perp_A : A \in \mathcal{A}_\infty \text{ and } A \cap S \text{ is infinite} \} \cup \downarrow B \subseteq \downarrow B = \downarrow B'. \quad (14)$$

The above inclusions are justified as follows. The first one follows from the definitions of S^+ and S^- . The second one from the fact that for every $A \in \mathcal{A}_\infty$ such that $A \cap S$ is infinite, the element $x_A \in B$ (defined in the paragraph before (13)) belongs to A , whence $\perp_A \leq x_A \in B$. The last one is justified right before Claim 6.11. Finally, by (14) there is $x_A \in B'$ such that $x \leq x_A$. Together with the definition of U_{\perp_A} , this implies that $x \in \downarrow x_A \subseteq U_{\perp_A} \in \mathcal{K}'$.

Next we consider the case where $x \notin S^+$. We have three subcases:

- (i) $\uparrow x \cap S = \emptyset$;
- (ii) $\uparrow x \cap S$ is infinite;
- (iii) $\uparrow x \cap S$ is nonempty and finite.

(i): Since $Y_1 = S_1 \subseteq S$, we get $x \notin \downarrow Y_1$. As Y_1 is the set of maximal elements of $\downarrow Z$ and X is image-finite, this implies that $x \notin \downarrow Z$. Thus, by (7), $x \in U_\top \in \mathcal{K}'$.

(ii): Since X is image-finite, necessarily $x \in X^+ \setminus X$. Thus, there is $A \in \mathcal{A}_\infty$ such that $x = \perp_A$. As we pointed out after the definition of X^+ , we have $\uparrow x = \{ \perp_A \} \cup A$. As $\perp_A \notin S$ (by definition of S), this implies that $A \cap S = \uparrow x \cap S$. Therefore, by assumption, $A \cap S$ is infinite, and hence $x = \perp_A \in S^+$, a contradiction with the assumption that $x \notin S^+$.

(iii): Let y be a minimal element in $\uparrow x \cap S$. Since $x \notin S^+$ and $S \subseteq S^+$, we have $x \notin S$, whence $x < y$. As $y \in S$, either $y \in \downarrow B$ or $y \in S \setminus \downarrow B = S^-$. If $y \in \downarrow B$, then there is $x_A \in B'$ such that $y \leq x_A$ (since $\downarrow B = \downarrow B'$). By (12) and $x \leq y \leq x_A$, we obtain $x \in U_{\perp_A} \in \mathcal{K}'$, as desired. Next suppose that $y \in S^-$. Suppose, with a view to contradiction, that $P_y \cap \uparrow x \neq \emptyset$, where P_y is defined in (9). Then there is $x \in P_y$ with $x \leq z$. By Claim 6.6, z is an immediate predecessor of y . Therefore, $x \leq z < y$. Moreover, from

$y \in S$ it follows that there is a positive integer k such that $y \in S_k$. By definition of S_{k+1} and $z \in P_y$, we have $z \in S_{k+1} \subseteq S$. Together with $x \leq z < y$, this contradicts the minimality of y in $\uparrow x \cap S$. Thus, $P_y \cap \uparrow x = \emptyset$, i.e., $x \notin \downarrow P_y$. Consequently, $x \in \downarrow y \setminus \downarrow P_y$. By Claim 6.6, $x \in \bigcup \mathcal{K}_y$. Since $y \in S^-$, we have $\bigcup \mathcal{K}_y \subseteq \bigcup \mathcal{K}'$. We conclude that $x \in \bigcup \mathcal{K}'$. \square

The final step consists in proving that the Priestley separation axiom holds in X^+ .

Definition 6.13. We call *distinguished* the elements of X^+ of the form \top , $\perp_{\{x,y\}}$, and \perp_A . The remaining elements of X^+ will be called *nondistinguished*.

Claim 6.14. *The following conditions hold for every $u, v \in X^+$:*

- (i) *The set $\downarrow u \setminus \{u\}$ is open in X^+ .*
- (ii) *If v is nondistinguished, the set $\downarrow v$ is open in X^+ .*

Proof. Set $U = \downarrow u \setminus \{u\}$ and $V = \downarrow v$. We verify that both U and V are open at once. Since \top is maximal and distinguished, it does not belong to $U \cup V$. Thus, (T1) holds trivially for U and V . Moreover, since U and V are downsets, Condition (T2) also holds trivially.

For (T3), suppose that an element of the form $\perp_{\{x,y\}}$ belongs to U , i.e., $\perp_{\{x,y\}} < u$. Since X^+ is a diamond system, it has width at most two, whence x and y are the only immediate successors of $\perp_{\{x,y\}}$ in X^+ . As X is image-finite, $\uparrow \perp_{\{x,y\}} \subseteq X$, and $\perp_{\{x,y\}} < u$, either $x \leq u$ or $y \leq u$. In both cases, $\downarrow x \cap \downarrow y \subseteq \downarrow u \setminus \{u\} = U$, as desired. This arguments can be adapted to the case of V . For suppose that $\perp_{\{x,y\}} \in V = \downarrow v$. As v is nondistinguished, this implies $\perp_{\{x,y\}} < v$. Thus, replicating the proof described for U , we obtain that (T3) holds for V as well.

Finally, let $A \in \mathcal{A}_\infty$ and $\perp_A \in U$. Then $\perp_A < u$. By definition of the order on X^+ , we have that $u \in A$. As A is infinite and $\uparrow u$ is finite, there is $x \in A$ such that $u \not\leq x$. Since A is downward directed (being a diamond sequence), we may assume without loss of generality that $x < u$. Then $\downarrow x \subseteq \downarrow u \setminus \{u\} = U$, whence U satisfies (T4). Again, this arguments can be adapted to the case of V as follows. Suppose that $\perp_A \in V = \downarrow v$. Since v is nondistinguished, this implies that $\perp_A < v$, so we can repeat the argument for U . \square

Claim 6.15. *Let $x, y, u \in X$ be such that x and y are incomparable and $\downarrow x \cap \downarrow y \neq \emptyset$. If $\perp_{\{x,y\}} \not\leq u$, then either $\downarrow x \cap \downarrow y \cap \downarrow u = \emptyset$ or $u \in \downarrow x \cap \downarrow y$.*

Proof. Recall that x and y are the only immediate successors of $\perp_{\{x,y\}}$ in X and hence in X^+ , and that $\uparrow \perp_{\{x,y\}} \subseteq X$. Consequently, from $\perp_{\{x,y\}} \not\leq u$ it follows that $x, y \not\leq u$. Suppose there is $z \in \downarrow x \cap \downarrow y \cap \downarrow u$. Since X^+ has width at most two and x, y are incomparable, we may assume by symmetry that u is comparable with x . Together with $x \not\leq u$, this yields $u < x$. Suppose, with a view to contradiction, that $u \not\leq y$. As $y \not\leq u$, this implies that u and y are incomparable. Since $z \leq u, x, y$ and $u \leq x$, we can apply the three point rule to obtain $y \leq x$, a contradiction. Thus, $u \leq y$, and so $u \in \downarrow x \cap \downarrow y$. \square

Claim 6.16. *Let $A \in \mathcal{A}_\infty$ and $u \in X$. If $\perp_A \not\leq u$, then there is $x \in A$ such that $\downarrow u \cap \downarrow x = \emptyset$.*

Proof. Since $\perp_A \not\leq u$, the definition of the order relation on X^+ gives that

$$a \not\leq u \text{ for all } a \in A. \quad (15)$$

Since $\uparrow u$ is finite (as $u \in X$ and X is image-finite) and A is infinite, there is $y \in A$ such that $u \not\leq y$. Then $u \in X$ and $y \in A$ are incomparable by (15). Consider $x \in A$ such that $x < y$. Its existence follows from the fact that A is infinite and downward directed and $\uparrow y$ is finite (as $y \in A \subseteq X$ and X is image-finite). Since u and y are incomparable and $x < y$, we get $u \not\leq x$. In addition, $x \not\leq u$ by (15). Thus, x and u are also incomparable. To conclude the proof, suppose that there is $z \in \downarrow u \cap \downarrow x$. As $z \leq u, x, y$ and x, u are incomparable and $x \leq y$, we can apply the three point rule to obtain $u \leq y$, a contradiction. \square

Claim 6.17. *If $u \in X^+$ is nondistinguished, then $\downarrow u$ is clopen in X^+ .*

Proof. By Claim 6.14(ii), the set $\downarrow u$ is open in X^+ . Then it only remains to verify that $X^+ \setminus \downarrow u$ is also open. First, observe that Condition (T1) holds for $X^+ \setminus \downarrow u$ (just take $Z = \{u\}$). To prove Condition (T2), let $x \in X^+ \setminus \downarrow u$. Since $u \notin \uparrow x$ and $\downarrow x \setminus \downarrow u \subseteq X^+ \setminus \downarrow u$, we may take $Z = \{u\}$.

To prove Condition (T3), suppose that $X^+ \setminus \downarrow u$ contains an element of the form $\perp_{\{x,y\}}$. Since $\perp_{\{x,y\}} \in \downarrow x \cap \downarrow y$ and $\perp_{\{x,y\}} \not\leq u$, we may apply Claim 6.15 to obtain that either $\downarrow x \cap \downarrow y \cap \downarrow u = \emptyset$ or $u \in \downarrow x \cap \downarrow y$. If $\downarrow x \cap \downarrow y \cap \downarrow u = \emptyset$, then $\downarrow x \cap \downarrow y \subseteq X^+ \setminus \downarrow u$, while if $u \in \downarrow x \cap \downarrow y$, then we take $Z = \{u\}$. Thus, Condition (T3) holds in both cases.

It only remains to check (T4). Consider $A \in \mathcal{A}_\infty$ such that $\perp_A \in X^+ \setminus \downarrow u$. In particular, $\perp_A \not\leq u$. Since u is nondistinguished by assumption and $X^+ \setminus X$ is a set of distinguished elements, $u \in X$. Therefore, $u \in X$ and $\perp_A \not\leq u$. Thus, we may apply Claim 6.16 to obtain $x \in A$ such that $\downarrow x \subseteq X^+ \setminus \downarrow u$. \square

Claim 6.18. *If u and v are incomparable elements in X , then the set $\downarrow u \cap \downarrow v$ is clopen in X^+ .*

Proof. Define $U = X^+ \setminus (\downarrow u \cap \downarrow v)$. Since u and v are incomparable,

$$\downarrow u \cap \downarrow v = (\downarrow u \setminus \{u\}) \cap (\downarrow v \setminus \{v\}).$$

Therefore, in view of Claim 6.14, the set $\downarrow u \cap \downarrow v$ is open in X^+ .

It remains to show that U is also open. First, observe that $X^+ \setminus \downarrow\{u, v\} \subseteq U$, whence U satisfies (T1). To prove (T2), let $x \in U$. By definition of U , either $x \not\leq u$ or $x \not\leq v$. By symmetry, we may assume that $x \not\leq u$. Together with the obvious inclusion $\downarrow x \setminus \downarrow u \subseteq U$, this implies that Condition (T2) holds for U . To prove (T3), suppose that U contains an element of the form $\perp_{\{x,y\}}$. Then either $\perp_{\{x,y\}} \not\leq u$ or $\perp_{\{x,y\}} \not\leq v$. By symmetry we may assume that $\perp_{\{x,y\}} \not\leq u$, whence $\perp_{\{x,y\}} \not\leq u$. Now apply Claim 6.15 to obtain that either $\downarrow x \cap \downarrow y \cap \downarrow u = \emptyset$ or $u \in \downarrow x \cap \downarrow y$. Thus, Condition (T3) follows from the inclusion $\downarrow x \cap \downarrow y \subseteq U$ in the first case and from the inclusion $(\downarrow x \cap \downarrow y) \setminus \downarrow u \subseteq U$ in the second case.

Lastly, to prove (T4), consider $A \in \mathcal{A}_\infty$ such that $\perp_A \in U$. By definition of U , either $\perp_A \not\leq u$ or $\perp_A \not\leq v$. By symmetry, we may assume that $\perp_A \not\leq u$, so $\perp_A \not\leq u$. Moreover, $u \in X$ by assumption. Thus, we may apply Claim 6.16 to obtain $x \in A$ such that $\downarrow u \cap \downarrow x = \emptyset$. Consequently, also $\downarrow u \cap \downarrow v \cap \downarrow x = \emptyset$, and hence $\downarrow x \subseteq X^+ \setminus (\downarrow u \cap \downarrow v) = U$. \square

Claim 6.19. *The space X^+ satisfies the Priestley separation axiom.*

Proof. Let $u, v \in X^+$ with $u \not\leq v$. We must find a clopen downset D of X^+ such that $u \notin D$ and $v \in D$. If v is nondistinguished, we may take $D = \downarrow v$ by Claim 6.17. Suppose v is distinguished. We have the following cases:

- (i) $v = \top$;
- (ii) $v = \perp_{\{x,y\}}$ for some $x, y \in X$;
- (iii) $v = \perp_A$ for some $A \in \mathcal{A}_\infty$.

(i): Since X is image-finite, there is a maximal element u° in X^+ such that $u \leq u^\circ$. Notice that $u^\circ \neq \top$ as $u \not\leq v = \top$. Since \top is maximal, this implies $v = \top \notin \downarrow u^\circ$. Being maximal in X^+ and different from \top , the element u° is nondistinguished. Thus, $\downarrow u^\circ$ is clopen by Claim 6.17. Since u° is maximal, we can apply Proposition 4.2 and obtain that the downset $\downarrow u^\circ$ is also an upset of X^+ . Therefore, the set $D := X^+ \setminus \downarrow u^\circ$ is a clopen downset of X^+ such that $u \notin D$ and $v \in D$.

(ii): If $u \notin \downarrow x \cap \downarrow y$, then we may take $D = \downarrow x \cap \downarrow y$ by Claim 6.18. Suppose $u \in \downarrow x \cap \downarrow y$. By Proposition 4.2, for every $z \in \downarrow x \cap \downarrow y$ there is a unique immediate predecessor z° of x and y such that $z \leq z^\circ$. In particular, u° is an immediate predecessor of x and y . Define

$$D = (\downarrow x \cap \downarrow y) \setminus \downarrow u^\circ.$$

Clearly $u \notin D$. Moreover, since u° and v are immediate predecessors of x and y , from $v \leq u^\circ$ it follows that $v = u^\circ$. This contradicts the fact that $u \leq u^\circ$ and $u \not\leq v$. Therefore, $v \not\leq u^\circ$, and hence $v = \perp_{\{x,y\}} \in D$.

We show that D is a clopen downset of X^+ . As X^+ has width at most two, its elements may have at most two immediate successors. It follows that among immediate predecessors of x and y there exists just one distinguished element, namely $\perp_{\{x,y\}}$. Since $u \not\leq v = \perp_{\{x,y\}}$, the element u° is nondistinguished. Thus, by Claims 6.17 and 6.18, the set D is clopen. Finally, to prove that D is a downset, we show that

$$z \notin D \iff z^\circ = u^\circ \text{ for every } z \in \downarrow x \cap \downarrow y. \quad (16)$$

Consider $z \in \downarrow x \cap \downarrow y$. If $z^\circ = u^\circ$, then $z \in \downarrow u^\circ$ (since $z \leq z^\circ$), whence $z \notin D$. Conversely, if $z \notin D$, then $z \in \downarrow u^\circ$. Therefore, u° is the unique immediate predecessor of x and y above z . Consequently, $z^\circ = u^\circ$. This establishes (16). Lastly, consider $z \in D$ and $t \leq z$. Since $t \leq z \leq x, y$, we get $t^\circ = z^\circ$. Moreover, (16) and $z \in D$ yield $z^\circ \neq u^\circ$. Thus, also $t^\circ \neq u^\circ$. With another application of (16) we conclude that $t \in D$. Hence, D is a downset.

(iii): Since $u \neq v$, we have $u \neq \perp_A$. Because \perp_A is the unique element whose set of strict successors is A , from $u \neq \perp_A$ it follows that there is $z \in A$ with $u \not\leq z$. If $z = \top$ or $z = \perp_{\{x,y\}}$ for some $x, y \in X$, then we take the clopen downset D constructed in the previous cases. Otherwise, z is nondistinguished and we may take $D = \downarrow z$ by Claim 6.17. \square

In view of Claims 6.5, 6.12, and 6.19, we conclude that X^+ is an Esakia space. \square

Corollary 6.20. *Let X be an image-finite root system. Then there is an Esakia space X^+ whose underlying poset is also a root system such that X is the image-finite part of X^+ .*

Proof. If X is a root system, the poset X^+ constructed in the proof of Proposition 6.2 is also a root system. Therefore, the result follows from Proposition 6.2. \square

Corollary 6.21. *Let n be a positive integer and X a diamond system of depth $\leq n$. Then X is Esakia representable.*

Proof. If X is of depth $\leq n$, the poset X^+ constructed in the proof of Proposition 6.2 coincides with X . Consequently, X is Esakia representable by Proposition 6.2. \square

7. CONSEQUENCES

By Theorems 4.4 and 6.1, a variety of Heyting algebras is such that its profinite members are profinite completions if and only if it omits the finite algebras $\text{Up}(P_1)$, $\text{Up}(P_2)$, $\text{Up}(P_3)$, and $\text{Up}(P_4)$. Consequently, we obtain:

Theorem 7.1.

- (i) *The problem of determining whether a finite set of equations axiomatizes a variety of Heyting algebras whose profinite members are profinite completions is decidable.*
- (ii) *The problem of determining whether a finite set of finite Heyting algebras generates a variety whose profinite members are profinite completions is decidable.*

Proof. (i): Let Σ be a finite set of equations and \mathbb{V} the variety axiomatized by it. By Theorems 4.4 and 6.1, the profinite members of \mathbb{V} are profinite completions if and only if \mathbb{V} validates the Jankov formulas $\mathcal{J}(P_1)$, $\mathcal{J}(P_2)$, $\mathcal{J}(P_3)$, and $\mathcal{J}(P_4)$. By Jankov's Lemma, this happens precisely when \mathbb{V} contains none of $\text{Up}(P_1)$, $\text{Up}(P_2)$, $\text{Up}(P_3)$, and $\text{Up}(P_4)$. This condition, in turn, can be decided by checking whether the equations in Σ hold in any of the finite algebras $\text{Up}(P_1)$, $\text{Up}(P_2)$, $\text{Up}(P_3)$, and $\text{Up}(P_4)$.


(ii): The variety generated by a finite set K of finite Heyting algebras is such that its profinite members are profinite completions precisely when K validates $\mathcal{J}(P_1)$, $\mathcal{J}(P_2)$, $\mathcal{J}(P_3)$, and $\mathcal{J}(P_4)$. Since K is a finite set of finite algebras, the latter property is decidable. \square

We next provide some properties of diamond Heyting algebras. Recall that a variety is *finitely based* if it can be axiomatized by finitely many equations, and that a class of (similar) algebras is a *quasivariety* if it is closed under the formation of reduced products, subalgebras, and isomorphic copies, see, e.g., [34, Sec. 1.5]. A variety is called *primitive* if all its subquasivarieties are varieties [34, Sec. 5.1.4].

Theorem 7.2. *Varieties of diamond Heyting algebras are locally finite, primitive, and finitely based. Moreover, there are only countably many of them.*

Proof. That varieties of diamond Heyting algebras are locally finite follows from Theorem 3.3 and the fact that $\text{DHA} \subseteq \text{CHA}$. In [19, 20] it is proved that a variety of Heyting algebras is primitive if and only if it excludes the algebras of upsets of the rooted posets in Figure 12. (The original proof in [20] is in Russian. For an English version of the proof, based on Esakia duality, see [13].)

From the fact that varieties of diamond Heyting algebras omit $\text{Up}(P_1)$, $\text{Up}(P_2)$, $\text{Up}(P_3)$, and $\text{Up}(P_4)$, it easily follows that they also omit the algebras of upsets of the posets P_1 , P_2 , P_5 , F_3 , and P_7 . Consequently,

FIGURE 12. The posets P_1 , P_2 , P_5 , F_3 and P_7 .

varieties of diamond Heyting algebras are primitive. As it is known that there are only countably many primitive varieties of Heyting algebras and that all of them are finitely based (see for instance [13, Thm. 9.1]), we conclude that the same holds for varieties of diamond Heyting algebras. It only remains to prove that there are infinitely many varieties of diamond Heyting algebras. But this follows, for instance, from the fact that there are infinitely many varieties of Gödel algebras, which are all varieties of diamond Heyting algebras. \square

A variety V of Heyting algebras is said to be *canonical* provided that $\text{Up}(X_A) \in V$ for all $A \in V$. The terminology comes from the fact that, given a Heyting algebra A , the completion $\text{Up}(X_A)$ is called the *canonical extension* of A [32]. Furthermore, recall that the smallest variety containing a class of similar algebras V is $\text{HISP}(V)$ [17, Thm. II.9.5].

Theorem 7.3. *The variety DHA is canonical.*

Proof. It is well known that if \mathcal{P} is an elementary class of posets, then the variety of Heyting algebras generated by $\{\text{Up}(X) : X \in \mathcal{P}\}$ is canonical (see, e.g., [33, Thm. 7.1]). Let then \mathcal{P} be the class of diamond systems. Notice that \mathcal{P} is elementary because the conditions in Definition 4.1 can be expressed by first-order sentences, so the variety

$$V := \text{HISP}\{\text{Up}(X) : X \in \mathcal{P}\}$$

is canonical. Thus, to conclude the proof, it suffices to show that $V = \text{DHA}$.

A straightforward adaptation of the proof of the implication (iii) \Rightarrow (i) in Theorem 4.4 shows that $\text{Up}(X) \in \text{DHA}$ for all $X \in \mathcal{P}$. Consequently, $V \subseteq \text{DHA}$. To prove the other inclusion, consider $A \in \text{DHA}$. From Theorem 4.4 it follows that X_A is a diamond system, whence $X_A \in \mathcal{P}$. Together with the fact that A embeds into $\text{Up}(X_A)$ via the map γ_A , this implies $A \in V$, as desired. \square

Intermediate logics algebraized, in the sense of [14], by varieties of diamond Heyting algebras have interesting metalogical properties. First, an intermediate logic is said to be *hereditarily structurally complete* if all its finitary extensions [31] are *structurally complete* in the sense that their admissible rules are derivable (see for instance [60]). On the other hand, an intermediate logic L has the *infinite Beth definability property* if implicit definitions can be turned explicit in L (we refer to [16, 52] for the technical details).

Theorem 7.4. *Intermediate logics algebraized by varieties of diamond algebras are hereditarily structurally complete and have the infinite Beth definability property.*

Proof. As detailed in [60] and [16], an intermediate logic is hereditarily structurally complete (resp. has the infinite Beth definability property) precisely when it is algebraized by a variety of Heyting algebras that is primitive (resp. in which epimorphisms are surjective). The fact that varieties of diamond Heyting algebras are primitive follows from Theorem 7.2, while the fact that epimorphisms are surjective in them is a consequence of [53, Thm. 9.4] (see also [11]). \square

We close this paper by drawing connection to the classical Representation Problem mentioned in the introduction.

Definition 7.5. We call a variety \mathcal{V} of Heyting algebras *representable* if X is Esakia representable for every poset X such that $\text{Up}(X) \in \mathcal{V}$.

Recall that D_n is the variety of Heyting algebras of depth $\leq n$. Our aim is to prove the following:

Theorem 7.6. *A variety \mathcal{V} of Heyting algebras is representable if and only if $\mathcal{V} \subseteq D_n \cap \text{DHA}$ for some positive integer n .*

To this end, we rely on the following folklore observation.

Proposition 7.7. *For a variety \mathcal{V} of Heyting algebras, the following are equivalent :*

- (i) *the members of \mathcal{V} have depth $\leq n$ for some positive integer n ;*
- (ii) *\mathcal{V} omits all infinite chains;*
- (iii) *\mathcal{V} omits some chain;*
- (iv) *\mathcal{V} validates $\mathcal{J}(C)$ for some finite chain C .*

Proof of Theorem 7.6. Let Z^- be the poset of negative integers with the standard order. First suppose that \mathcal{V} is representable. By Proposition 5.2 and Lemma 5.3, \mathcal{V} omits the algebras $\text{Up}(P)$ for all $P \in \{P_1, P_2, P_3, P_4\}$. In view of Theorem 4.4, we conclude that $\mathcal{V} \subseteq \text{DHA}$. Moreover, $\text{Up}(Z^-) \notin \mathcal{V}$ as Z^- is not Esakia representable by Proposition 2.3(ii). Thus, by Proposition 7.7 the members of \mathcal{V} have depth $\leq n$ for some positive integer n .

Conversely, suppose that $\mathcal{V} \subseteq D_n \cap \text{DHA}$ for some positive integer n . Consider a poset X such that $\text{Up}(X) \in \mathcal{V}$. We show that X is image-finite. It suffices to prove that X has bounded depth and width. Since $\text{Up}(X)$ has depth $\leq n$, the same holds for X (see, e.g., [18, Prop. 2.38]). Therefore, by Theorem 4.4, DHA has width ≤ 2 . Thus, $\text{Up}(X)$ also has width ≤ 2 , and hence so does X (see, e.g., [18, Prop. 2.39]). Consequently, X is image-finite. As X is image-finite and $\text{Up}(X)$ is a diamond Heyting algebra, by Corollary 4.8 we obtain that X is a diamond system of depth $\leq n$. Thus, X is Esakia representable by Corollary 6.21. \square

Acknowledgements. The third author was partially supported by the Beatriz Galindo grant BEAGAL-18/00040 and by the I+D+i research project PID2019-110843GA-I00, both funded by the Ministry of Science and Innovation and the Ministry of Universities of Spain. The fourth author would like to express his deep gratitude to the logic group in the Institute of Computer Science of the Czech Academy of Sciences, especially to Petr Cintula, for the hospitality and support that allowed him to conduct this research.

REFERENCES

- [1] S. Abramsky. A Cook’s Tour of the Finitary Non-Well-Founded Sets. *Invited Lecture at BCTCS*, 1988. Available at arXiv:1111.7148.
- [2] J. Adámek, H. Herrlich, and G. E. Strecker. *Abstract and concrete categories. The joy of cats*. Online edition, 2004. Reprints in Theory and Applications of Categories, No. 17 (2006), 1–507.
- [3] R. Balbes. On the partially ordered set of prime ideals of a distributive lattice. *Canadian Journal of Mathematics*, 23:866–874, 1971.
- [4] R. Balbes and P. Dwinger. *Distributive lattices*. University of Missouri Press, Columbia, Missouri, 1974.
- [5] C. Bergman. *Universal Algebra: Fundamentals and Selected Topics*. Chapman & Hall Pure and Applied Mathematics. Chapman and Hall/CRC, 2011.
- [6] G. Bezhanishvili. Varieties of monadic Heyting algebras. III. *Studia Logica*, 64(2):215–256, 2000.
- [7] G. Bezhanishvili and N. Bezhanishvili. Profinite Heyting algebras. *Order*, 25(3):211–227, 2008.
- [8] G. Bezhanishvili, M. Gehrke, R. Mines, and P. J. Morandi. Profinite completions and canonical extensions of Heyting algebras. *Order*, 23(2–3):143–161, 2006.
- [9] G. Bezhanishvili and J. Harding. Compact Hausdorff Heyting algebras. *Algebra universalis*, 76:301–304, 2016.
- [10] G. Bezhanishvili and P. J. Morandi. Profinite Heyting algebras and profinite completions of Heyting algebras. *Georgian Mathematical Journal*, 16(1):29–47, 2009.
- [11] G. Bezhanishvili, T. Moraschini, and J. G. Raftery. Epimorphisms in varieties of residuated structures. *Journal of Algebra*, 492:185–211, 2017.
- [12] N. Bezhanishvili. *Lattices of Intermediate and Cylindric Modal Logics*. Ph. D. Dissertation DS-2006-02, Institute for Logic, Language and Computation, University of Amsterdam, 2006.

- [13] N. Bezhanishvili and T. Moraschini. Citkin’s description of hereditarily structurally complete intermediate logics via Esakia duality. *Submitted manuscript*, 2020.
- [14] W. J. Blok and D. Pigozzi. *Algebraizable logics*, volume 396 of *Mem. Amer. Math. Soc.* A.M.S., Providence, January 1989.
- [15] P. Blackburn, M. de Rijke, and Y. Venema. *Modal logic*. Number 53 in Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, Cambridge, 2001.
- [16] W. J. Blok and E. Hoogland. The Beth property in Algebraic Logic. *Studia Logica*, 83(1–3):49–90, 2006.
- [17] S. Burris and H. P. Sankappanavar. *A course in Universal Algebra*. Available in internet <https://www.math.uwaterloo.ca/~snburris/htdocs/ualg.html>, the millennium edition, 2012.
- [18] A. Chagrov and M. Zakharyashev. *Modal Logic*, volume 35 of *Oxford Logic Guides*. Oxford University Press, 1997.
- [19] A. Citkin. On structurally complete superintuitionistic logics. *Soviet Mathematics Doklady*, 19:816–819, 1978.
- [20] A. Citkin. Structurally complete superintuitionistic logics and primitive varieties of pseudoBoolean algebras. *Mat. Issled. Neklass. Logiki*, 98, 134–151, 1987 (in Russian).
- [21] D. M. Clark, B. A. Davey, and M. G. Jackson. The axiomatizability of topological prevarieties. *Advances in Mathematics*, 218(5):1604–1653, 2008.
- [22] W. Cornish. On H. Priestley’s dual of the category of bounded distributive lattices. *Mat. Vesnik*, 12(27)(4):329–332, 1975.
- [23] D. van Dalen. Intuitionistic logic. In *Handbook of Philosophical Logic. Volume III*, 225–339, Springer, 1986.
- [24] D. van Dalen. *Logic and structure*. Universitext. Springer-Verlag, Berlin, fourth edition, 2004.
- [25] M. Dummett. A propositional calculus with denumerable matrix. *The Journal of Symbolic Logic*, 24:97–106, 1959.
- [26] L. Esakia. Topological Kripke models. *Soviet Math. Dokl.*, 15:147–151, 1974.
- [27] L. Esakia. On a locally finite variety of Heyting algebras. In *XVII Soviet Algebraic Conference. Part II, Minsk*, 280–281, 1983.
- [28] L. Esakia. *Heyting Algebras. Duality Theory*. Springer, English translation of the original 1985 book. 2019.
- [29] L. Esakia and R. Grigolia. The criterion of Brouwerian and closure algebras to be finitely generated. *Bulletin of the Section of Logic*, 6(2):46–52, 1977.
- [30] K. Fine. Logics containing K4. I. *J. Symbolic Logic*, 39:31–42, 1974.
- [31] J. M. Font. *Abstract Algebraic Logic - An Introductory Textbook*, volume 60 of *Studies in Logic - Mathematical Logic and Foundations*. College Publications, London, 2016.
- [32] M. Gehrke and B. Jónsson. Bounded distributive lattices with operators. *Mathematica Japonica*, 40:207–215, 1994.
- [33] R. Goldblatt. Canonical extensions and ultraproducts of polarities *Algebra Universis*, 79:1–28, 2018.
- [34] V. A. Gorbunov. *Algebraic theory of quasivarieties*. Siberian School of Algebra and Logic. Consultants Bureau, New York, 1998. Translated from Russian.
- [35] G. Grätzer. *Lattice Theory. First Concepts and Distributive Lattices*. W. H. Freeman and Company, San Francisco, 1971.
- [36] P. Hájek. Metamathematics of fuzzy logic. *Trends in Logic—Studia Logica Library*. Kluwer Academic Publishers, Dordrecht, 1998.
- [37] M. Hochster Prime ideal structure in commutative rings. *Trans. Amer. Math. Soc.*, 142:43–60, 1969.
- [38] T. Hosoi. On intermediate logics I. *J. Fac. Sci. Univ. Tokyo*, 14:293–293–312, 1967.
- [39] T. Hosoi. Axiomatization of certain intermediate logics. *Journal of Tsuda, College* 10:9–12, 1978.
- [40] V. A. Jankov On the relation between deducibility in intuitionistic propositional calculus and finite implicative structures. *Doklady Akademii Nauk SSSR*, 151:1293–1294, 1963.
- [41] V. A. Jankov The construction of a sequence of strongly independent superintuitionistic propositional calculi *Soviet Mathematics Doklady*, 9:806–807, 1968.
- [42] V. A. Jankov Conjunctively irresolvable formulae in propositional calculi. *Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya*, 33:18–38, 1969.
- [43] P. T. Johnstone. *Stone Spaces*, volume 3 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 1982.
- [44] A. Joyal. Spectral spaces and distributive lattices. *Notices Amer. Math. Soc.*, 18:393–394, 1971.
- [45] I. Kaplansky. *Commutative rings*. Revised edition. The University of Chicago Press, Chicago, Ill.-London, 1974.
- [46] C. Kupke, A. Kurz, and Y. Venema. Stone coalgebras. *Theoretical Computer Science*, 327(1-2):109–134, 2004.
- [47] A. V. Kuznetsov On superintuitionistic logics. *Proceedings of the International Congress of Mathematicians, Vancouver*, 1:243–249, 1974.
- [48] W. J. Lewis. The spectrum of a ring as a partially ordered set. *J. Algebra*, 25:419–435, 1973.
- [49] L. L. Maksimova. Pretabular intuitionist logics. *Algebra and Logic*, 11:308–314–308–314, 1972.
- [50] L. L. Maksimova. On locally finite varieties of pseudoboolean algebras. *XVI All-Union Algebraic Conference*, Leningrad, 1:99–100, 1981.
- [51] S. I. Mardaev. Two Sequences of Locally Tabular Superintuitionistic Logics. *Studia Logica*, 50(2):333–342, 1991.
- [52] T. Moraschini, J. G. Raftery, and J. J. Wannenburg. Epimorphisms, definability and cardinalities. *Studia Logica*, 108:255–275, 2020.
- [53] T. Moraschini and J. J. Wannenburg. Epimorphisms surjectivity in varieties of Heyting algebras. *Annals of Pure and Applied Logic*, 171(9), 2020.
- [54] A. Di Nola and R. Grigolia. Gödel spaces and perfect MV-algebras. *Journal of Applied Logic*, 13(3):270–284, 2015.
- [55] H. Ono. Kripke models and intermediate logics. *Publ. RIMS Kyoto Univ.*, 6:461–476, 1970/71.
- [56] H. A. Priestley. Representation of distributive lattices by means of ordered Stone spaces. *Bull. London Math. Soc.*, 2:186–190, 1970.

- [57] H. A. Priestley. Ordered topological spaces and the representation of distributive lattices. *Proceedings of the London Mathematical Society. Third Series*, 24:507–530, 1972.
- [58] H. A. Priestley. Spectral sets. *Journal of Pure and Applied Algebra*, 94(1):101–114, 1994.
- [59] H. Rasiowa and R. Sikorski *The mathematics of metamathematics*. Monografie Matematyczne, Tom 41. Państwowe Wydawnictwo Naukowe, Warsaw, 1963.
- [60] V. V. Rybakov. *Admissibility of logical inference rules*, volume 136 of *Studies in Logic*. Elsevier, Amsterdam, etc., 1997.
- [61] F. M. Schneider and J. Zúbrägel. Profinite algebras and affine boundedness. *Adv. Math.*, 305:661–681, 2017.
- [62] Z. Semadeni. *Banach spaces of continuous functions*, volume 1 of *Monografie Matematyczne, Tom 55*. PWN—Polish Scientific Publishers, Warsaw, 1971.
- [63] S. K. Sobolev. Finite-dimensional superintuitionistic logics. *Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya*, 41(5):963–986, 1977.
- [64] T. P. Speed. Profinite posets. *Bull. Austral. Math. Soc.*, 6:177–183, 1972.
- [65] M. H. Stone. The theory of representations for Boolean algebras. *Transactions of the American Mathematical Society*, 39:37–111, 1936.
- [66] M. H. Stone. Applications of the theory of Boolean rings to general topology. *Transactions of the American Mathematical Society*, 41(3):375–481, 1937.

GURAM BEZHANISHVILI: DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVERSITY, LAS CRUCES NM 88003, USA

Email address: guram@nmsu.edu

NICK BEZHANISHVILI: INSTITUTE FOR LOGIC, LANGUAGE AND COMPUTATION, UNIVERSITY OF AMSTERDAM, POSTBUS 94242, 1090GE AMSTERDAM, THE NETHERLANDS

Email address: N.Bezhanishvili@uva.nl

TOMMASO MORASCHINI: DEPARTMENT OF PHILOSOPHY, UNIVERSITY OF BARCELONA, CARRER DE MONTALEGRE 6, 08001, BARCELONA, SPAIN

Email address: tommaso.moraschini@ub.edu

MICHAŁ STRONKOWSKI: INSTITUTE OF COMPUTER SCIENCE, ACADEMY OF SCIENCES OF CZECH REPUBLIC, POD VODÁRENSKOU VĚŽÍ 271/2, 182 07 PRAGUE 8, CZECH REPUBLIC
and

FACULTY OF MATHEMATICS AND INFORMATION SCIENCES, WARSAW UNIVERSITY OF TECHNOLOGY, UL. KOSZYKOWA 75, 00-662 WARSAW, POLAND

Email address: m.stronkowski@mini.pw.edu.pl